

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 October 2006 (26.10.2006)

PCT

(10) International Publication Number
WO 2006/113331 A1

(51) International Patent Classification:

C07K 14/715 (2006.01) *C07K 16/28* (2006.01)
A61P 17/02 (2006.01) *C07K 16/24* (2006.01)
A61K 38/17 (2006.01) *A61K 39/395* (2006.01)

(21) International Application Number:

PCT/US2006/013829

(22) International Filing Date: 13 April 2006 (13.04.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/671,374 14 April 2005 (14.04.2005) US

(71) Applicants (for all designated States except US):

WYETH [US/US]; Five Giralda Farms, Madison, New Jersey 07940 (US). **THE GOVERNMENT OF THE UNITED STATES AS REPRESENTED BY THE SECRETARY OF THE DEPARTMENT OF HEALTH AND HUMAN SERVICES** [US/US]; National Institutes of Health, Office of Technology Transfer, 6011 Executive Boulevard, Suite 325, Rockville, Maryland 20852-3804 (US). **THE PRESIDENT AND FELLOWS OF HARVARD COLLEGE** [US/US]; Holyoke Center, Suite 727, 1350 Massachusetts Ave., Cambridge, Massachusetts 02138 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **YOUNG, Deborah, A.** [US/US]; 39 Nelson Road, Melrose, Massachusetts 02176 (US). **WYNN, Thomas, A.** [US/US]; 1822 Tilton Drive, Silver Spring, Maryland 20902 (US). **COLLINS, Mary** [US/US]; 54 Rathbun Road, Natick, Massachusetts 01760 (US). **GRUSBY, Michael** [US/US]; 5 Nightingale Path, Newton, Massachusetts 02459 (US).

(74) Agents: **MANDRA, Raymond, R.** et al.; Fitzpatrick, Cella, Harper & Scinto, 30 Rockefeller Plaza, New York, New York 10112-3801 (US).

(81) Designated States (unless otherwise indicated, for every


kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),

[Continued on next page]

(54) Title: METHODS FOR TREATING AND PREVENTING FIBROSIS BY IL-21 / IL-21R ANTAGONISTS



(57) Abstract: The present invention provides methods of screening for compositions useful for treating, ameliorating, or preventing fibrosis and/or fibrosis-associated conditions by measuring changes in the level(s) of IL-21 and/or IL-21 receptor (IL-21R) (e.g., the level of expression of IL-21 and/or IL-21 R protein and/or mRNA, the level of activity of IL-21 and/or IL-21R, the level of interaction of IL-21 with IL-21R). The invention further provides antagonists of IL-21 or IL-21R for the treatment of fibrosis and/or fibrosis-associated conditions. Further provided herein are methods of diagnosing, prognosing, and monitoring the progress (e.g., the course of treatment) of fibrosis and/or fibrosis-associated conditions by measuring the level of IL-21 and/or IL-21R (i.e., the level of activity of IL-21 and/or IL-21R, the level of expression of IL-21 and/or IL-21R (e.g., the level of IL-21 and/or IL-21 R gene products), and/or the level of interaction of IL-21 with IL-21R).

WO 2006/113331 A1



European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report*

— *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

TITLE

METHODS FOR TREATING AND PREVENTING FIBROSIS

BY IL-21 / IL-21R ANTAGONISTS

Related Applications

[0001] This application claims the benefit of priority to U.S. Provisional Patent Application No. 60/671,374, filed April 14, 2005, which is hereby incorporated by reference herein in its entirety.

BACKGROUND OF THE INVENTION

Field of the Invention

[0002] The present invention relates to methods for treating and preventing fibrosis and fibrosis-associated conditions.

Related Background Art

[0003] Injury to any organ typically leads to a physiological response involving platelet-induced hemostasis, followed by an influx of inflammatory cells and activated fibroblasts. Cytokines produced by these cell types drive the formation of new extracellular matrix and blood vessels, which collectively form granulation tissue. The formation of fibrous tissue is part of the normal beneficial process of healing following injury; fibrosis, however, is a condition characterized by an abnormal accumulation of a collagen matrix following injury or inflammation that alters the structure and function of various tissues.

Progressive fibrosis in the kidney, liver, lung, heart, bone, bone marrow, and skin is a major cause of, or contributor to, death.

[0004] Many of the diseases associated with the proliferation of fibrous tissue are chronic and often debilitating and include, for example, skin diseases such as scleroderma. Some, including pulmonary fibrosis, may be fatal, due in part to the fact that the current treatments have significant side effects and are generally not effective in slowing or halting the progression of fibrosis. There is, accordingly, a continuing need for new anti-fibrotic agents.

[0005] The IL-21 receptor (IL-21R) is a newly discovered member of the class I cytokine receptor family (Parrish-Novak et al. (2000) *Nature* 408:57-63; Ozaki et al. (2000) *Proc. Natl. Acad. Sci. U.S.A.* 97:11439-44). The IL-21 receptor shows significant sequence and structural homology with the IL-4 receptor alpha (IL-4R α) chain and is adjacent to the IL-4R α in the human and mouse genomes, while its ligand, IL-21, shares significant homology with the cytokines IL-2, IL-4, and IL-15 (Sivakumar et al. (2004) *Immunology* 112:177-82; Habib et al. (2003) *J. Allergy Clin. Immunol.* 112:1033-45). IL-21 and IL-21R are thus newly described members of the gamma chain (γ c)-dependent cytokine network because of their homology with cytokines and receptors that require the γ c for functional signaling (Vosshenrich and Di Santo (2001) *Curr. Biol.* 11:R175-77). Because all members of the γ c network exhibit important and unique roles in host immunity, there has been growing interest in dissecting the novel functions of the IL-21R during antigen-triggered immune responses *in vivo*.

[0006] Initial studies examining the function of IL-21 showed that NK cell expansion is antagonized, whereas antigen-specific T cell immunity is promoted by IL-21, including anti-tumor immunity (Ma et al. (2003) *J. Immunol.* 171:608-15; Kishida et al. (2003) *Mol. Ther.* 8:552-58; Di Carlo et al. (2004) *J. Immunol.* 172:1540-47), findings that suggest that IL-21 serves as a bridge between innate and adaptive immune responses (Collins et al. (2003) *Immunol. Res.* 28:131-40). IL-21 also regulates B cell and CD8⁺ T cell function *in vivo* (Ozaki et al. (2002) *Science* 298:1630-34; Suto et al. (2002) *Blood* 100:4565-73; Mehta et al. (2003) *J. Immunol.* 170:4111-18; Pene et al. (2004) *J. Immunol.* 172:5154-57; Jin et al. (2004) *J. Immunol.* 173:657-65; Zeng et al. (2005) *J. Exp. Med.* 201:139-48). Additional studies suggest that IL-21 is a T_H2 cytokine that can inhibit the

differentiation of naïve T_H cells into IFN- γ -secreting T_H1 cells (Wurster et al. (2002) *J. Exp. Med.* 196:969-77). Indeed, exogenous treatment with IL-21 potently inhibited IFN- γ production without affecting other T_H1/T_H2-associated cytokines, suggesting that the repression of IFN- γ by IL-21 is highly specific. Thus, by virtue of its ability to suppress the development of T_H1 cells, it was hypothesized that IL-21 might promote T_H2 responses (Wurster et al., *supra*). Nevertheless, the involvement of the IL-21R signaling pathway in T_H2 response development was not previously investigated in any T_H2-dependent disorder.

[0007] In schistosomiasis, T_H2 cytokines play an indispensable role in the pathogenesis of the disease (Wynn (2004) *Nat. Rev. Immunol.* 4:583-594; Pearce and MacDonald (2002) *Nat. Rev. Immunol.* 2:499-511). Indeed, IL-4/IL-13-, IL-4R α -, and Stat6-deficient mice all show significantly impaired granuloma formation and liver fibrosis following infection with *S. mansoni* (Chiaramonte et al. (1999) *J. Clin. Invest.* 104:777-85; Kaplan et al. (1998) *J. Immunol.* 160:1850-56; Jankovic et al. (1999) *J. Immunol.* 163:337-42; Fallon et al. (2000) *J. Immunol.* 164:2585-91). Given the recent classification of IL-21 as a T_H2 cytokine (Wurster et al. (2002), *supra*; Mehta et al. (2005). *Proc. Natl. Acad. Sci. U.S.A.* 102:2016-21), the striking similarities between the IL-4 and IL-21 receptors (Sivakumar et al., *supra*; Habib et al., *supra*), and the critical role of the related IL-4R α /Stat6-signaling pathway in this disease as well as in other T_H2 cytokine-driven inflammatory disorders (Wynn (2003) *Annu. Rev. Immunol.* 21:425-56), an important question evolving from these studies was whether IL-21R signaling plays a significant role in the initiation and/or maintenance of T_H2 immunity.

SUMMARY OF THE INVENTION

[0008] The present invention provides methods of treating, ameliorating or preventing fibrosis or a fibrosis-associated disorder, as well as methods of screening for compounds and compositions useful in those methods. The invention also provides methods of diagnosing, prognosing, and monitoring the progress (e.g., the course of treatment) of fibrosis and/or fibrosis-associated conditions. These methods are related to measuring and/or modulating the level of IL-21 and/or IL-21R (i.e., the level of activity of IL-21 and/or IL-21R, the

level of expression of IL-21 and/or IL-21R (e.g., the level of IL-21 and/or IL-21R gene products), and/or the level of interaction of IL-21 with IL-21R). The invention further provides antagonists of IL-21 or IL-21R for the treatment of fibrosis and/or fibrosis-associated conditions.

[0009] In one embodiment, the invention provides a method for treating, ameliorating, or preventing fibrosis or a fibrosis-associated disorder in a subject (e.g., a human) comprising administering to the subject a therapeutically effective amount of an agent that reduces the level of IL-21 and/or IL-21R in the subject. In a further embodiment, the agent is an IL-21/IL-21R antagonist selected from the group consisting of an anti-IL-21R antibody, an anti-IL-21 antibody, an antigen-binding fragment of an anti-IL-21R antibody, an antigen-binding fragment of an anti-IL-21 antibody, and a soluble fragment of an IL-21R. In another further embodiment, the agent is a soluble fragment of an IL-21R, and the soluble fragment of the IL-21R comprises an amino acid sequence that is at least 90% identical to an amino acid sequence selected from the group consisting of amino acids 1-538 of SEQ ID NO:2, amino acids 20-538 of SEQ ID NO:2, amino acids 1-235 of SEQ ID NO:2, amino acids 20-235 of SEQ ID NO:2, amino acids 1-236 of SEQ ID NO: 2, amino acids 20-236 of SEQ ID NO:2, amino acids 1-529 of SEQ ID NO:5, amino acids 20-529 of SEQ ID NO:5, amino acids 1-236 of SEQ ID NO:5, and amino acid 20-236 of SEQ ID NO:5. In another embodiment, the soluble fragment of the IL-21R binds to an IL-21 polypeptide.

[0010] In another embodiment, the agent is a soluble fragment of an IL-21R, and the soluble fragment of the IL-21R comprises an amino acid sequence that is substantially identical to the amino acid sequence set forth in SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, or SEQ ID NO:27. In another embodiment, the amino acid sequence of the soluble fragment of the IL-21R comprises an amino acid sequence that is substantially identical to the amino acid sequence set forth in SEQ ID NO:11 or SEQ ID NO:13. In another embodiment, the agent is a soluble fragment of an IL-21R, and the soluble fragment of the IL-21R is encoded by a nucleotide sequence that is substantially identical to the nucleic acid sequence set forth in SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, or SEQ ID NO:26. In another embodiment, the soluble fragment of

the IL-21R is encoded by a nucleotide sequence that is substantially identical to the nucleic acid sequence set forth in SEQ ID NO:12 or SEQ ID NO:16.

[0011] In another embodiment, the agent is a soluble fragment of an IL-21R, and the soluble fragment of the IL-21R comprises an extracellular domain of IL-21R and an immunoglobulin Fc fragment. In a further embodiment, the amino acid sequence of the extracellular domain of the IL-21R comprises an amino acid sequence that is at least 90% identical to amino acids 1-235 of SEQ ID NO:2 or amino acids 20-235 of SEQ ID NO:2. In another embodiment, the immunoglobulin Fc fragment has an altered function. In another further embodiment, the immunoglobulin Fc fragment has the amino acid sequence of amino acids 244-467 of SEQ ID NO:17.

[0012] In another embodiment, the fibrosis or fibrosis-associated disorder affects the liver, epidermis, endodermis, muscle, tendon, cartilage, heart, pancreas, lung, uterus, nervous system, testis, ovary, adrenal gland, artery, vein, colon, small intestine, biliary tract, or stomach. In a further embodiment, the fibrosis or fibrosis-associated disorder is interstitial lung fibrosis. In another embodiment, the fibrosis or fibrosis-associated disorder is the result of an infection with schistosoma. In another embodiment, the fibrosis or fibrosis-associated disorder is the result of wound healing. In a further embodiment, the wound healing results from a surgical incision.

[0013] In another embodiment, the invention further comprises administering to the subject at least one additional therapeutic agent. In another embodiment, the at least one additional therapeutic agent is selected from the group consisting of cytokine inhibitors, growth factor inhibitors, immunosuppressants, anti-inflammatory agents, metabolic inhibitors, enzyme inhibitors, cytotoxic agents, and cytostatic agents. In a further embodiment, the at least one additional therapeutic agent is selected from the group consisting of TNF antagonists, anti-TNF agents, IL-12 antagonists, IL-15 antagonists, IL-17 antagonists, IL-18 antagonists, IL-22 antagonists, T cell-depleting agents, B cell-depleting agents, cyclosporin, FK506, CCI-779, etanercept, infliximab, rituximab, adalimumab, prednisolone, azathioprine, gold, sulphasalazine, hydroxychloroquine, minocycline, anakinra, abatacept, methotrexate, leflunomide, rapamycin, rapamycin analogs, Cox-2 inhibitors, cPLA2 inhibitors, NSAIDs, p38 inhibitors, antagonists of B7.1, B7.2, ICOSL, ICOS and/or CD28, and agonists of CTLA4.

[0014] In another embodiment, the invention provides a method for identifying a compound for treating, ameliorating or preventing fibrosis or a fibrosis-associated disorder in a subject, comprising: (a) measuring the level of IL-21 and/or IL-21R in a cell or sample of interest; (b) contacting the cell or sample of interest with a compound; and (c) measuring the level of IL-21 and/or IL-21R in the cell or sample of interest following contact with the compound, wherein a lower level of IL-21 and/or IL-21R in the contacted cell or sample of interest, in comparison to the level of IL-21 and/or IL-21R in a noncontacted cell or sample of interest, identifies the compound as a compound useful for treating, ameliorating, or preventing fibrosis or a fibrosis-associated condition in a subject.

[0015] In another embodiment, the invention provides a method for identifying a compound for treating, ameliorating or preventing fibrosis or a fibrosis-associated disorder in a subject, comprising: (a) measuring the level of IL-21 and/or IL-21R in a cell or sample of interest; (b) contacting the cell or sample of interest with a compound; (c) measuring the level of IL-21 and/or IL-21R in the cell or sample of interest following contact with the compound; and (d) comparing the level of IL-21 and/or IL-21R in the contacted cell or sample of interest with a reference level of IL-21 and/or IL-21R, wherein a lower level of IL-21 and/or IL-21R in the contacted cell or sample of interest, in comparison to the reference level of IL-21 and/or IL-21R, identifies the compound as a compound useful for treating, ameliorating, or preventing fibrosis or a fibrosis-associated condition in a subject.

[0016] In another embodiment, the invention provides a method for monitoring the progress of fibrosis or a fibrosis-associated condition in a subject, comprising: (a) measuring the level of IL-21 and/or IL-21R in a cell or sample of interest from the subject at a first time point; and (b) measuring the level of IL-21 and/or IL-21R in a cell or sample of interest from the subject at a second time point, wherein a lower level of IL-21 and/or IL-21R in the cell or sample of interest from the subject at the second time point, in comparison to the level of IL-21 and/or IL-21R in the cell or sample of interest from the subject at the first time point, provides an indication that the fibrosis or fibrosis-associated condition has decreased in severity.

[0017] In another embodiment, the invention provides a method for prognosing fibrosis or a fibrosis-associated condition in a subject, comprising: (a) measuring

the level of IL-21 and/or IL-21R in a cell or sample of interest from the subject at a first time point; and (b) measuring the level of IL-21 and/or IL-21R in a cell or sample of interest from the subject at a second time point, wherein a lower level of IL-21 and/or IL-21R in the cell or sample of interest from the subject at the second time point, in comparison to the level of IL-21 and/or IL-21R in the cell or sample of interest from the subject at the first time point, indicates a decreased likelihood that the subject will develop fibrosis or the fibrosis-associated condition or a decreased likelihood that the fibrosis or fibrosis-associated condition will worsen in the subject.

[0018] In another embodiment, the invention provides a method for prognosing fibrosis or a fibrosis-associated condition in a subject, comprising: (a) measuring the level of IL-21 and/or IL-21R in a cell or sample of interest from the subject; and (b) comparing the level of IL-21 and/or IL-21R in the cell or sample of interest from the subject to a reference level of IL-21 and/or IL-21R, wherein a lower level of IL-21 and/or IL-21R in the cell or sample of interest from the subject, in comparison to the reference level of IL-21 and/or IL-21R, indicates a decreased likelihood that the subject will develop fibrosis or the fibrosis-associated condition or a decreased likelihood that the fibrosis or fibrosis-associated condition will worsen in the subject.

[0019] In another embodiment, the invention provides a method for diagnosing fibrosis or a fibrosis-associated condition in a subject, comprising: (a) measuring the level of IL-21 and/or IL-21R in a cell or sample of interest from the subject, and (b) comparing the level of IL-21 and/or IL-21R in the cell or sample of interest from the subject with a reference level of IL-21 and/or IL-21R, wherein a higher level of IL-21 and/or IL-21R in the cell or sample of interest from the subject, in comparison to the reference level of IL-21 and/or IL-21R, indicates the presence of fibrosis or the fibrosis-associated condition in the subject.

BRIEF DESCRIPTION OF THE DRAWINGS

[0020] FIG. 1 depicts the full-length cDNA sequence of murine IL-21R/MU-1. The nucleotide sequence corresponds to nucleotides 1-2628 of SEQ ID NO:4.

[0021] FIGS. 2A and 2B depict the amino acid sequences of murine and human IL-21R/MU-1. FIG. 2A depicts the amino acid sequence of murine

IL-21R/MU-1 (corresponding to the amino acids 1-529 of SEQ ID NO:5). There is a leader sequence (predicted by SPScan with a score of 10.1) at amino acids 1-19 (boldface type). A predicted transmembrane domain (underlined) is found at amino acids 237-253 of SEQ ID NO:5. Predicted signaling motifs include the "Box 1" motif at amino acids 265-274 and the "Box 2" motif at amino acids 311-324 (bold and underlined). Six tyrosines are located at amino acid positions 281, 319, 361, 368, 397, and 510, of SEQ ID NO:5. The WSXWS motif (SEQ ID NO:3) is located at amino acid residues 214-218 (in large, boldface type). Potential Stat docking sites include amino acids 393-398 and amino acids 510-513 of SEQ ID NO:5. FIG. 2B depicts the amino acid sequence of human IL-21R/MU-1 (corresponding to SEQ ID NO:2). The location of the predicted signal sequence (about amino acids 1-19 of SEQ ID NO:2); WSXWS motif (about amino acids 213-217 of SEQ ID NO:2); and transmembrane domain (about amino acids 236-252 (or 236-253, or 236-254) of SEQ ID NO:2 (underlined)) are indicated. Potential JAK binding sites, Box 1 and 2 signaling motifs, and Stat docking sites are indicated by labeled arrows.

[0022] FIG. 3 depicts the GAP comparison of human and murine IL-21R/MU-1 cDNA sequences (corresponding to nucleic acids 1-2665 of SEQ ID NO:1 and nucleic acids 1-2628 of SEQ ID NO:4, respectively). huMU-1=human IL-21R/MU-1, murMU-1=murine IL-21R/MU-1. Gap Parameters: Gap Weight=50; Average Match=10.000; Length Weight=3; Average Mismatch=0.000; Percent Identity=66.116.

[0023] FIG. 4 depicts a GAP comparison of human IL-21R/MU-1 protein (corresponding to amino acids of SEQ ID NO:2) and murine IL-21R/MU-1 protein (corresponding to amino acids of SEQ ID NO:5). The alignment was generated by BLOSUM62 amino acid substitution matrix (Henikoff and Henikoff (1992) *Proc. Natl. Acad. Sci. U.S.A.* 89:10915-19). Gap parameters=Gap Weight: 8; Average Match=2.912; Length Weight=2; Average Mismatch=-2.003; Percent Identity=65.267.

[0024] FIG. 5 depicts a multiple sequence alignment of the amino acids of human IL-21R/MU-1 (corresponding to SEQ ID NO:2), murine IL-21R/MU-1 (corresponding to SEQ ID NO:5), and human IL-2 beta chain (GENBANK® Accession No. M26062). Leader and transmembrane domains are underlined.

Conserved cytokine receptor module motifs are indicated by boldface type.

Potential signaling regions are indicated by underlining and boldface type.

[0025] FIG. 6 depicts signaling through IL-21R/MU-1. IL-21R/MU-1 phosphorylates Stat 5 in Clone E7 EPO IL-21R/MU-1 chimera expressing cells stimulated with EPO. Treatment of controls or chimeric BAF-3 cells with IL-3 resulted in phosphorylation of Stat 3, but not Stat 1 or 5.

[0026] FIGS. 7A-7B depict an alignment of the nucleotide and amino acid sequences of mature human IL-21R fused at the amino terminal to a honeybee leader sequence and His₆ and Flag tags. The nucleotide and amino acid sequences of the fusion protein depicted in FIGS. 7A-7B are set forth in SEQ ID NO:10 and SEQ ID NO:11, respectively. The amino acid sequences of the mature human IL-21R fragment and the honeybee leader / His tags fragment of the fusion protein correspond to amino acids 20-235 of SEQ ID NO:2 and amino acids 1-44 of SEQ ID NO:11, respectively.

[0027] FIGS. 8A-8C depict an alignment of the nucleotide and amino acid sequences of human IL-21R extracellular domain fused at the C-terminus via a linker to human immunoglobulin G1 (IgG1) Fc sequence. The nucleotide and amino acid sequences of the fusion protein depicted in FIGS. 8A-8C are set forth in SEQ ID NO:12 and SEQ ID NO:13, respectively. The amino acid sequences of the human IL-21R extracellular domain, the linker, and the human immunoglobulin G1 (IgG1) Fc sequence correspond to amino acids 1-235 of SEQ ID NO:2, amino acids 236-243 of SEQ ID NO:13, and amino acids 244-467 of SEQ ID NO:13, respectively.

[0028] FIGS. 9A-9C depict an alignment of the nucleotide and amino acid sequences of human IL-21R extracellular domain fused at the C-terminus via a linker to human immunoglobulin G1 (IgG1) Fc sequence and His₆ sequence tag. The nucleotide and amino acid sequences of the fusion protein depicted in FIGS. 9A-9C are set forth in SEQ ID NO:14 and SEQ ID NO:15, respectively. The amino acid sequences of the human IL-21R extracellular domain, the linker, the human immunoglobulin G1 (IgG1) Fc sequence, and the His₆ sequence tag correspond to amino acids 1-235 of SEQ ID NO:2, amino acids 236-243 of SEQ ID NO:15, amino acids 244-467 of SEQ ID NO:15, and amino acids 468-492 of SEQ ID NO:15, respectively.

[0029] FIGS. 10A-10C depict an alignment of the nucleotide and amino acid sequences of human IL-21R extracellular domain fused at the C-terminus via a linker to human immunoglobulin G1 (IgG1) Fc-mutated sequence. The human Fc sequence has been mutated at residues 254 and 257 from the wild-type sequence to reduce Fc-receptor binding. The nucleotide and amino acid sequences of the fusion protein depicted in FIGS. 10A-10C are set forth in SEQ ID NO:16 and SEQ ID NO:17, respectively. The amino acid sequences of the human IL-21R extracellular domain, the linker, and the human immunoglobulin G1 (IgG1) Fc-mutated sequence correspond to amino acids 1-235 of SEQ ID NO:2, amino acids 236-243 of SEQ ID NO:17, and amino acids 244-467 of SEQ ID NO:17, respectively.

[0030] FIGS. 11A-11B depict an alignment of the nucleotide and amino acid sequences of human IL-21R extracellular domain fused at the C-terminus to a rhodopsin sequence tag. The nucleotide and amino acid sequences of the fusion protein are set forth in SEQ ID NO:18 and SEQ ID NO:19, respectively. The amino acid sequence of the human IL-21R extracellular domain corresponds to amino acids 1-235 of SEQ ID NO:2.

[0031] FIGS. 12A-12C depict an alignment of the nucleotide and amino acid sequences of human IL-21R extracellular domain fused at the C-terminus to an EK cleavage site and mutated IgG1 Fc region. The nucleotide and amino acid sequences of the fusion protein depicted in FIGS. 12A-12C are set forth in SEQ ID NO:20 and SEQ ID NO:21, respectively. The amino acid sequences of the human IL-21R extracellular domain, and the EK cleavage site/mutated IgG1 Fc region correspond to amino acids 1-235 of SEQ ID NO:2 and amino acids 236-470 of SEQ ID NO:21, respectively.

[0032] FIGS. 13A-13B depict an alignment of the nucleotide and amino acid sequences of murine IL-21R extracellular domain fused at the C-terminus to mouse immunoglobulin G2a (IgG2a). The nucleotide and amino acid sequences are set forth in SEQ ID NO:22 and SEQ ID NO:23, respectively.

[0033] FIGS. 14A-14B depict an alignment of the nucleotide and amino acid sequences of murine IL-21R extracellular domain fused at the C-terminus to Flag and His₆ sequence tags. The nucleotide and amino acid sequences are set forth in SEQ ID NO:24 and SEQ ID NO:25, respectively.

[0034] FIGS. 15A-15B depict an alignment of the nucleotide and amino acid sequences of (honeybee leader) murine IL-21R extracellular domain fused at the N-terminus to Flag and His₆ sequence tags. The nucleotide and amino acid sequences are set forth in SEQ ID NO:26 and SEQ ID NO:27, respectively.

[0035] FIG. 16 shows IL-21 and IL-21R expression profiles during highly polarized type-1 and type-2 immune responses. Groups of five IL-10/IL-4 KO (TH1, Δ) and IL-10/IL-12 KO (TH2, \bullet) mice were sensitized i.p. with *S. mansoni* eggs and challenged i.v. 14 days later. Lung RNA specimens were prepared individually for real-time RT-PCR analysis of IL-13 and IFN- γ (FIG. 16A) and IL-21R and IL-21 (FIG. 16B). The means \pm SEM in gene expression were expressed as fold-increases over unchallenged WT controls after normalization to HPRT. Asterisks denote significant differences between groups at the given time point, * $p < 0.05$.

[0036] FIG. 17 shows that type-2 cytokine production is reduced in the lungs of schistosome egg-challenged IL-21R^{-/-} mice. Groups of naïve WT (open bars) and IL-21R^{-/-} mice (filled bars) were i.v. challenged with live *S. mansoni* eggs and sacrificed on days 4, 7, and 14 post-challenge. (A) RNA was prepared from lung tissues and analyzed individually (N=5 per group/time point) by real-time RT-PCR. Results are shown as box-and-whisker plots with five-number summary bars showing the median, the quartiles, and the smallest and greatest percentiles in the distribution; bars (from bottom to top) indicate the 10th, 25th, 50th, 75th, and 90th percentiles, respectively, of the tested samples. The asterisks denote significant differences from wild-type values at the given time point, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. (B) Spleens (Spl) and lung-associated lymph nodes (LN) were each pooled (2 separate groups, 3-4 mice per group) and single cell suspensions were assayed for IL-5, IL-10, IL-13, and IFN- γ after a 72-h incubation in the presence of Con A (CON, 1 $\mu\text{g/ml}$) or soluble egg antigen (SEA, 20 $\mu\text{g/ml}$). Cytokines were below the level of detection in unstimulated cultures. (C) Granuloma size (volume, $\text{mm}^3 \times 10^{-3}$) and the percentage of eosinophils in granulomas were quantified microscopically. (D) Real-time PCR analysis of T_H2-regulated inflammatory genes in granulomatous lung tissue. All data are representative of at least 2 separate experiments.

[0037] FIG. 18 shows that the type-2 response is impaired in *N. brasiliensis*-infected IL-21R^{-/-} mice. Lungs (A) and lung-associated lymph nodes (LALN) (B) were removed on day 7 from individual *N. brasiliensis*-infected and untreated C57BL/6 or IL-21R^{-/-} mice (5/treatment group). RNA was isolated and cDNA was generated as described in the legend to FIG. 17. mRNA was analyzed individually for IL-13, IL-4, AMCase, Ym1, and FIZZ1 by real-time quantitative PCR. Fold changes are based on comparisons of infected mice to naive animals.

[0038] FIG. 19 shows that type-2 cytokine-driven inflammation is reduced in IL-21R^{-/-} mice. WT (open bars) and IL-21R^{-/-} mice (filled bars) were sensitized i.p. with eggs, challenged i.v. two weeks later with live *S. mansoni* eggs and then sacrificed on days 4 and 7 post-challenge. (A) RNA was prepared from lung tissues and analyzed individually (N=5 per group/time point) by real-time RT-PCR as described above in the legend to FIG. 17. (B) Spleens (Spl) and lung-associated lymph nodes (LN) were assayed for IL-5, IL-10, IL-13, and IFN- γ following antigen (SEA) or mitogen stimulation (CON). (C) Granuloma size (mm³ X 10⁻³) and the percentage of eosinophils in granulomas were quantified microscopically in WT (mice per group: N=10, day 4; N=15, day 7) and IL-21R^{-/-} mice (N=11, day 4; N=16, day 7). (D) Real-time PCR analysis of T_H2 inflammatory genes in granulomatous lung tissue (N=5 per group/time point). The asterisks denote significant differences from wild-type values at the given time point, * p<0.05, ** p<0.01, *** p<0.001. Data shown are the combined results of 3 separate experiments.

[0039] FIG. 20 shows that chronic liver disease following percutaneous *S. mansoni* infection is reduced in the absence of IL-21R. WT (open bars) and IL-21R^{-/-} mice (filled bars) were infected with 25-30 *S. mansoni* cercariae. All animals were sacrificed at week 9 (acute) or week 12 (chronic) post-infection. (A) RNA was isolated from liver tissues and analyzed individually (N=8-10 per group/time point) by real-time RT-PCR as described above in the legend to FIG. 17. (B) Spleens (Spl) and mesenteric lymph nodes (LN) were pooled in groups of 2-4 mice and single cell suspensions were assayed for IL-5, IL-10, and IFN- γ . The data shown are the averages of three separate pooled groups. (C) Granuloma size (mm³ X 10⁻³) and the percentage of eosinophils in granulomas were evaluated microscopically in WT mice (mice per group: N=30 for week 9,

N=17 for week 12) and IL-21R^{-/-} mice (mice per group: N=27 for week 9, N=19 for week 12). (D) Real-time PCR analysis of Th2 inflammatory genes in granulomatous liver tissue (N=8-10 per group/time point). Data shown are the combined results of 3 separate experiments conducted on week 9 and two performed on week 12. The asterisks denote significant differences from wild-type values at the given time point, * p<0.05, ** p<0.01, *** p<0.001.

[0040] FIG. 21 shows that the cellular composition of granulomas is unchanged by IL-21R deficiency. (A) The cellular composition of granulomas was evaluated in the livers of 9 week infected WT (N=10) and IL-21R^{-/-} (N=9) mice. The average \pm SEM of small lymphocytes (Sm Lym), large lymphocytes (Lg Lym), macrophages (Mac), Fibroblasts (Fibro), Eosinophils (Eos), and Mast Cells (Mc) are shown. (B) Lymphocytes were isolated from the perfused lungs of naïve WT and IL-21R^{-/-} mice (top panels) and on day 7 following i.v. challenge with 5000 *S. mansoni* eggs (bottom panels). The numbers in the histograms indicate the percentages of CD4⁻ and CD4⁺ T cells among total lung lymphocytes.

[0041] FIG. 22 shows that IL-21R-deficiency significantly slows the progression of T_H2 cytokine-dependent fibrosis. WT (open bars) and IL-21R^{-/-} mice (filled bars) were infected with *S. mansoni* cercariae. Animals were sacrificed at 9 (acute), 12 (chronic) (panels A-D) or 29 weeks (late chronic) (panel E) post-infection. (A) The average worm pairs, total worms and eggs/worm pair in thousands \pm SE are shown for each group. No difference in infection intensity was noted in any experiment (*n*=number of mice). (B) Mice were bled at the time of sacrifice and SEA isotype-specific Ab titers were determined by ELISA. (C) Total serum IgE values in μ g/ml. (D-F) Fibrosis was assessed by the amount of hydroxyproline in micromoles detected in the liver per 10,000 eggs (panel D) or in total liver (panels E and F). In panel F, infected WT C57BL/6 mice were treated with either an IgG2a control antibody (cIg – open bar) or with sIL-21R-Fc (filled bar) for 6 weeks. The asterisks denote significant differences from wild-type values at the given time point, * p<0.05, ** p<0.01, *** p<0.001.

[0042] FIG. 23 shows that IL-21 signaling promotes alternative macrophage activation by modulating IL-13 receptor expression. Bone marrow-derived macrophages were treated with various combinations of IL-4 (20 ng/ml), IL-13 (20 ng/ml), and IL-21 (20 ng/ml) overnight. Macrophages treated with IL-4,

IL-13, and IL-21 were pretreated with IL-21 for 6 hours prior to administration of IL-4 and IL-13. Cells were lysed 20 hr later and RNA was analyzed individually by real-time RT-PCR. (A) The ability of IL-21 to promote alternative macrophage activation was assessed by measuring Arg-1 and FIZZ1 gene expression. (B) Arginase activity was quantified in cell lysates by measuring the conversion of L-arginine to urea (mg/dL \pm SEM, triplicate measurements) (C) Expression of IL-4R α and IL-13R α 1 mRNA was evaluated by real-time PCR. IL-13R α 2 mRNA was nearly undetectable in all conditions (not shown). The data shown in panels A, B, and C are representative of 3 separate experiments. (D) Naïve C57BL/6 mice were challenged intravenously with 5000 live *S. mansoni* eggs and treated with PBS or rIL-21 (2 μ g/dose) every other day from day 1 through day 6. Animals (5 per group) were sacrificed on day 7 and lung IL-13R α 2 mRNA levels were assayed by real-time PCR and expressed as fold-increase over untreated controls (open bar). Mice were also bled at the time of sacrifice and the amount of sIL-13R α 2 in individual serum samples was assayed by ELISA. The asterisks denote significant differences, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

[0043] FIG. 24 shows that alternatively activated macrophages do not produce significant quantities of active TGF- β 1. The right panel shows active TGF- β 1 following macrophage activation, while the left panel shows total TGF- β 1 levels following macrophage activation. Bone marrow-derived macrophages were treated with various combinations of IL-4 (20 ng/ml), IL-13 (20 ng/ml), and IL-21 (20 ng/ml) overnight. Macrophages treated with IL-4, IL-13, and IL-21 were pretreated with IL-21 for 6 hours prior to administration of IL-4 and IL-13. 20 hours post-activation, supernatants were assayed for total and active TGF- β 1 by ELISA. High levels of total TGF- β 1 were detected in all groups (e.g., compare left panel "IL-4" with "Untreated") except for cells treated with IL-21 alone (compare left panel "IL-21" with "Untreated"). Although total TGF- β 1 expression was high, active TGF- β 1 was minimal in all groups (right panel). The data shown are representative of 3 separate experiments producing similar results.

DETAILED DISCRIPTION OF THE INVENTION

[0044] To examine the role of the IL-21/IL-21R signaling pathway in the pathogenesis of fibrosis, immune responses were compared in mice lacking a functional IL-21R (IL-21R^{-/-}) and in wild type mice using various models of lung and liver inflammation. In one model, live schistosome eggs were injected intravenously into naïve or antigen-sensitized animals to study primary and secondary granulomatous inflammation in the lung. In another model, mice were infected percutaneously with *S. mansoni* cercariae and the development of egg-induced inflammation and fibrosis was observed in the liver. In another model, mice were infected with *N. brasiliensis*. Using these models, the influence of the IL-21R on type-2 cytokine-driven pathology in acute and chronic disease settings was studied. The results demonstrate an important role for the IL-21R in the generation of polarized type-2 responses *in vivo*, particularly in type-2 cytokine-mediated inflammation and fibrosis.

[0045] The present invention therefore provides methods for treating, ameliorating, or preventing fibrosis or fibrosis-associated disorders in a subject (e.g., a human, e.g., a human patient) using an agent(s) that reduces the level of IL-21 and/or IL-21R (e.g., the level of expression of IL-21 and/or IL-21R (e.g., the level of IL-21 and/or IL-21R gene products (i.e., protein and/or mRNA)), the level of activity of IL-21 and/or IL-21R, the level of interaction of IL-21 with IL-21R, etc.) relative to an untreated control (e.g., a control subject afflicted with fibrosis or a fibrosis-associated condition, a control subject not afflicted with fibrosis or a fibrosis-associated condition) or relative to an appropriate reference level. In relation to identifying an agent(s) that reduces the level of IL-21 and/or IL-21R, measuring “the level of IL-21 and/or IL-21R” includes, but is not limited to, (1) measuring the level of expression of IL-21 and/or IL-21R (e.g., measuring the level of IL-21 and/or IL-21R gene products (e.g., protein and/or its corresponding mRNA)); (2) measuring the level of activity of IL-21 and/or IL-21R; and (3) measuring the level of interaction of IL-21 with IL-21R (e.g., in a cell or sample of interest, e.g., from a subject (e.g., a human patient, a control subject)). As described in further detail herein, exemplary agents useful to treat, ameliorate, and/or prevent fibrosis or fibrosis-associated conditions or disorders include anti-IL-21R antibodies, antigen-binding fragments of anti-IL-21R antibodies, anti-IL-21 antibodies, antigen-binding fragments of anti-IL-21

antibodies, and soluble fragments of IL-21R polypeptides. The invention further provides methods for monitoring the course of treatment of fibrosis or a fibrosis-associated disorder, diagnosing and prognosing the same, and screening for compounds useful to treat fibrosis or a fibrosis-associated disorder.

[0046] As used herein, “IL-21” or “IL-21R” means any polypeptide that is substantially identical to the naturally occurring IL-21 or IL-21 receptor protein, respectively. The nucleotide and amino acid sequences encoding human interleukin-21 (IL-21) and its receptor (IL-21R) are described, for example, in WO 00/53761, WO 01/85792, Parrish-Novak et al. (2000) *Nature* 408:57-63, and Ozaki et al. (2000) *Proc. Natl. Acad. Sci. U.S.A.* 97:11439-44. Desirably, the IL-21 polypeptide binds IL-21R or the IL-21R polypeptide binds IL-21, and upon interaction there is an increase in the signaling activity of the IL-21/IL-21R pathway by at least 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or more than 100% above control levels, as measured by any standard method. IL-21R is also known as “MU-1,” “NLR,” and “zalphal1.”

[0047] An agent that decreases the level of IL-21 and/or IL-21R encompasses any agent that decreases the signaling activity of the IL-21/IL-21R pathway, the level of activity of IL-21 and/or IL-21R, the level of expression of IL-21 and/or IL-21R, and/or the level of interaction of IL-21 and IL-21R by at least 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100%. Optionally, the decrease in levels of IL-21 and/or IL-21R is assessed by measuring the level of reduction in fibrosis. An agent that decreases the level of interaction of IL-21 with IL-21R encompasses any agent that decreases the interaction of IL-21 with IL-21R by at least 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100%. These aforementioned agents that decrease the level of activity or expression of IL-21 and/or IL-21R and/or decrease the level of interaction of IL-21 with IL-21R (i.e., agents that decrease the level of IL-21 and/or IL-21R) may be referred to herein as “antagonists” of IL-21 and/or IL-21R.

[0048] An “IL-21 gene” or “IL-21R gene” is defined as a nucleic acid that encodes an IL-21 or IL-21R polypeptide, respectively.

[0049] “Fibrosis” is defined as any pathological condition resulting from an overproduction or aberrant production of fibrous tissue. Fibrosis may occur in any organ including, for example, kidney, lung, liver, skin, central nervous system, bone, bone marrow, cardiovascular system, an endocrine organ or the

gastrointestinal system. By “fibrosis-associated condition” is meant any condition that is related to fibrosis. Thus, fibrosis-associated conditions may be caused by, be concomitant with, or cause fibrosis.

[0050] Decreasing the level of activity of IL-21 and/or IL-21R may refer to a reduction in the level or biological activity of IL-21 relative to the level or biological activity of IL-21 and/or IL-21R in an untreated control or reference sample (e.g., a reference level). Such level or activity may be decreased by at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100%. A decrease in the activity of IL-21 and/or IL-21R may also be associated with a reduction in type-2 (“Th2”) cytokine expression and/or function, which may include a modulation in, e.g., IL-4, IL-13, AMCase, Ym1, and Fizz1/RELM α -levels and activity.

[0051] Decreasing the level of interaction of IL-21 with IL-21R may refer to a decrease in the interaction in a treated cell or sample relative to the level of interaction of IL-21 with IL-21R in an untreated control or reference sample. Such level of interaction may be decreased by at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100%. The level of interaction may be assessed by several well-known molecular biology techniques, e.g., ELISA and Western blotting.

[0052] Decreasing the level IL-21 and/or IL-21R gene products refers to a decrease in the mRNA and/or protein expression level in a treated cell or sample relative to the gene or protein expression level of IL-21 and/or IL-21R in an untreated control or reference sample. Such expression may be decreased by at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100%. The level of expression may be assessed by a number of well-known molecular biology techniques, e.g., Northern blotting or Western blotting.

[0053] By “treating or ameliorating fibrosis” is meant decreasing the level of fibrosis relative to an untreated control, as measured by any standard method. A reduction in fibrosis may also be measured by a reduction in any symptom associated with fibrosis or a fibrosis-associated condition. The examples disclosed herein provide exemplary methods of determining whether the level of fibrosis is decreased relative to a control.

[0054] By “treating or ameliorating a fibrosis-associated condition (or disorder)” is meant decreasing such condition before or after it has occurred. As compared with an equivalent untreated control, such reduction or degree of prevention is at least 5%, 10%, 20%, 40%, 50%, 60%, 80%, 90%, 95%, or 100% as measured by any standard technique. A subject who is being treated for a fibrosis-associated condition is one who a medical practitioner has diagnosed as having such a condition. Diagnosis may be by any suitable means. A subject in whom the development of a fibrosis-associated condition is being prevented may or may not have received such a diagnosis. One in the art will understand that these subjects (e.g., patients) may have been subjected to standard tests for diagnosing fibrosis-associated conditions or may have been identified, without examination, as one at high risk due to the presence of one or more risk factors. The examples disclosed herein provide exemplary methods of determining whether the level of a fibrosis-associated disorder is decreased relative to a control.

[0055] “Preventing” refers to delaying the onset of a fibrosis or a fibrosis-associated condition, or prohibiting the onset of fibrosis or a fibrosis-associated condition in a subject likely to develop such a condition.

[0056] By “an effective amount” is meant an amount of a compound, alone or in a combination, required to treat, ameliorate, reduce or prevent fibrosis or a fibrosis-associated condition in a mammal. The effective amount of active compound(s) varies depending upon the route of administration, age, body weight, and general health of the subject. Ultimately, the attending physician or veterinarian will decide the appropriate amount and dosage regimen.

[0057] By “substantially identical,” when referring to a protein or polypeptide, is meant a protein or polypeptide exhibiting at least 75%, but preferably 85%, more preferably 90%, most preferably 95%, or even 99% identity to a reference amino acid sequence, e.g., SEQ ID NO:2, SEQ ID NO:5, and fusion proteins such as those set forth in SEQ ID NOs:11, 13, 15, 17, 19, 21, 23, 25 and 27. For proteins or polypeptides, the length of comparison sequences will generally be at least 20 amino acids, preferably at least 30 amino acids, more preferably at least 40 amino acids, and most preferably 50 amino acids or the full-length protein or polypeptide. Nucleic acids that encode such “substantially identical” proteins or polypeptides constitute examples of “substantially identical” nucleic acids. It is recognized that, due to the redundancy of the genetic code, several nucleic acids

may encode a given protein or polypeptide; such nucleic acids are within the scope of the invention if they encode a polypeptide that is “substantially identical” to a reference polypeptide.

[0058] The nucleic acids related to the present invention may comprise DNA or RNA and may be wholly or partially synthetic. Reference to a nucleotide sequence as set forth herein encompasses a DNA molecule with the specified sequence (or a complement thereof), and encompasses an RNA molecule with the specified sequence in which U is substituted for T, unless context requires otherwise.

[0059] The isolated polynucleotides related to the present invention may be used as hybridization probes and primers to identify and isolate nucleic acids having sequences identical to or similar to those encoding the disclosed polynucleotides. Hybridization methods for identifying and isolating nucleic acids include polymerase chain reaction (PCR), Southern hybridization, *in situ* hybridization and Northern hybridization, and are well known to those skilled in the art.

[0060] Hybridization reactions may be performed under conditions of different stringency. The stringency of a hybridization reaction includes the difficulty with which any two nucleic acid molecules will hybridize to one another. Preferably, each hybridizing polynucleotide hybridizes to its corresponding polynucleotide under reduced stringency conditions, more preferably stringent conditions, and most preferably highly stringent conditions. Examples of stringency conditions are shown in Table 1 below: highly stringent conditions are those that are at least as stringent as, for example, conditions A-F; stringent conditions are at least as stringent as, for example, conditions G-L; and reduced stringency conditions are at least as stringent as, for example, conditions M-R.

Table 1: Stringency Conditions

Stringency Condition	Poly-nucleotide Hybrid	Hybrid Length (bp) ¹	Hybridization Temperature and Buffer ²	Wash Temperature and Buffer ²
A	DNA:DNA	>50	65°C; 1xSSC -or- 42°C; 1xSSC, 50% formamide	65°C; 0.3xSSC
B	DNA:DNA	<50	T _B *; 1xSSC	T _B *; 1xSSC
C	DNA:RNA	>50	67°C; 1xSSC -or- 45°C; 1xSSC, 50% formamide	67°C; 0.3xSSC

Stringency Condition	Poly-nucleotide Hybrid	Hybrid Length (bp) ¹	Hybridization Temperature and Buffer ²	Wash Temperature and Buffer ²
D	DNA:RNA	<50	T _D *; 1xSSC	T _D *; 1xSSC
E	RNA:RNA	>50	70°C; 1xSSC -or- 50°C; 1xSSC, 50% formamide	70°C; 0.3xSSC
F	RNA:RNA	<50	T _F *; 1xSSC	T _F *; 1xSSC
G	DNA:DNA	>50	65°C; 4xSSC -or- 42°C; 4xSSC, 50% formamide	65°C; 1xSSC
H	DNA:DNA	<50	T _H *; 4xSSC	T _H *; 4xSSC
I	DNA:RNA	>50	67°C; 4xSSC -or- 45°C; 4xSSC, 50% formamide	67°C; 1xSSC
J	DNA:RNA	<50	T _J *; 4xSSC	T _J *; 4xSSC
K	RNA:RNA	>50	70°C; 4xSSC -or- 50°C; 4xSSC, 50% formamide	67°C; 1xSSC
L	RNA:RNA	<50	T _L *; 2xSSC	T _L *; 2xSSC
M	DNA:DNA	>50	50°C; 4xSSC -or- 40°C; 6xSSC, 50% formamide	50°C; 2xSSC
N	DNA:DNA	<50	T _N *; 6xSSC	T _N *; 6xSSC
O	DNA:RNA	>50	55°C; 4xSSC -or- 42°C; 6xSSC, 50% formamide	55°C; 2xSSC
P	DNA:RNA	<50	T _P *; 6xSSC	T _P *; 6xSSC
Q	RNA:RNA	>50	60°C; 4xSSC -or- 45°C; 6xSSC, 50% formamide	60°C; 2xSSC
R	RNA:RNA	<50	T _R *; 4xSSC	T _R *; 4xSSC

1: The hybrid length is that anticipated for the hybridized region(s) of the hybridizing polynucleotides. When hybridizing a polynucleotide to a target polynucleotide of unknown sequence, the hybrid length is assumed to be that of the hybridizing polynucleotide. When polynucleotides of known sequence are hybridized, the hybrid length can be determined by aligning the sequences of the polynucleotides and identifying the region or regions of optimal sequence complementarity.

2: SSPE (1xSSPE is 0.15M NaCl, 10mM NaH₂PO₄, and 1.25mM EDTA, pH 7.4) can be substituted for SSC (1xSSC is 0.15M NaCl and 15mM sodium citrate) in the hybridization and wash buffers; washes are performed for 15 minutes after hybridization is complete.

T_B* - T_R*: The hybridization temperature for hybrids anticipated to be less than 50 base pairs in length should be 5-10°C less than the melting temperature (T_m) of the hybrid, where T_m is determined according to the following equations. For hybrids less than 18 base pairs in length, T_m(°C) = 2(# of A + T bases) + 4(# of G + C bases). For hybrids between 18 and 49 base pairs in length, T_m(°C) = 81.5 + 16.6(log₁₀Na⁺) + 0.41(%G+C) - (600/N), where N is the number of bases in the hybrid, and Na⁺ is the concentration of sodium ions in the hybridization buffer (Na⁺ for 1xSSC = 0.165M).

Additional examples of stringency conditions for polynucleotide hybridization are provided in Sambrook, J., E.F. Fritsch, and T. Maniatis, 1989, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, chapters 9 and 11, and *Current Protocols in Molecular Biology*, 1995, F.M. Ausubel et al., eds., John Wiley & Sons, Inc., sections 2.10 and 6.3-6.4, incorporated herein by reference.

[0061] The isolated polynucleotides related to the present invention may be used as hybridization probes and primers to identify and isolate DNA having sequences encoding allelic variants of the disclosed polynucleotides. Allelic variants are naturally occurring alternative forms of the disclosed polynucleotides that encode polypeptides that are identical to or have significant similarity to the

polypeptides encoded by the disclosed polynucleotides. Preferably, allelic variants have at least 90% sequence identity (more preferably, at least 95% identity; most preferably, at least 99% identity) with the disclosed polynucleotides. Alternatively, significant similarity exists when the nucleic acid segments will hybridize under selective hybridization conditions (e.g., highly stringent hybridization conditions) to the disclosed polynucleotides.

[0062] The isolated polynucleotides related to the present invention may also be used as hybridization probes and primers to identify and isolate DNAs having sequences encoding polypeptides homologous to the disclosed polynucleotides. These homologs are polynucleotides and polypeptides isolated from a different species than that of the disclosed polypeptides and polynucleotides, or within the same species, but with significant sequence similarity to the disclosed polynucleotides and polypeptides. Preferably, polynucleotide homologs have at least 50% sequence identity (more preferably, at least 75% identity; most preferably, at least 90% identity) with the disclosed polynucleotides, whereas polypeptide homologs have at least 30% sequence identity (more preferably, at least 45% identity; most preferably, at least 60% identity) with the disclosed polypeptides. Preferably, homologs of the disclosed polynucleotides and polypeptides are those isolated from mammalian species.

[0063] Calculations of "homology" or "sequence identity" between two sequences are performed by means well known to those of skill in the art. For example, one general means for calculating sequence identity is described as follows. The sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and nonhomologous sequences can be disregarded for comparison purposes). In a preferred embodiment, the length of a reference sequence aligned for comparison purposes is at least 30%, preferably at least 40%, more preferably at least 50%, even more preferably at least 60%, and even more preferably at least 70%, 80%, 90%, 100% of the length of the reference sequence. The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a

function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

[0064] The comparison of sequences and determination of percent sequence identity between two sequences may be accomplished using a mathematical algorithm. In a preferred embodiment, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch ((1970) *J. Mol. Biol.* 48:444-53) algorithm, which has been incorporated into the GAP program in the GCG software package (available at www.gcg.com), using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another preferred embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package (available at www.gcg.com), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. One preferred set of parameters is a Blossum 62 scoring matrix with a gap penalty of 12, a gap extend penalty of 4, and a frameshift gap penalty of 5. The percent identity between two amino acid or nucleotide sequences can also be determined using the algorithm of Meyers and Miller ((1989) *CABIOS* 4:11-17), which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

[0065] "Substantially pure" is defined as a nucleic acid, polypeptide, or other molecule that has been separated from the components that naturally accompany it, e.g., genetic material, associated proteins, membranes, and cell debris. Typically, a polypeptide is substantially pure if it is at least 60%, 70%, 80%, 90%, 95%, or even 99%, by weight, free from the proteins and naturally occurring organic molecules with which it naturally associates. For example, a substantially pure polypeptide may be obtained by extraction from a natural source, by expression of a recombinant nucleic acid in a cell that does not normally express that protein, or by chemical synthesis.

[0066] The term "isolated DNA" is defined as DNA that is relatively or substantially free of the genes and other DNA sequences that flank the DNA in the naturally occurring genome of the organism from which the given DNA is

derived. Thus, the term “isolated DNA” encompasses, for example, cDNA, cloned genomic DNA, and synthetic DNA.

[0067] “IL-21 fusion” polypeptide or protein, or “IL-21R fusion” polypeptide or protein is defined as all or part of an IL-21 or IL-21R amino acid sequence, e.g., the IL-21R extracellular fragment from amino acids 1-235 of SEQ ID NO:2, linked to a second, heterologous amino acid sequence. In one embodiment, the second, heterologous amino acid sequence is a tag sequence. Common tag sequences include myc tags, his tags, flag tags, etc. In another embodiment of the invention, the second, heterologous amino acid sequence is an immunoglobulin sequence, e.g., an Fc fragment. Such fusion proteins and polypeptides, which are described in greater detail herein, are encoded by nucleic acid sequences referred to as “IL-21 fusion genes” or “IL-21R fusion genes.”

[0068] In the screening methods a “compound” refers to a chemical, whether naturally occurring or artificially derived. Such compounds may include, for example, peptides, polypeptides, synthetic organic molecules, naturally occurring organic molecules, nucleic acid molecules, peptide nucleic acid molecules, and components and derivatives thereof. For example, a useful compound according to the present invention reduces binding of IL-21 to IL-21R.

[0069] Antagonists of IL-21 and IL-21R for use in treating, ameliorating, or preventing fibrosis or a fibrosis-associated condition may also consist of small molecules. The term “small molecule” refers to compounds that are not macromolecules (see, e.g., Karp (2000) *Bioinformatics Ontology* 16:269-85; Verkman (2004) *AJP-Cell Physiol.* 286:465-74). Thus, small molecules are often considered those compounds that are, e.g., less than one thousand daltons (e.g., Voet and Voet, *Biochemistry*, 2nd ed., ed. N. Rose, Wiley and Sons, New York, 14 (1995)). For example, Davis et al. (2005) *Proc. Natl. Acad. Sci. USA* 102:5981-86, use the phrase small molecule to indicate folates, methotrexate, and neuropeptides, while Halpin and Harbury (2004) *PLoS Biology* 2:1022-30, use the phrase to indicate small molecule gene products, e.g., DNAs, RNAs and peptides. Examples of natural and synthesized small molecules include, but are not limited to, cholesterol, neurotransmitters, siRNAs, and various chemicals listed in numerous commercially available small molecule databases, e.g., FCD (Fine Chemicals Database), SMID (Small Molecule Interaction Database), ChEBI (Chemical Entities of Biological Interest), and CSD (Cambridge

Structural Database) (see, e.g., Alfarano et al. (2005) *Nuc. Acids Res. Database Issue* 33:D416-24).

[0070] The term “pharmaceutical composition” means any composition that contains at least one therapeutically or biologically active agent and is suitable for administration to a subject. Any of these formulations can be prepared by well-known and accepted methods of the art. See, for example, *Remington: The Science and Practice of Pharmacy*, 21st Ed., (ed. A.R. Gennaro), Lippincott Williams & Wilkins, Baltimore, MD (2005).

[0071] The present invention provides significant advantages over standard therapies for treatment and prevention of fibrosis-associated conditions. As described herein, administration of a therapeutic agent that reduces the level of activity or expression of IL-21 and/or IL-21R or decreases the interactions between IL-21 and IL-21R (i.e., reduces the level of IL-21 and/or IL-21R activity) results in amelioration, reduction or prevention of fibrosis and fibrosis-associated conditions. In addition, the compound screening methods, provided by this invention, allow one to identify novel therapeutics that modify the injury process, rather than merely mitigating the symptoms.

Fibrotic Disorders

[0072] The generation of granulation tissue is a carefully orchestrated process in which the expression of protease inhibitors and extracellular matrix proteins is upregulated and the expression of proteases is reduced, leading to the accumulation of extracellular matrix. Abnormal accumulation of fibrous materials, however, may ultimately lead to organ failure (e.g., Border et al. (1994) *New Engl. J. Med.* 331:1286-92). The development of fibrotic conditions, whether induced or spontaneous, is caused at least in part by the stimulation of fibroblast activity. The influx of inflammatory cells and activated fibroblasts into the injured organ depends on the ability of these cell types to interact with the interstitial matrix, which contains primarily collagens. Exemplary tissues that may be affected by fibrosis include the kidney, lung, liver, skin, central nervous system, bone, bone marrow, tissues of the cardiovascular system, endocrine organs, and tissues of the gastrointestinal system.

[0073] The methods and compositions of the present invention are useful for any fibrosis or fibrosis-associated condition affecting any tissue including, for

example, fibrosis of an internal organ, a cutaneous or dermal fibrosing disorder, and fibrotic conditions of the eye. Fibrosis of internal organs (e.g., liver, lung, kidney, heart blood vessels, gastrointestinal tract) occurs in disorders such as pulmonary fibrosis, idiopathic fibrosis, autoimmune fibrosis, myelofibrosis, liver cirrhosis, veno-occlusive disease, mesangial proliferative glomerulonephritis, crescentic glomerulonephritis, diabetic nephropathy, renal interstitial fibrosis, renal fibrosis in subjects receiving cyclosporin, allograft rejection, HIV associated nephropathy. Other fibrosis-associated disorders include systemic sclerosis, eosinophilia-myalgia syndrome, and fibrosis-associated CNS disorders such as intraocular fibrosis. Dermal fibrosing disorders include, for example, scleroderma, morphea, keloids, hypertrophic scars, familial cutaneous collagenoma, and connective tissue nevi of the collagen type. Fibrotic conditions of the eye include conditions such as diabetic retinopathy, post-surgical scarring (for example, after glaucoma filtering surgery and after crossed-eyes (strabismus) surgery), and proliferative vitreoretinopathy. Additional fibrotic conditions that may be treated by the methods of the present invention may result, for example, from rheumatoid arthritis, diseases associated with prolonged joint pain and deteriorated joints; progressive systemic sclerosis, polymyositis, dermatomyositis, eosinophilic fascitis, morphea, Raynaud's syndrome, and nasal polyposis. As described herein, an IL-21/IL-21R pathway antagonist may be administered to treat or prevent fibrosis and fibrosis-associated disorders, or to ameliorate one or more of the symptoms associated with these disorders.

IL-21 or IL-21R Antagonists (IL-21/IL-21R Antagonists)

[0074] The IL-21 antagonists or IL-21R antagonists of the invention interact with IL-21 or IL-21R (e.g., mammalian IL-21 or IL-21R such as human, bovine, rat, mouse, horse, or dog), respectively, and reduce the level of IL-21 and/or IL-21R, e.g., reduce one or more biological activities associated with IL-21 and/or IL-21R. When this interaction involves direct binding, antagonists bind to IL-21 or IL-21R with high affinity (e.g., with an affinity constant of at least about 10^7 M^{-1} , preferably about 10^8 M^{-1} , and more preferably, about 10^9 M^{-1} to 10^{10} M^{-1} or stronger).

[0075] The level of IL-21 and/or IL-21R is desirably reduced by 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or even 100%. An antagonist may, for

example, reduce the activity of IL-21R by neutralizing IL-21. An antagonist may be a fusion protein that includes a fragment of an IL-21R fused to a non-IL-21R fragment such as an immunoglobulin Fc region, e.g., the fusion proteins set forth in SEQ ID NOs:11, 13, 15, 17, 19, 21, 23, 25 and 27. Other exemplary antagonists are anti-IL-21R or anti-IL-21 antibodies or antigen-binding fragments thereof, soluble forms of IL-21R, peptides, inhibitory polynucleotides (e.g., siRNAs, SNPs, and aptamers) and small molecules.

[0076] In one embodiment, the IL-21/IL-21R antagonist is an anti-IL-21R or anti-IL-21 antibody, or an antigen-binding fragment thereof. In a further embodiment, the antibody is a neutralizing antibody. If desired, the antibody may be a monoclonal or single-specificity antibody that binds to IL-21 or IL-21R or an antigen-binding fragment thereof (e.g., an Fab, F(ab')₂, Fv or a single chain Fv fragment). The antibody may be human, humanized, chimeric, or *in vitro*-generated antibody to IL-21 or IL-21R polypeptide.

[0077] Alternatively, the IL-21 antagonist or IL-21R antagonist may be a full length (e.g., a mutated sequence) or a fragment of an IL-21 polypeptide or an IL-21R polypeptide (e.g., human). Exemplary antagonists include, for example, an inhibitory IL-21 receptor-binding domain of an IL-21 polypeptide (e.g., human) or the extracellular domain of murine or human IL-21R. The IL-21 antagonist may have an amino acid sequence that is substantially identical to (e.g., having at least 85%, 90%, 95%, 98%, 99% sequence identity with) the naturally occurring IL-21R (e.g., SEQ ID NO:2 (human) or SEQ ID NO:5 (murine)) or a fragment thereof (see Table 2). Alternatively, the antagonist may have an amino acid sequence encoded by a nucleotide sequence that is substantially identical to the naturally occurring mammalian IL-21R or a fragment thereof (e.g., SEQ ID NO:1 (human) or SEQ ID NO:4 (murine)) or by a nucleotide sequence that hybridizes to one of the foregoing nucleotide sequences under stringent conditions, e.g., highly stringent conditions (see Table 1).

Table 2: Summary of Sequences

Designation	Description
SEQ ID NO:1	Nucleic acid sequence of human IL-21R cDNA
SEQ ID NO:2	Amino acid sequence of human IL-21R
SEQ ID NO:3	Amino acid sequence of conserved WSXWS motif
SEQ ID NO:4	Nucleic acid sequence of mouse IL-21R cDNA
SEQ ID NO:5	Amino acid sequence of mouse IL-21R
SEQ ID NO:6	Amino acid sequence of Fc fragment
SEQ ID NO:7	Nucleic acid (cDNA) sequence of human IL-21
SEQ ID NO:8	Amino acid sequence of human IL-21
SEQ ID NO:9	Peptide signal sequence
SEQ ID NO:10	Nucleic acid (cDNA) sequences of human IL-21R monomer fused to a honeybee leader sequence and amino terminal His ₆ and Flag tags
SEQ ID NO:11	Amino acid sequence of human IL-21R monomer (20-235) fused to a honeybee leader sequence and amino terminal His ₆ and Flag tags
SEQ ID NO:12	Nucleic acid (cDNA) sequence of human IL-21R extracellular domain (1-235) fused to an IgG fragment
SEQ ID NO:13	Amino acid sequence of human IL-21R extracellular domain (1-235) fused to an IgG fragment
SEQ ID NO:14	Nucleic acid (cDNA) sequence of human IL-21R extracellular domain (1-235) fused to an IgG fragment and a His ₆ tag
SEQ ID NO:15	Amino acid sequence of human IL-21R extracellular domain (1-235) fused to an IgG fragment and a His ₆ tag
SEQ ID NO:16	Nucleic acid (cDNA) sequence of human IL-21R extracellular domain (1-235) fused to an IgG fragment mutated at residues 254 and 257
SEQ ID NO:17	Amino acid sequence of human IL-21R extracellular domain (1-235) fused to an IgG fragment mutated at residues 254 and 257
SEQ ID NO:18	Nucleic acid (cDNA) sequence of human IL-21R monomer fused to a Rhodopsin tag
SEQ ID NO:19	Amino acid sequence of human IL-21R monomer (20-235) fused to a Rhodopsin tag
SEQ ID NO:20	Nucleic acid (cDNA) sequence of human IL-21R extracellular domain (1-235) fused to EK cleavage sites and an IgG1 fragment with a mutated Fc region
SEQ ID NO:21	Amino acid sequence of human IL-21R extracellular domain (1-235) fused to EK cleavage sites and an IgG1 fragment with a mutated Fc region
SEQ ID NO:22	Nucleic acid sequence of mouse IL-21R extracellular domain fused to a mouse genomic IgG2a fragment

Designation	Description
SEQ ID NO:23	Amino acid sequence of mouse IL-21R extracellular domain fused to a mouse genomic IgG2a fragment
SEQ ID NO:24	Nucleic acid (genomic) sequence of mouse IL-21R extracellular domain fused to Flag and His ₆ tags
SEQ ID NO:25	Amino acid sequence of mouse IL-21R extracellular domain fused to fused to Flag and His ₆ tags
SEQ ID NO:26	Nucleic acid sequence of mouse IL-21R extracellular domain fused to a honeybee leader sequence and amino terminal Flag and His ₆ tags
SEQ ID NO:27	Amino acid sequence of mouse IL-21R extracellular domain fused to a honeybee leader sequence and amino terminal Flag and His ₆ tags
SEQ ID NO:28	Sense primer for mouse IL-21
SEQ ID NO:29	Antisense primer for mouse IL-21
SEQ ID NO:30	Sense primer for mouse IL-21R
SEQ ID NO:31	Antisense primer for mouse IL-21R
SEQ ID NO:32	Sense primer for mouse IFN- γ
SEQ ID NO:33	Antisense primer for mouse IFN- γ

[0078] Optionally, the IL-21R polypeptide may be a soluble polypeptide incapable of membrane anchoring. Such soluble polypeptides include, for example, IL-21R polypeptides that lack a sufficient portion of their membrane spanning domain or are modified such that the membrane spanning domain is nonfunctional. For example, the IL-21R polypeptide may be a soluble fragment of an IL-21R (e.g., a fragment of an IL-21R containing the extracellular domain of murine or human IL-21R, including an amino acid sequence from about amino acids 1-235, 1-236, 20-235, or 20-236 of SEQ ID NO:2 (human), or from about amino acids 1-236, or 20-236 of SEQ ID NO:5 (murine)). Exemplary IL-21 antagonists may have an amino acid sequence that is substantially identical to amino acids 20-538 of SEQ ID NO:2 (mature human IL-21R), amino acids 1-235 of SEQ ID NO:2 (extracellular domain of human IL-21R), amino acids 1-236 of SEQ ID NO:2, amino acids 20-235 of SEQ ID NO:2, amino acids 20-236 of SEQ ID NO:2, amino acids 1-236 of SEQ ID NO:5, or amino acids 20-236 of SEQ ID NO:5.

[0079] An IL-21 antagonist of the invention may also be encoded by nucleic acids that hybridize to the nucleotide sequence set forth in SEQ ID NO:1, SEQ

ID NO:4, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, or SEQ ID NO:26, under highly stringent conditions, e.g., as set forth in Table 1. Isolated polynucleotides which encode IL-21R proteins or fusion proteins, but which differ from the nucleotide sequence set forth in SEQ ID NOs:1, 4, 10, 12, 14, 16, 18, 20, 22, 24, or 26, by virtue of the degeneracy of the genetic code, are also encompassed by the present invention. Variations in the nucleotide sequence as set forth in SEQ ID NOs:1, 4, 10, 12, 14, 16, 18, 20, 22, 24, or 26, which are caused by point mutations or by induced modifications, are also included in the invention.

[0080] If desired, a soluble IL-21R polypeptide may include, or be fused to, a second moiety such as a polypeptide (e.g., an immunoglobulin chain, a GST, Lex-A or MBP polypeptide sequence). For example, a fusion protein may include a fragment of an IL-21R polypeptide, which is capable of binding IL-21, such as a soluble fragment of an IL-21R (e.g., a fragment containing the extracellular domain of murine or human IL-21R such as amino acids 1-235, 1-236, 20-235 or 20-236 of SEQ ID NO:2 (human), or amino acids 1-236 or 20-236 of SEQ ID NO:5 (murine)) fused to a second moiety (e.g., an immunoglobulin chain, an Fc fragment, a heavy chain constant regions of various immunoglobulin isotypes, including IgG1, IgG2, IgG3, IgG4, IgM, IgA1, IgA2, IgD, and IgE).

[0081] Desirably, the IL-21 antagonist of the invention reduces at least one biological activity associated with the naturally occurring IL-21R, including, for example, the ability to interact with or bind to an IL-21 polypeptide, the ability to associate with signal transduction molecules such as γc or JAK1, the ability to stimulate phosphorylation and/or activation of stat proteins (e.g., Stat 5 and/or Stat 3), and the ability to modulate (e.g., stimulate or decrease) proliferation, differentiation, effector cell function, cytolytic activity, cytokine secretion, and/or survival of immune cells such as T cells (CD8+ and CD4+ T cells, including Th1 and Th2 cells), NK cells, B cells, macrophages, and megakaryocytes.

[0082] According to the present invention, an IL-21 polypeptide is a cytokine showing sequence homology to IL-2, IL-4 and IL-15 (Parrish-Novak et al. (2000) *Nature* 408:57-63). Despite low sequence homology among interleukin cytokines, cytokines share a common secondary motif, i.e., a "four-helix-bundle"

structure that is representative of the family. IL-21 is expressed primarily in activated CD4⁺ T cells, and has been reported to have effects on NK, B and T cells (Parrish-Novak et al. (2000) *supra*; Kasaian et al. (2002) *Immunity* 16:559-69). IL-21 binds to IL-21R (also referred to as "MU-1," "NILR," and "alpha11"). Upon IL-21 binding, activation of IL-21R leads to Stat5 and/or Stat3 signaling (Ozaki et al. (2000) *supra*).

[0083] The amino acid sequences of IL-21 polypeptides are publicly known. For example, the nucleotide sequence and amino acid sequence of a human IL-21 is available at GenBank Acc. No. NM_021803. The disclosed human IL-21 nucleotide sequence is presented below:

```

1   gctgaagtga aaacgagacc aaggtctagc tctactgttg gtacttatga gatccagtcc
61  tggcaacatg gagaggattg tcactctgtc gatggctatc ttcttgggga cactgggtcca
121 caaatcaagc tcccaaggtc aagatcgcca catgattaga atgcgtcaac ttatagatat
181 tggtgatcag ctgaaaaatt atgtgaatga cttgggtcct gaatttctgc cagctccaga
241 agatgtagag acaaaactgtg agtggtcagc tttttcctgc tttcagaagg cccaactaaa
301 gtcagcaa at acaggaaaca atgaaaggat aatcaatgta tcaattaaaa agctgaagag
361 gaaaccacct tccacaaatg cagggaagag acagaaacac agactaacat gcccttcatg
421 tgattcttat gagaaaaaac caccxaaaga attcctagaa agattcaaat cacttctcca
481 aaagatgatt catcagcatc tgtcctctag aacacacgga agtgaagatt cctgaggatc
541 taacttgcatg ttggacacta tgttacatac tctaataatag tagtgaaagt catttcttg
601 tattccaagt ggaggag (SEQ ID NO:7)

```

[0084] The amino acid sequence of the disclosed human IL-21 polypeptide is presented below:

```

MRSSPGNMERIVICLMVIFLGLTLVHKSSSQGQDRHMIRMRLIDIVDQLKNYVND
LVPEFLPAPEDVETNCEWSAFSCFQKAQLKSANTGNNERI INVS IKKLKRKPPST
NAGRRQKHRLTCPSYKPKPEFLERFKSLQKMIHQHLSSRTHGSEDS
(SEQ ID NO:8)

```

[0085] Thus, an IL-21 polypeptide refers to a protein that is capable of interacting with or binding to IL-21R and having one of the following features: (i) an amino acid sequence substantially identical to a naturally occurring mammalian IL-21 or a fragment thereof (e.g., SEQ ID NO:8 (human)); (ii) an amino acid sequence which is encoded by a nucleotide sequence that is substantially identical to a naturally occurring mammalian IL-21 nucleotide sequence or a fragment thereof (e.g., SEQ ID NO:7 (human) or a fragment thereof); (iii) an amino acid sequence encoded by a nucleotide sequence degenerate to a naturally occurring IL-21 nucleotide sequence or a fragment thereof, e.g., SEQ ID NO:7 (human) or a fragment thereof; or (iv) a nucleotide

sequence that hybridizes to one of the foregoing nucleotide sequence sequences under stringent conditions.

[0086] In all foregoing aspects of the invention, the IL-21 or IL-21R polypeptides may be provided as a variant polypeptide having a mutation in the naturally occurring IL-21 or IL-21R sequence (wild type) that results in higher affinity (relative to the nonmutated sequence) binding to IL-21R or IL-21, respectively. Such mutations may be useful, for example, to increase resistance to proteolysis (relative to the nonmutated sequence). Some amino acid sequences in the disclosed sequences can be varied without significantly modifying IL-21 or IL-21R structure or function. In general, it is possible to replace residues that form IL-21 or IL-21R protein tertiary structure, provided that residues that perform a similar function are used. In other instances, the type of residue may be completely irrelevant if an alteration occurs in a noncritical area. Thus, the invention further includes IL-21 and IL-21R variants that show substantial IL-21-type biological activity. Such variants include deletions, insertions, inversions, repeats, and type substitutions (for example, substituting one hydrophilic residue for another, but not a strongly hydrophilic residue for a strongly hydrophobic residue). Small changes or "neutral" amino acid substitutions will often have little impact on protein function. (Taylor (1986) *J. Theor. Biol.* 119:205-18). Conservative substitutions may include, but are not limited to, replacements among the aliphatic amino acids, substitutions between amide residues, exchanges of basic residues, and replacements among the aromatic residues. Further guidance concerning which amino acid change is likely to be phenotypically silent (i.e., is unlikely to significantly affect function) can be found in Bowie et al. (1990) *Science* 247:1306-10 and Zvelebil et al. (1987) *J. Mol. Biol.* 195:957-61.

[0087] Optionally, the IL-21 or IL-21R antagonist is a fusion protein containing the IL-21 or IL-21R polypeptides or fragments thereof described herein fused to a second moiety such as an immunoglobulin chain, e.g., an Fc fragment, an epitope (tag) sequence, e.g., GST or myc, and additional well-known sequences such as Lex-A, or MBP polypeptide sequence. If desired, the fusion protein may include a fragment of an IL-21R polypeptide that is capable of binding IL-21, such as a soluble fragment of an IL-21R (e.g., a fragment of an IL-21R containing the extracellular domain of murine or human IL-21R from about amino acids 1-235,

1-236, 20-235, or 20-236 of SEQ ID NO:2 (human), or from about amino acids 1-236, or 20-236 of SEQ ID NO:5 (murine) or a fragment identical to, or substantially identical to, a polypeptide encoded by SEQ ID NOs:1 or 4) fused to a second moiety (e.g., an immunoglobulin chain, an Fc fragment, a heavy chain constant region(s) of the various isotypes, including: IgG1, IgG2, IgG3, IgG4, IgM, IgA1, IgA2, IgD, and IgE). Alternatively, the human Fc sequence has been mutated at one or more amino acids (e.g., mutated at residues 254 and 257 of SEQ ID NO:16) in the naturally occurring sequence to reduce Fc receptor binding. In other embodiments, the fusion protein may include the extracellular domain of murine IL-21R (from about amino acids 1-236, or 20-235 of SEQ ID NO:5 (murine)) fused to a murine immunoglobulin Fc chain (including, but not limited to, murine IgG, e.g., murine IgG2a or a mutated form of murine IgG2a).

[0088] Examples of antagonistic fusion proteins that may be used in the methods of the invention are shown in FIGS. 7-15. The fusion protein may include an amino acid sequence substantially identical to SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, or SEQ ID NO:27 or an amino acid sequence encoded by a nucleotide sequence that is substantially identical to SEQ ID NOs:10, 12, 14, 16, 18, 20, 22, 24, or 26. One exemplary fusion protein contains the human IL-21R extracellular domain (e.g., amino acids 1-235 of SEQ ID NO:2) fused at the C-terminus via a linker (corresponding to amino acids 236-243 of SEQ ID NO:17) to human immunoglobulin G1 (IgG1) Fc mutated sequence (corresponding to amino acids 244-467 of SEQ ID NO:17). The human Fc sequence has been mutated at residues 254 and 257 from the wild type sequence to reduce Fc receptor binding. The nucleotide and amino acid sequences are shown as SEQ ID NO:16 and SEQ ID NO:17, respectively.

[0089] The second polypeptide is preferably soluble. Optionally, the second polypeptide enhances the half-life, (e.g., the serum half-life) of the linked polypeptide. If desired, the second polypeptide includes a sequence that facilitates association of the fusion polypeptide with a second IL-21R or IL-21 polypeptide. The second polypeptide may include at least a region of an immunoglobulin polypeptide. Immunoglobulin fusions polypeptides are known in the art and are described in, e.g., U.S. Patent Nos. 5,225,538; 5,428,130; 5,514,582; 5,714,147; and 5,455,165.

[0090] Optionally, the second polypeptide is a full-length immunoglobulin polypeptide or a fragment thereof (e.g., a heavy chain, light chain, Fab, Fab₂, Fv, or Fc).

[0091] In one example, the second polypeptide has less effector function than the effector function of an Fc region of a wild-type immunoglobulin heavy chain. Fc effector function includes for example, Fc receptor binding, complement fixation and T cell-depleting activity (see, for example, U.S. Patent No. 6,136,310). Methods for assaying T cell-depleting activity, Fc effector function, and antibody stability are known in the art. In one embodiment, the second polypeptide has low or no affinity for the Fc receptor. In an alternative embodiment, the second polypeptide has low or no affinity for complement protein C1q.

[0092] A preferred second polypeptide sequence includes the amino acid sequence of SEQ ID NO:6. This sequence includes an Fc region. Underlined amino acids are those that differ from the amino acids found in the corresponding positions of the wild-type immunoglobulin sequence:

HTCPPCPAPEALGAPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAK
TKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPVP^{IEKTISKAKGQPREPQVYTLPP}
SREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQ
QGNVFSCSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO:6)

[0093] Fusion proteins may additionally include a linker sequence(s) joining the first moiety to the second moiety. For example, the fusion protein may include a peptide linker of about 4 to 20 amino acids, more preferably 5 to 10 amino acids in length, and most preferably about 8 amino acids in length. The amino acids in the peptide linker may include, e.g., Gly, Ser, Asn, Thr and Ala. Thus, a peptide linker may consist of a Gly-Ser element. In other embodiments, the fusion protein includes a peptide linker having the formula (Ser-Gly-Gly-Gly-Gly)_y, wherein "y" is 1, 2, 3, 4, 5, 6, 7, or 8.

[0094] In other embodiments, additional amino acid sequences can be added to the N- or C-terminus of the fusion protein to facilitate expression, detection and/or isolation or purification. For example, an IL-21/IL-21R fusion protein may be linked to one or more additional moieties, e.g., GST (i.e., glutathione S-transferase), His, FLAG, or myc tags. For example, the fusion protein may additionally include a GST peptide in which the fusion protein sequences are

fused to the C-terminus of the GST sequences. Such fusion proteins facilitate the purification or identification of the IL-21R/MU-1 fusion protein. In other embodiments, additional amino acid sequences may be added to the N- or C-terminus of the fusion protein to facilitate expression, steric flexibility, detection, and/or isolation or purification.

[0095] The fusion protein may also include a heterologous signal sequence at its N-terminus. For example, the native IL-21R signal sequence may be removed and replaced with a signal sequence from another protein. In certain host cells (e.g., mammalian host cells), expression and/or secretion of IL-21R may be increased using a heterologous signal sequence. A signal peptide that can be included in the fusion protein is MPLLLLLLLLLPSPLHP (SEQ ID NO:9). If desired, one or more amino acids can additionally be inserted between the first polypeptide moiety comprising the IL-21R/MU-1 moiety and the second polypeptide moiety.

[0096] The IL-21/IL-21R antagonists described herein may be derivatized or linked to another functional molecule, e.g., another peptide or protein (e.g., an Fab² fragment). For example, the fusion protein or an antibody, or antigen-binding portion, can be functionally linked (e.g., by chemical coupling, genetic fusion, noncovalent association or otherwise) to one or more other molecular entities, such as an antibody (e.g., a bispecific or a multispecific antibody), toxins, radioisotopes, cytotoxic or cytostatic agents, among others.

[0097] A chimeric or fusion protein of the invention may be produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different polypeptide sequences are ligated together in-frame in accordance with conventional techniques, e.g., by employing blunt-ended or sticky-ended termini for ligation, restriction enzyme digestion to create appropriate termini, filling-in of sticky ends as appropriate, alkaline phosphatase treatment to avoid undesirable ligation, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers that give rise to complementary overhangs between two consecutive gene fragments that can subsequently be annealed and reamplified to generate a chimeric gene sequence (see, for example, Ausubel et al., *supra*). Moreover, many expression vectors that encode a fusion moiety (e.g., an Fc

region of an immunoglobulin heavy chain) are commercially available. An IL-21R/MU-1 or IL-21 encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the immunoglobulin protein. In some embodiments, IL-21R/MU-1 or IL-21 fusion polypeptides exist as oligomers, such as dimers or trimers. An IL-21R/MU-1 or IL-21 monomer, and/or nucleic acids encoding an IL-21R/MU-1 or IL-21, can be constructed using methods known in the art.

Production of Nucleic Acids

[0098] The isolated polynucleotides of the invention may be operably linked to an expression control sequence, such as the pMT2 or pED expression vectors disclosed in Kaufman et al. (1991) *Nuc. Acids Res.* 19:4485-90, in order to produce the IL-21R or IL-21 polypeptides (including fragments and fusions thereof) recombinantly. Many suitable expression control sequences are known in the art. General methods of expressing recombinant proteins are also known and are exemplified in Kaufman (1990) *Meth. Enzym.* 185:537-66. As defined herein "operably linked" means enzymatically or chemically ligated to form a covalent bond between the isolated polynucleotide of the invention and the expression control sequence in such a way that the IL-21R or IL-21 polypeptide is expressed by a host cell that has been transformed (transfected) with the ligated polynucleotide/expression control sequence.

[0099] The term "vector," as used herein, is intended to refer to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid," which refers to a circular double stranded DNA loop into which additional DNA segments may be ligated. Another type of vector is a viral vector, wherein additional DNA segments may be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., nonepisomal mammalian vectors) can be integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "recombinant expression vectors" (or simply, "expression vectors"). In general,

expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, "plasmid" and "vector" may be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

[0100] The recombinant expression constructs of the invention may carry additional sequences, such as regulatory sequences (i.e., sequences that regulate either vector replication, e.g., origins of replication, transcription of the nucleic acid sequence encoding the polypeptide (or peptide) of interest, or expression of the encoded polypeptide), tag sequences such as histidine, and selectable marker genes. The term "regulatory sequence" is intended to include promoters, enhancers and any other expression control elements (e.g., polyadenylation signals, transcription splice sites) that control transcription, replication or translation. Such regulatory sequences are described, for example, in Goeddel, *Gene Expression Technology: Methods in Enzymology*, Academic Press, San Diego, CA (1990). It will be appreciated by those skilled in the art that the design of the expression vector, including the selection of regulatory sequences, will depend on various factors, including choice of the host cell and the level of protein expression desired. Preferred regulatory sequences for expression of proteins in mammalian host cells include viral elements that direct high levels of protein expression, such as promoters and/or enhancers derived from the FF-1a promoter and BGH poly A, cytomegalovirus (CMV) (e.g., the CMV promoter/enhancer), Simian virus 40 (SV40) (e.g., the SV40 promoter/enhancer), adenovirus (e.g., the adenovirus major late promoter (AdMLP)), and polyoma. Viral regulatory elements, and sequences thereof, are described in, e.g., U.S. Patent Nos. 5,168,062; 4,510,245; and 4,968,615.

[0101] The recombinant expression vectors of the invention may carry additional sequences, such as sequences that regulate replication of the vector in host cells (e.g., origins of replication and terminator sequences) and selectable marker genes. The selectable marker gene facilitates selection of host cells into which the vector has been introduced (see, e.g., U.S. Patents Nos. 4,399,216, 4,634,665 and 5,179,017, all by Axel et al.). For example, typically the selectable marker gene confers resistance of the host cell transfected or

transformed with the selectable marker to compounds such as G418 (geneticin), hygromycin or methotrexate. Preferred selectable marker genes include the dihydrofolate reductase (DHFR) gene (for use in *dhfr*⁻ host cells with methotrexate selection/amplification), the *neo* gene (for G418 selection), and genes conferring tetracycline and/or ampicillin resistance to bacteria.

[0102] Suitable vectors, containing appropriate regulatory sequences, including promoter sequences, terminator sequences, polyadenylation sequences, enhancer sequences, marker genes and other sequences as appropriate, may be either chosen or constructed. Inducible expression of proteins, achieved by using vectors with inducible promoter sequences, such as tetracycline-inducible vectors, e.g., pTet-On™ and pTet-Off™ (Clontech, Palo Alto, CA), may also be used in the disclosed method. For further details regarding expression vectors, see, for example, Sambrook et al., *supra*. Many known techniques and protocols for manipulation of nucleic acids, for example, in preparation of nucleic acid constructs, mutagenesis, sequencing, introduction of DNA into cells, gene expression, and analysis of proteins, are also described in detail in Sambrook et al., *supra*.

[0103] A number of types of cells may act as suitable host cells for expression of the IL-21R/MU-1 or IL-21 or fusion protein thereof. Any cell type capable of expressing functional IL-21R/MU-1 or IL-21 protein or fusion thereof may be used. Suitable mammalian host cells include, for example, monkey COS cells, Chinese Hamster Ovary (CHO) cells, human kidney 293 cells, human epidermal A431 cells, human Colo205 cells, 3T3 cells, CV-1 cells, other transformed primate cell lines, normal diploid cells, cell strains derived from *in vitro* culture of primary tissue, primary explants, HeLa cells, mouse L cells, BHK, HL-60, U937, HaK, C3H10T1/2, Rat2, BaF3, 32D, FDCP-1, PC12, M1x or C2C12 cells.

[0104] The IL-21R or IL-21 polypeptide or fusion protein thereof may also be produced by operably linking the isolated polynucleotide of the invention to suitable control sequences in one or more insect expression vectors, and employing an insect expression system. Materials and methods for baculovirus/Sf9 expression systems are commercially available in kit form (e.g., the MAXBAC® kit, Invitrogen, Carlsbad, CA). Soluble forms of the

polypeptides described herein may also be produced in insect cells using appropriate isolated polynucleotides as described above.

[0105] Alternatively, the IL-21R or IL-21 polypeptide or fusion protein thereof may be produced in lower eukaryotes such as yeast, or in prokaryotes such as bacteria. Suitable yeast strains include *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces strains*, *Candida*, or any yeast strain capable of expressing heterologous proteins. Suitable bacterial strains include *Escherichia coli*, *Bacillus subtilis*, *Salmonella typhimurium*, or any bacterial strain capable of expressing heterologous proteins. Expression in bacteria may result in formation of inclusion bodies incorporating the recombinant protein. Thus, refolding of the recombinant protein may be required in order to produce active or more active material. Several methods for obtaining correctly folded heterologous proteins from bacterial inclusion bodies are known in the art. These methods generally involve solubilizing the protein from the inclusion bodies, then denaturing the protein completely using a chaotropic agent. When cysteine residues are present in the primary amino acid sequence of the protein, it is often necessary to accomplish the refolding in an environment that allows correct formation of disulfide bonds (a redox system). General methods of refolding are disclosed in Kohno (1990) *Meth. Enzym.* 185:187-95, EP 0433225, and U.S. Patent No. 5,399,677.

[0106] A protein of the invention (or a fragment or fusion thereof) may also be expressed as a product of transgenic animals, e.g., as a component of the milk of transgenic cows, goats, pigs, or sheep which are characterized by somatic or germ cells containing a polynucleotide sequence, e.g., encoding the IL-21R or IL-21 or fusion protein thereof. Accordingly, the protein may be prepared by growing a culture transformed host cells under culture conditions necessary to express the desired protein. The resulting expressed protein may then be purified from the culture medium or cell extracts. Soluble forms of the protein may be purified from conditioned media. Membrane-bound forms of IL-21R protein of the invention can be purified by preparing a total membrane fraction from the expressing cell and extracting the membranes with a nonionic detergent such as TRITON® X-100 (EMD Biosciences, San Diego, CA).

[0107] The polypeptides described herein may be purified using methods known to those skilled in the art. For example, the protein of the invention may be

concentrated using a commercially available protein concentration filter, for example, by using an AMICON[®] or PELLICON[®] ultrafiltration unit (Millipore, Billerica, MA). Following the concentration step, the concentrate may be applied to a purification matrix such as a gel filtration medium. Alternatively, an anion exchange resin may be employed, for example, a matrix or substrate having pendant diethylaminoethyl (DEAE) or polyethyleneimine (PEI) groups. The matrices may be acrylamide, agarose, dextran, cellulose or other types commonly employed in protein purification. Alternatively, a cation exchange step may be employed. Suitable cation exchangers include various insoluble matrices comprising sulfopropyl or carboxymethyl groups. Sulfopropyl groups are preferred (e.g., S-SEPHAROSE[®] columns, Sigma-Aldrich, St. Louis, MO). The purification of the IL-21R/MU-1 protein or fusion protein from culture supernatant may also include one or more column steps over such affinity resins such as concanavalin A-agarose, AF-HEPARIN650, heparin-TOYOPEARL[®] or Cibacron blue 3GA SEPHAROSE[®] (Tosoh Biosciences, San Francisco, CA); or by hydrophobic interaction chromatography using such resins as phenyl ether, butyl ether, or propyl ether; or by immunoaffinity chromatography. Finally, one or more reverse-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, e.g., silica gel having pendant methyl or other aliphatic groups, can be employed to further purify the IL-21R/MU-1 or IL-21 protein. Affinity columns including antibodies to the protein of the invention may also be used in purification in accordance with known methods. Some or all of the foregoing purification steps, in various combinations or with other known methods, may also be employed to provide a substantially purified isolated recombinant protein. Preferably, the isolated protein is purified so that it is substantially free of other mammalian proteins.

Production of Antibodies

[0108] The IL-21 or IL-21R polypeptides of the invention may be used to immunize animals to obtain polyclonal and monoclonal antibodies that specifically react with the IL-21 or IL-21R and regulate the expression or activity of IL-21 and/or IL-21R, or regulate the level of interaction of IL-21 with IL-21R. Such antibodies may be obtained, for example, using the entire IL-21R or fragments thereof as immunogens. The peptide immunogens may additionally

contain a cysteine residue at the carboxyl terminus and be conjugated to a hapten such as keyhole limpet hemocyanin (KLH). Additional peptide immunogens may be generated by replacing tyrosine residues with sulfated tyrosine residues. Methods for synthesizing such peptides are known in the art, for example, as in Merrifield (1963) *J. Amer. Chem. Soc.* 85: 2149-54 and Krstenansky and Mao (1987) *FEBS Lett.* 211:10-16.

[0109] Human monoclonal antibodies (mAbs) directed against IL-21 or IL-21R may be generated using transgenic mice carrying the human immunoglobulin genes rather than the mouse system. Splenocytes from these transgenic mice immunized with the antigen of interest are used to produce hybridomas that secrete human mAbs with specific affinities for epitopes from a human protein (see, e.g., WO 91/00906, WO 91/10741, WO 92/03918, WO 92/03917, Lonberg et al. (1994) *Nature* 368:856-59, Green et al. (1994) *Nat. Genet.* 7:13-21, Morrison et al. (1994) *Proc. Natl. Acad. Sci. U.S.A.* 81:6851-55, and Tuaillon et al. (1993) *Proc. Natl. Acad. Sci. U.S.A.* 90:3720-24).

[0110] Monoclonal antibodies may also be generated by other methods known to those skilled in the art of recombinant DNA technology. One exemplary method, referred to as the "combinatorial antibody display" method, has been developed to identify and isolate antibody fragments having a particular antigen specificity, and can be utilized to produce monoclonal antibodies (for descriptions of combinatorial antibody display see, e.g., Sastry et al. (1989) *Proc. Natl. Acad. Sci. U.S.A.* 86:5728-32; Huse et al. (1989) *Science* 246:1275-81; and Orlandi et al. (1989) *Proc. Natl. Acad. Sci. U.S.A.* 86:3833-37). After immunizing an animal with an immunogen as described above, the antibody repertoire of the resulting B cell pool is cloned. The DNA sequence of the variable regions of a diverse population of immunoglobulin molecules may be obtained using a mixture of oligomer primers and PCR. For instance, mixed oligonucleotide primers corresponding to the 5' leader (signal peptide) sequences and/or framework 1 (FR1) sequences, as well as primer to a conserved 3' constant region primer may be used for PCR amplification of the heavy and light chain variable regions from a number of murine antibodies (Larrick et al. (1991) *BioTechniques* 11:152-56). A similar strategy may also be used to amplify human heavy and light chain variable regions from human antibodies (Larrick et al. (1991) *Methods: Companion to Methods in Enzymology* 2:106-10).

[0111] Chimeric antibodies, including chimeric immunoglobulin chains, may also be produced by recombinant DNA techniques known in the art. For example, a gene encoding the Fc constant region of a murine (or other species) monoclonal antibody molecule is digested with restriction enzymes to remove the region encoding the murine Fc, and the equivalent portion of a gene encoding a human Fc constant region is substituted (see PCT/US86/02269; EP 184,187; EP 171,496; EP 173,494; WO 86/01533; U.S. Patent No. 4,816,567; EP 125,023; Better et al. (1988) *Science* 240:1041-43; Liu et al. (1987) *Proc. Natl. Acad. Sci. U.S.A.* 84:3439-43; Liu et al. (1987) *J. Immunol.* 139:3521-26; Sun et al. (1987) *Proc. Natl. Acad. Sci. U.S.A.* 84:214-18; Nishimura et al. (1987) *Canc. Res.* 47:999-1005; Wood et al. (1985) *Nature* 314:446-49; and Shaw et al. (1988) *J. Natl. Cancer Inst.* 80:1553-59).

[0112] If desired, an antibody or an immunoglobulin chain may be humanized by methods known in the art. Humanized antibodies, including humanized immunoglobulin chains, may be generated by replacing sequences of the Fv variable region that are not directly involved in antigen binding with equivalent sequences from human Fv variable regions. General methods for generating humanized antibodies are provided by Morrison (1985) *Science* 229:1202-07; Oi et al. (1986) *BioTechniques* 4:214-21; and U.S. Patent Nos. 5,585,089, 5,693,761 and 5,693,762, all of which are hereby incorporated by reference in their entireties. Those methods include isolating, manipulating, and expressing the nucleic acid sequences that encode all or part of immunoglobulin Fv variable regions from at least one of a heavy or light chain. Sources of such nucleic acid are well known to those skilled in the art and, for example, may be obtained from a hybridoma producing an antibody against a predetermined target. The recombinant DNA encoding the humanized antibody, or fragment thereof, may then be cloned into an appropriate expression vector.

[0113] Humanized or CDR-grafted antibody molecules or immunoglobulins may be produced by CDR grafting or CDR substitution, wherein one, two, or all CDRs of an immunoglobulin chain can be replaced. See e.g., U.S. Patent No. 5,225,539; Jones et al. (1986) *Nature* 321:552-25; Verhoeyan et al. (1988) *Science* 239:1534-36; and Beidler et al. (1988) *J. Immunol.* 141:4053-60, all of which are hereby incorporated by reference in their entireties. U.S. Patent No. 5,225,539 describes a CDR-grafting method that may be used to prepare

humanized antibodies of the present invention (see also, GB 2188638A). All of the CDRs of a particular human antibody may be replaced with at least a portion of a nonhuman CDR, or only some of the CDRs may be replaced with nonhuman CDRs. It is only necessary to replace the number of CDRs required for binding of the humanized antibody to a predetermined antigen.

[0114] Monoclonal, chimeric and humanized antibodies, which have been modified by, e.g., deleting, adding, or substituting other portions of the antibody, e.g., the constant region, are also within the scope of the invention. For example, an antibody may be modified as follows: (i) by deleting the constant region; (ii) by replacing the constant region with another constant region, e.g., a constant region meant to increase half-life, stability or affinity of the antibody, or a constant region from another species or antibody class; or (iii) by modifying one or more amino acids in the constant region to alter, for example, the number of glycosylation sites, effector cell function, Fc receptor (FcR) binding, complement fixation, among others.

[0115] Methods for altering an antibody constant region are known in the art. Antibodies with altered function (e.g., altered affinity for an effector ligand, such as FcR on a cell, or the C1 component of complement) may be produced by replacing at least one amino acid residue in the constant portion of the antibody with a different residue (see, e.g., EP 388,151 A1, U.S. Patent Nos. 5,624,821 and 5,648,260, all of which are hereby incorporated by reference in their entireties). Similar types of alterations may also be applied to murine immunoglobulins and immunoglobulins from other species. For example, it is possible to alter the affinity of an Fc region of an antibody (e.g., an IgG, such as a human IgG) for an FcR (e.g., Fc gamma R1) or for C1q binding by replacing the specified residue(s) with a residue(s) having an appropriate functionality on its side chain, or by introducing a charged functional group, such as glutamate or aspartate, or perhaps an aromatic nonpolar residue such as phenylalanine, tyrosine, tryptophan or alanine (see, e.g., U.S. Patent No. 5,624,821).

[0116] Human antibodies to IL-21 and/or IL-21R may additionally be produced using transgenic nonhuman animals that are modified so as to produce fully human antibodies rather than the animal's endogenous antibodies in response to challenge by an antigen. See, e.g., PCT publication WO 94/02602. The endogenous genes encoding the heavy and light immunoglobulin chains in the

nonhuman host have been incapacitated, and active loci encoding human heavy and light chain immunoglobulins are inserted into the host's genome. The human genes are incorporated, for example, using yeast artificial chromosomes containing the requisite human DNA segments. An animal which provides all the desired modifications is then obtained as progeny by crossbreeding intermediate transgenic animals containing fewer than the full complement of the modifications. One embodiment of such a nonhuman animal is a mouse, and is termed the XENOMOUSETM as disclosed in PCT publications WO 96/33735 and WO 96/34096. This animal produces B cells that secrete fully human immunoglobulins. The antibodies can be obtained directly from the animal after immunization with an immunogen of interest, as, for example, a preparation of a polyclonal antibody, or alternatively from immortalized B cells derived from the animal, such as hybridomas producing monoclonal antibodies. Additionally, the genes encoding the immunoglobulins with human variable regions can be recovered and expressed to obtain the antibodies directly, or can be further modified to obtain analogs of antibodies such as, for example, single chain Fv molecules.

[0117] Other protein-binding molecules may also be employed to modulate the activity of IL-21 and/or IL-21R. Such protein-binding molecules include small modular immunopharmaceutical (SMIPTM) drugs (Trubion Pharmaceuticals, Seattle, WA). SMIPs are single-chain polypeptides composed of a binding domain for a cognate structure such as an antigen, a counterreceptor or the like, a hinge-region polypeptide having either one or no cysteine residues, and immunoglobulin CH2 and CH3 domains (see also www.trubion.com). SMIPs and their uses and applications are disclosed in, e.g., U.S. Published Patent Appln. Nos. 2003/0118592, 2003/0133939, 2004/0058445, 2005/0136049, 2005/0175614, 2005/0180970, 2005/0186216, 2005/0202012, 2005/0202023, 2005/0202028, 2005/0202534, and 2005/0238646, and related patent family members thereof, all of which are hereby incorporated by reference herein in their entireties.

[0118] As discussed herein, neutralizing or nonneutralizing antibodies (preferably monoclonal antibodies) binding to IL-21R or IL-21 or a fusion protein thereof may be useful in the treatment of immune conditions such as fibrosis and fibrosis-associated conditions.

[0119] Accordingly, the present invention further provides for compositions comprising an antibody that specifically reacts with an IL-21 or IL-21R or a fusion protein thereof.

Screening Assays

[0120] The IL-21R or IL-21 polypeptides or fusion proteins of the invention may also be used to screen for agents that are capable of binding to IL-21 or IL-21R, and regulating the level of IL-21 and/or IL-21R, e.g., the level of activity of IL-21 and/or IL-21R, the level of expression of IL-21 and/or IL-21R (e.g., the level of IL-21 and/or IL-21R gene products), and/or the level of interaction between IL-21 and IL-21R. Binding assays using a protein of the invention, or a binding partner thereof, which may be free or immobilized to a support, are well known in the art. Purified cell-based or protein-based (cell-free) screening assays may be used to identify binding partners and/or ligands (natural or synthetic, e.g., test compounds) to IL-21R or IL-21 polypeptides or fusion proteins of the invention. For example, IL-21R may be immobilized in purified form on a carrier and binding of potential ligands to IL-21R may be measured.

Methods for Diagnosing, Prognosing, and Monitoring the Progress of Fibrosis and Fibrosis-Associated Disorders, Related to IL-21

[0121] The present invention provides methods for diagnosing, prognosing, and monitoring the progress (e.g., monitoring the course of treatment) of disorders, i.e., fibrosis or fibrosis-associated conditions or disorders, related to IL-21 and/or IL-21R by, e.g., detecting and/or measuring the level of IL-21 and/or IL-21R, wherein the phrase "level of IL-21 and/or IL-21R" and equivalents thereof includes, but is not limited to, (1) the level of expression of IL-21 and/or IL-21R gene products (e.g., the level of IL-21 and/or IL-21R protein and/or mRNA in a cell or sample of interest); (2) the level of activity of IL-21 protein and/or IL-21R protein (e.g., Th2 cytokine expression and/or function in a cell or sample of interest); and (3) the level of interaction of IL-21 with IL-21R (e.g., in a cell or sample of interest). For example, the invention provides methods of diagnosing, prognosing and monitoring, e.g., by detecting the upregulation or downregulation of IL-21 and/or IL-21R gene products and/or activity, and/or by measuring the interaction of IL-21 with IL-21R, etc. (including but not limited to the use of such

methods in human subjects). IL-21 and/or IL-21R levels may also be measured in a reference cell or sample of interest to produce or obtain a reference level of IL-21 and/or IL-21R, or such reference level may be obtained through other methods, or may be generally known, by one of skill in the art. These methods may be performed by, e.g., utilizing prepackaged diagnostic kits comprising at least one of the group comprising an IL-21 or IL-21R polynucleotide (or fragments thereof); an IL-21 or IL-21R polypeptide (or fragments and/or fusion proteins thereof); an antibody to an IL-21 or IL-21R polypeptide (or derivatives thereof, or antigen-binding fragments thereof); or modulators of IL-21 or IL-21R polynucleotides and/or polypeptides as described herein, which may be conveniently used, for example, in a clinical setting.

[0122] “Diagnostic” or “diagnosing” means identifying the presence or absence of a pathologic condition. Diagnostic methods include, but are not limited to, detecting upregulation of the level of IL-21 and/or IL-21R by determining a test amount of the gene products (e.g., RNA, cDNA, or polypeptide, including fragments thereof) of IL-21 and/or IL-21R, by measuring the activity of IL-21 and/or IL-21R, and/or by measuring the level of interaction of IL-21 with IL-21R, in a biological sample from a subject (human or nonhuman mammal), and comparing the test amount with, e.g., a normal amount or range (e.g., an amount or range from an individual(s) known not to suffer from disorders related to IL-21, or from an individual known not to suffer from fibrosis or a fibrosis-associated condition). Although a particular diagnostic method may not provide a definitive diagnosis of disorders related to IL-21, it suffices if the method provides a positive indication that aids in diagnosis.

[0123] The present invention also provides methods for prognosing such disorders by detecting, for example, the upregulation of levels of IL-21 and/or IL-21R, e.g., by detecting upregulation of IL-21 and/or IL-21R gene products and/or activity, and/or by measuring the level of interaction of IL-21 with IL-21R, etc. “Prognostic” or “prognosing” means predicting the probable development and/or severity of a pathologic condition. Prognostic methods include determining the test amount of a gene product of IL-21 and/or IL-21R in a biological sample from a subject, and comparing the test amount to a prognostic amount or range (i.e., an amount or range from individuals with varying severities of disorders, i.e., fibrosis and/or a fibrosis-associated condition, e.g., related to IL-21) for the level

of IL-21 and/or IL-21R. Various amounts related to the level of IL-21 and/or IL-21R in a test sample are consistent with certain prognoses for disorders, e.g., fibrosis or fibrosis-associated conditions or disorders, related to IL-21 and/or IL-21R. The detection of an amount of IL-21 and/or IL-21R level at a particular prognostic level provides a prognosis for the subject.

[0124] The present invention also provides methods for monitoring the progress or course of such disorders related to IL-21 by detecting, for example, the upregulation of IL-21 and/or IL-21R levels, e.g., by detecting upregulation of IL-21 and/or IL-21R gene products, activity, and/or the interaction of IL-21 with IL-21R. Monitoring methods include determining the test amounts of a gene product of IL-21 in biological samples taken from a subject at a first and second time, and comparing the amounts. A change in amount of an IL-21 and/or IL-21R gene product between the first and second times indicates a change in the course of an IL-21-related disorder, with a decrease in amount indicating remission of such disorders, and an increase in amount indicating progression of such disorders. Such monitoring assays are also useful for evaluating the efficacy of a particular therapeutic intervention in patients being treated for fibrosis or a fibrosis-associated disorder.

Measuring the Level of IL-21 and/or IL-21R

[0125] The level of IL-21 and/or IL-21R (e.g., the level of IL-21 and/or IL-21 gene products, activity, and/or interaction) in methods of the invention (e.g., methods for screening for and/or identifying a compound for treating, ameliorating, or preventing fibrosis or a fibrosis-associated condition, methods of diagnosing, prognosing, and/or monitoring the progress of fibrosis or a fibrosis-associated condition, and methods of treating, ameliorating, or preventing fibrosis or a fibrosis-associated condition) outlined herein may be measured in a variety of biological samples, including bodily fluids (e.g., whole blood, plasma, and urine), cells (e.g., whole cells, cell fractions, and cell extracts), and other tissues. Biological samples also include sections of tissue, such as biopsies and frozen sections taken for histological purposes. Preferred biological samples include blood, plasma, lymph, tissue biopsies, urine, CSF (cerebrospinal fluid), synovial fluid, and BAL (bronchoalveolar lavage). It will be appreciated that analysis of a biological sample need not necessarily require removal of cells or tissue from the

subject. For example, appropriately labeled agents that bind IL-21 and/or IL-21R gene products (e.g., antibodies, nucleic acids) may be administered to a subject and visualized (when bound to the target) using standard imaging technology (e.g., CAT, NMR (MRI), and PET).

[0126] In the methods of treating, ameliorating, or preventing fibrosis or a fibrosis-associated disorder, in the methods for identifying a compound for treating, ameliorating or preventing fibrosis or a fibrosis-associated disorder in a subject, and in the diagnostic, prognostic, and monitoring assays and methods of the present invention, the level of IL-21 and/or IL-21R is detected and measured to yield a test amount. The test amount is then compared with, e.g., a normal amount or range. For example, an amount above (e.g., a higher level) the normal amount or range is a positive sign in the diagnosis of disorders related to IL-21.

[0127] Normal amounts or baseline levels of IL-21 and/or IL-21R may be determined for any particular sample type and population. Generally, baseline (normal) levels of IL-21 and/or IL-21R are determined by measuring respective levels of IL-21 and/or IL-21R in a biological sample type from normal (i.e., healthy) subjects. Alternatively, normal levels of IL-21 and/or IL-21R may be determined by measuring the amount in healthy cells or tissues taken from the same subject from which the diseased (or possibly diseased) test cells or tissues were taken. The level of IL-21 and/or IL-21R (either the normal amount or the test amount) may be determined or expressed on a per cell, per total protein, or per volume basis. To determine the cell amount of a sample, one can measure the level of a constitutively expressed gene product or other gene product expressed at known levels in cells of the type from which the biological sample was taken.

[0128] It will be appreciated that the assay methods of the present invention do not necessarily require measurement of absolute values for the level of IL-21 and/or IL-21R because relative values are sufficient for many applications of these methods. It will also be appreciated that in addition to the quantity or abundance of IL-21 and/or IL-21R levels, variant or abnormal IL-21 and/or IL-21R levels or their expression patterns (e.g., mutated transcripts, truncated polypeptides) may be identified by comparison to normal levels and expression patterns.

[0129] Whether the level of a particular gene or protein in two samples is increased (i.e., higher) or reduced (i.e., lower), e.g., significantly above or

significantly below a given level, respectively, depends on the gene itself and, *inter alia*, its variability in expression, activity, and/or interaction with a ligand between different individuals or different samples. It is within the skill in the art to determine whether IL-21 and/or IL-21R levels are significantly similar or different among samples. Factors such as genetic variation between individuals, species, organs, tissues, or cells may be taken into consideration (when and where necessary) for determining whether the level of IL-21 and/or IL-21R between two samples is increased or reduced. As a result of the natural heterogeneity in IL-21 and/or IL-21R levels between individuals, species, organs, tissues, or cells, phrases such as “significantly above” or “significantly below” cannot be defined as a precise percentage or value, but rather can be ascertained by one skilled in the art upon practicing the invention. Particular methods of detection and measurement of IL-21 and/or IL-21R gene products, activity, and interaction are described herein.

Assays for Measuring IL-21 and/or IL-21R Gene Products

[0130] The methods of the present invention involve detecting and quantifying the level of IL-21 and/or IL-21R, e.g., the level of the gene products of IL-21 and/or IL-21R, the level of activity of IL-21 and/or IL-21R, and/or the level of interaction of IL-21 with IL-21R, in biological samples. IL-21 and IL-21R gene products include mRNAs and polypeptides, and both can be measured using methods well known to those skilled in the art.

[0131] For example, mRNA can be directly detected and quantified using hybridization-based assays, such as Northern hybridization, *in situ* hybridization, dot and slot blots, and oligonucleotide arrays. Hybridization-based assays refer to assays in which a probe nucleic acid is hybridized to a target nucleic acid. In some formats, the target, the probe, or both are immobilized. The immobilized nucleic acid may be DNA, RNA, or another oligonucleotide or polynucleotide, and may comprise naturally or nonnaturally occurring nucleotides, nucleotide analogs, or backbones. Methods of selecting nucleic acid probe sequences for use in the present invention (e.g., based on the nucleic acid sequence of IL-21) are well known in the art.

[0132] Alternatively, mRNA may be amplified before detection and quantitation. Such amplification-based assays are well known in the art and include

polymerase chain reaction (PCR), reverse-transcription-PCR (RT-PCR), PCR-enzyme-linked immunosorbent assay (PCR-ELISA), and ligase chain reaction (LCR). Primers and probes for producing and detecting amplified IL-21 gene products (e.g., mRNA or cDNA) may be readily designed and produced without undue experimentation by those of skill in the art based on the nucleic acid sequences of IL-21 and/or IL-21R. Amplified IL-21 and/or IL-21R gene products may be directly analyzed, for example, by gel electrophoresis; by hybridization to a probe nucleic acid; by sequencing; by detection of a fluorescent, phosphorescent, or radioactive signal; or by any of a variety of well-known methods. In addition, methods are known to those of skill in the art for increasing the signal produced by amplification of target nucleic acid sequences. One of skill in the art will recognize that, whichever amplification method is used, a variety of quantitative methods known in the art (e.g., quantitative PCR) may be used if quantitation of gene products is desired.

[0133] An IL-21 and/or IL-21R polypeptide (or fragments thereof) may be detected using various well-known immunological assays employing the respective anti-IL-21 and/or IL-21R antibodies that may be generated as described herein. Immunological assays refer to assays that utilize an antibody (e.g., polyclonal, monoclonal, chimeric, humanized, scFv, and/or fragments thereof) that specifically binds to, e.g., an IL-21 polypeptide (or a fragment thereof). Such well-known immunological assays suitable for the practice of the present invention include ELISA, radioimmunoassay (RIA), immunoprecipitation, immunofluorescence, fluorescence-activated cell sorting (FACS), and Western blotting. The ordinarily skilled artisan will also recognize that an IL-21 polypeptide may also be detected using a labeled IL-21R polypeptide(s). One of skill in the art will understand that the aforementioned methods may be applied to disorders related to IL-21, especially fibrosis or a fibrosis-associated condition.

Assays for Measuring the Activity of IL-21 and/or IL-21R

[0134] The activity of IL-21/IL-21R (e.g., in response to IL-21/IL-21R antagonists as modulators of cytokine production and cell proliferation/differentiation) can be tested using any one of a number of routine factor-dependent cell proliferation assays for cell lines including, without

limitation, 32D, DA2, DA1G, T10, B9, B9/11, BaF3, MC9/G, M+(preB M+), 2E8, RB5, DA1, 123, T1165, HT2, CTLL2, TF-1, Mo7e and CMK.

[0135] Assays for T cell or thymocyte proliferation are described, for example, in *Current Protocols in Immunology*, Coligan et al. (eds.) Greene Pub. Assoc. & Wiley-Interscience, NY, NY (1991) (Chapter 3, "In Vitro assays for Mouse Lymphocyte Function" and Chapter 7, "Immunologic studies in Humans"); Takai et al. (1986) *J. Immunol.* 137:3494-500; Bertagnolli et al. (1990) *J. Immunol.* 145:1706-12; Bertagnolli et al. (1991) *Cell. Immunol.* 133:327-41; Bertagnolli et al. (1992) *J. Immunol.* 149:3778-83; Bowman et al. (1994) *J. Immunol.* 152:1756-61. Assays for cytokine production and/or proliferation of spleen cells, lymph node cells or thymocytes are described, for example, in "Polyclonal T cell stimulation" Kruisbeek and Shevach in *Current Protocols in Immunology*, Vol. 1 Coligan et al. (eds.) pp. 3.12.1-14, John Wiley and Sons, Toronto (1994); and "Measurement of mouse and human Interferon gamma" Schreiber, R.D. in *Current Protocols in Immunology*, Vol. 1 Coligan et al. (eds.) pp. 6.8.1-8, John Wiley and Sons, Toronto (1994).

[0136] Assays for proliferation and differentiation of hematopoietic and lymphopoietic cells are described in, for example, "Measurement of Human and Murine Interleukin 2 and Interleukin 4" Bottomly et al. in *Current Protocols in Immunology*, Vol. 1 Coligan et al. (eds.) pp. 6.3.1-12, John Wiley and Sons; Toronto (1991); deVries et al. (1991) *J. Exp. Med.* 173:1205-11; Moreau et al. (1988) *Nature* 336:690-92; Greenberger et al. (1983) *Proc. Natl. Acad. Sci. U.S.A.* 80:2931-38; "Measurement of mouse and human interleukin 6" Nordan, R. in *Current Protocols in Immunology*, Vol. 1 Coligan et al. (eds.) pp. 6.6.1-5, John Wiley and Sons, Toronto (1991); Smith et al. (1986) *Proc. Natl. Acad. Sci. U.S.A.* 83:1857-61; "Measurement of human Interleukin 11" Bennett et al. in *Current Protocols in Immunology*, Vol. 1 Coligan et al. (eds.) p. 6.15.1, John Wiley and Sons, Toronto (1991); "Measurement of mouse and human Interleukin 9" Ciarletta et al. in *Current Protocols in Immunology*, Vol. 1. Coligan et al. (eds.) p. 6.13.1, John Wiley and Sons, Toronto (1991).

[0137] Assays for T cell clone responses to antigens (which will identify, among others, proteins that affect APC-T cell interactions as well as direct T cell effects by measuring proliferation and cytokine production) include, for example, those described in: *Current Protocols in Immunology*, Coligan et al. (eds.) Greene Pub.

Assoc. and Wiley-Interscience, NY, NY (1991) (Chapter 3, "In Vitro assays for Mouse Lymphocyte Function"; Chapter 6, "Cytokines and their cellular receptors"; Chapter 7, "Immunologic studies in Humans"); Weinberger et al. (1980) *Proc. Natl. Acad. Sci. U.S.A.* 77:6091-95; Weinberger et al. (1981) *Eur. J. Immunol.* 11:405-11; Takai et al. (1986) *J. Immunol.* 137:3494-500; Takai et al. (1988) *J. Immunol.* 140:508-12.

Assays for Measuring the Interaction of IL-21 with IL-21R

[0138] Methods for detecting and/or measuring the level of interaction of IL-21 with IL-21R are well known in the art. For example, such interactions between a cytokine and its receptor may be detected and/or measured with, but not limited to, such techniques as ELISA, Western blotting, immunoprecipitation, Biacore analysis, etc.

Pharmaceutical Compositions

[0139] In one aspect, the invention features a method of treating, ameliorating, or preventing an IL-21-related disorder, i.e., fibrosis or a fibrosis-associated condition. The method may comprise contacting a population of cells with (e.g., by administering to a subject suffering from or at risk for fibrosis or a fibrosis-associated disorder) an agent that reduces the level of IL 21 and/or IL 21R activity in the subject, e.g., an IL-21/IL-21R antagonist (e.g., an anti-IL-21R antibody, an anti-IL-21 antibody, an antigen-binding fragment of an anti-IL-21R antibody, an antigen-binding fragment of an anti-IL-21 antibody, and a soluble fragment of an IL-21R (e.g., an amino acid sequence that is at least 90% identical to an amino acid sequence selected from the group consisting of amino acids 1-538 of SEQ ID NO:2, amino acids 20-538 of SEQ ID NO:2, amino acids 1-235 of SEQ ID NO:2, amino acids 20-235 of SEQ ID NO:2, amino acids 1-236 of SEQ ID NO: 2, amino acids 20-236 of SEQ ID NO:2, amino acids 1-529 of SEQ ID NO:5, amino acids 20-529 of SEQ ID NO:5, amino acids 1-236 of SEQ ID NO:5, and amino acid 20-236 of SEQ ID NO:5)) in an amount sufficient to inhibit the activity of IL-21 in the cell or population.

[0140] IL-21/IL-21R antagonists for treating fibrosis or a fibrosis-associated condition may be used as a pharmaceutical composition when combined with a pharmaceutically acceptable carrier. Such a composition may contain, in addition

to the IL-21/IL-21R-antagonists and carrier, various diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. The term “pharmaceutically acceptable” means a nontoxic material that does not interfere with the effectiveness of the biological activity of the active ingredient(s). The characteristics of the carrier will depend on the route of administration, and are generally well known in the art.

[0141] The pharmaceutical composition of the invention may be in the form of a liposome in which an IL-21/IL-21R-antagonist(s) is combined with, in addition to other pharmaceutically acceptable carriers, amphipathic agents such as lipids which exist in aggregated form as micelles, insoluble monolayers, liquid crystals, or lamellar layers which exist in aqueous solution. Suitable lipids for liposomal formulation include, without limitation, monoglycerides, diglycerides, sulfatides, lysolecithin, phospholipids, saponin, bile acids, and the like. Preparation of such liposomal formulations is within the level of skill in the art, as disclosed, e.g., in U.S. Patent Nos. 4,235,871, 4,501,728, 4,837,028, and 4,737,323, all of which are incorporated herein by reference in their entireties.

[0142] As used herein, the term “therapeutically effective amount” means the total amount of each active component of the pharmaceutical composition or method that is sufficient to show a meaningful subject benefit, e.g., amelioration or reduction of symptoms of, prevention of, healing of, or increase in rate of healing of such conditions. When applied to an individual active ingredient, administered alone, the term refers to that ingredient alone. When applied to a combination, the term refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination, serially or simultaneously.

[0143] In practicing the method of treatment or use of the present invention, a therapeutically effective amount of an IL-21/IL-21R antagonist is administered to a subject, e.g., a mammal (e.g., a human). An IL-21/IL-21R antagonist(s) may be administered in accordance with the method of the invention either alone or in combination with other therapies as described in more detail herein. When coadministered with one or more agents, an IL-21 and/or IL-21R antagonist may be administered either simultaneously with the second agent, or sequentially. If administered sequentially, the attending physician will decide on the appropriate

sequence of administering an IL-21/IL-21R antagonist in combination with other agents.

[0144] Administration of an IL-21/IL-21R antagonist used in a pharmaceutical composition of the present invention or to practice a method of the present invention may be carried out in a variety of conventional ways, such as oral ingestion, inhalation, or cutaneous, subcutaneous, or intravenous injection. Intravenous administration to the subject is sometimes preferred. When a therapeutically effective amount of an IL-21/IL-21R agonist or antagonist is administered orally, the binding agent will be in the form of a tablet, capsule, powder, solution or elixir. When administered in tablet form, the pharmaceutical composition of the invention may additionally contain a solid carrier such as a gelatin or an adjuvant. The tablet, capsule, and powder contain from about 5 to 95% binding agent, and preferably from about 25 to 90% binding agent. When administered in liquid form, a liquid carrier such as water, petroleum, oils of animal or plant origin such as peanut oil (albeit keeping in mind the frequency of peanut allergies in the population), mineral oil, soybean oil, or sesame oil, or synthetic oils may be added. The liquid form of the pharmaceutical composition may further contain physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol, propylene glycol or polyethylene glycol. When administered in liquid form, the pharmaceutical composition contains from about 0.5 to 90% by weight of the binding agent, and preferably from about 1 to 50% of the binding agent.

[0145] When a therapeutically effective amount of an IL-21/IL-21R antagonist is administered by intravenous, cutaneous or subcutaneous injection, the binding agent will be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable protein solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred pharmaceutical composition for intravenous, cutaneous, or subcutaneous injection should contain, in addition to a binding agent, an isotonic vehicle such as sodium chloride injection, Ringer's injection, dextrose injection, dextrose and sodium chloride injection, lactated Ringer's injection, or other vehicle as known in the art. The pharmaceutical composition of the present invention may also contain stabilizers, preservatives, buffers, antioxidants, or other additive known to those of skill in the art.

[0146] The amount of an IL-21/IL-21R antagonist in the pharmaceutical composition of the present invention will depend upon the nature and severity of the condition being treated, and on the nature of prior treatments that the subject has undergone. Ultimately, the attending physician will decide the amount of binding agent with which to treat each individual subject. Initially, the attending physician will administer low doses of binding agent and observe the subject's response. Larger doses of binding agent may be administered until the optimal therapeutic effect is obtained for the subject, and at that point the dosage is not generally increased further. It is contemplated that the various pharmaceutical compositions used to practice the method of the present invention should contain about 0.1 µg to about 100 mg IL-21/IL-21R antagonist per kg body weight.

[0147] The duration of intravenous therapy using the pharmaceutical composition of the present invention will vary, depending on the severity of the disease being treated and the condition and potential idiosyncratic response of each individual subject. It is contemplated that the duration of each application of an IL-21/IL-21R antagonist may be in the range of 12 to 24 hours of continuous i.v. administration. Also contemplated is subcutaneous (s.c.) therapy using a pharmaceutical composition of the present invention. These therapies can be administered daily, weekly, or, more preferably, biweekly, or monthly. It is also contemplated that where the IL-21/IL-21R antagonist is a small molecule (e.g., for oral delivery), the therapies may be administered daily, twice a day, three times a day, etc. Ultimately the attending physician will decide on the appropriate duration of i.v. or s.c. therapy, or therapy with a small molecule, and the timing of administration of the therapy using the pharmaceutical composition of the present invention.

[0148] The polynucleotide and proteins of the present invention are expected to exhibit one or more of the uses or biological activities (including those associated with assays cited herein) identified below. Uses or activities described for proteins, antibodies, or polynucleotides of the present invention may be provided by administration or use of such proteins, or antibodies, or by administration or use of polynucleotides encoding such proteins or antibodies (such as, for example, in gene therapies or vectors suitable for introduction of DNA).

Combination Therapy

[0149] In one embodiment, a pharmaceutical composition comprising at least one IL-21R/IL-21 antagonist, e.g., an IL-21R/IL-21 antibody, and at least one therapeutic agent is administered in combination therapy. Such therapy is useful for treating pathological conditions or disorders, such as immune and/or inflammatory disorders. The term “in combination” in this context means that the antagonist composition and the therapeutic agent are given substantially contemporaneously, either simultaneously or sequentially. If given sequentially, at the onset of administration of the second compound, the first of the two compounds may still be detectable at effective concentrations at the site of treatment.

[0150] For example, the combination therapy can include at least one IL-21R/IL-21 antagonist coformulated with, and/or coadministered with, at least one additional therapeutic agent. Additional agents may include at least one cytokine inhibitor, growth factor inhibitor, immunosuppressant, anti-inflammatory agent, metabolic inhibitor, enzyme inhibitor, cytotoxic agent, or cytostatic agent, as described in more detail below. Such combination therapies may advantageously utilize lower dosages of the administered therapeutic agents, thus avoiding possible toxicities or complications associated with the various monotherapies. Moreover, the therapeutic agents disclosed herein act on pathways that differ from the IL-21/IL-21R pathway, and thus are expected to enhance and/or synergize with the effects of the IL-21R/IL-21 antagonists.

[0151] Therapeutic agents used in combination with IL-21R/IL-21 antagonists may be those agents that interfere at different stages in the autoimmune and subsequent inflammatory response. In one embodiment, at least one IL-21R/IL-21 antagonist described herein may be coformulated with, and/or coadministered with, at least one cytokine and/or growth factor antagonist. The cytokine and/or growth factor antagonists may include soluble receptors, peptide inhibitors, small molecules, ligand fusions, antibodies (that bind cytokines or growth factors or their receptors or other cell surface molecules), and “anti-inflammatory cytokines” and agonists thereof.

[0152] Nonlimiting examples of the agents that can be used in combination with the IL-21R/IL-21 antagonists described herein, include, but are not limited to, antagonists of at least one interleukin (e.g., IL-1, IL-2, IL-6, IL-7, IL-8, IL-12,

IL-13, IL-15, IL-16, IL-17, IL-18, and IL-22); cytokine (e.g., TNF α , LT, EMAP-II, and GM-CSF); or growth factor (e.g., FGF and PDGF). The agents may also include, but are not limited to, antagonists of at least one receptor for an interleukin, cytokine, and growth factor. IL-21R/IL-21 antagonists can also be combined with inhibitors of, e.g., antibodies to, cell surface molecules such as CD2, CD3, CD4, CD8, CD20 (e.g., the CD20 inhibitor rituximab (RITUXAN[®])), CD25, CD28, CD30, CD40, CD45, CD69, CD80 (B7.1), CD86 (B7.2), CD90, or their ligands, including CD154 (gp39 or CD40L), or LFA-1/ICAM-1 and VLA-4/VCAM-1 (Yusuf-Makagiansar et al. (2002) *Med. Res. Rev.* 22:146-67). Other compounds that can be used in combination with IL-21R/IL-21 antagonists described herein may include antagonists of the receptors for IL-1, IL-12, TNF α , IL-15, IL-17, IL-18 and IL-22.

[0153] Examples of agents useful in combination therapies with an IL-21R/IL-21 antagonist include IL-12 antagonists (such as antibodies that bind IL-12 (see e.g., WO 00/56772); IL-12 receptor inhibitors (such as antibodies to the IL-12 receptor); and soluble IL-12 receptor and fragments thereof. Examples of IL-15 antagonists include antibodies against IL-15 or its receptor, soluble fragments of the IL-15 receptor, and IL-15-binding proteins. Examples of IL-18 antagonists include antibodies to IL-18, soluble fragments of the IL-18 receptor, and IL-18 binding proteins (IL-18BP, Mallat et al. (2001) *Circ. Res.* 89:E41-45). Examples of IL-1 antagonists include interleukin-1-converting enzyme (ICE) inhibitors (such as Vx740), IL-1 antagonists (e.g., IL-1RA (anakinra (KINERET[™]), Amgen)), sIL-1RII (Immunex); and anti-IL-1 receptor antibodies.

[0154] Examples of TNF antagonists include antibodies to TNF (e.g., human TNF α), such as D2E7 (human anti-TNF α antibody, U.S. Patent No. 6,258,562, HUMIRA[™], Abbott Labs); CDP-571/CDP-870/BAY-10-3356 (humanized anti-TNF α antibodies, Celltech/Pharmacia); cA2 (chimeric anti-TNF α antibody, REMICADE[™], Centocor); and anti-TNF antibody fragments (e.g., CPD870). Other examples include soluble TNF receptor (e.g., human p55 or p75) fragments and derivatives, such as p55 kDTNFR-IgG (55 kD TNF receptor-IgG fusion protein, LENERCEPT[™]) and 75 kDTNFR-IgG (75 kD TNF receptor-IgG fusion protein, ENBRELE[™] (etanercept - Immunex)). See, e.g., van der Poll et al. (1997) *Blood*. 89:3727-34; Mori et al. (1996) *J. Immunol.* 157:3178-82. Further

examples include enzyme antagonists (e.g., $\text{TNF}\alpha$ converting enzyme inhibitors (TACE) such as alpha-sulfonyl hydroxamic acid derivative (WO 01/55112) or N-hydroxyformamide inhibitors (GW 3333, -005, or -022, GlaxoSmithKline) and TNF-bp/s-TNFR (soluble TNF binding protein, see, e.g., Lantz et al. (1991) *J Clin Invest.* 88:2026-31; Kapadia et al. (1995) *Amer. J. Physiol. Heart Circ. Phys.* 268:H517-25). TNF antagonists may be soluble TNF receptor (e.g., human p55 or p75) fragments and derivatives, such as 75 kDTNFR-IgG; and $\text{TNF}\alpha$ converting enzyme (TACE) inhibitors.

[0155] In other embodiments, the IL-21R/IL-21 antagonists described herein can be administered in combination with at least one of the following: IL-13 antagonists, such as soluble IL-13 receptors and/or anti-IL-13 antibodies; and IL-2 antagonists, such as IL-2 fusion proteins (e.g., DAB 486-IL-2 and/or DAB 389-IL-2 made by Seragen, see e.g., Sewell et al. (1993) *Arthritis Rheum.* 36:1223-33) and anti-IL-2R antibodies (e.g., anti-Tac (humanized antibody, Protein Design Labs, see Junghans et al. (1990) *Cancer Res.* 50:1495-502). Another combination includes IL-21R/IL-21 antagonists in combination with nondepleting anti-CD4 inhibitors such as IDEC-CE9.1/SB 210396 (anti-CD4 antibody, GlaxoSmithKline). Yet other combinations include IL-21R/IL-21 antagonists with CD80 (B7.1) and CD86 (B7.2) costimulatory pathway antagonists (such as antibodies, soluble receptors, or antagonistic ligands); P-selectin glycoprotein ligand (PSGL) and PSGL-1 inhibitors (such as antibodies to PSGL and/or PSGL-1 and small molecule inhibitors); T cell- and B cell-depleting agents (such as anti-CD4 or anti-CD22 antibodies), and anti-inflammatory cytokines and agonists thereof (e.g., antibodies). The anti-inflammatory cytokines may include IL-4 (e.g., Schering-Plough Biopharma); IL-10 (e.g., SCH 52000, recombinant IL-10, Schering-Plough Biopharma); IL-11; IL-13; and $\text{TGF}\beta$ or agonists thereof (e.g., agonist antibodies).

[0156] In other embodiments, at least one IL-21R/IL-21 antagonist can be coformulated with, and/or coadministered with, at least one anti-inflammatory drug, immunosuppressant, metabolic inhibitor, and enzymatic inhibitor. Nonlimiting examples of the drugs or inhibitors that can be used in combination with the IL-21R/IL-21 antagonists described herein, include, but are not limited to, at least one of: nonsteroidal anti-inflammatory drugs (NSAIDs) (including,

but not limited to, aspirin, salsalate, diflunisal, ibuprofen, ketoprofen, nabumetone, piroxicam, naproxen, diclofenac, indomethacin, sulindac, tolmetin, etodolac, ketorolac, oxaprozin, tenidap, meloxicam, piroxicam, aceclofenac, tolmetin, tiaprofenic acid, nimesulide, etc.); sulfasalazine; corticosteroids (such as prednisolone); cytokine suppressive anti-inflammatory drugs (CSAID); inhibitors of nucleotide biosynthesis (such as inhibitors of purine biosynthesis (e.g., folate antagonist such as methotrexate)); and inhibitors of pyrimidine biosynthesis, e.g., a dihydroorotate dehydrogenase (DHODH) inhibitor such as leflunomide (see, e.g., Kraan et al. (2004) *Ann. Rheum. Dis.* 63:1056-61). Therapeutic agents for use in combination with IL-21/IL-21R antagonists may include one or more NSAIDs, CSAIDs, DHODH inhibitors (such as leflunomide), and folate antagonists (such as methotrexate).

[0157] Examples of additional agents that may be used in combination with IL-21/IL-21R antagonists include at least one of: corticosteroid (oral, inhaled and local injection); immunosuppressant (such as cyclosporin and tacrolimus (FK-506)); an mTOR inhibitor (such as sirolimus (rapamycin) or a rapamycin analog and/or derivative, e.g., ester rapamycin derivative such as CCI-779 (see, e.g., Elit (2002) *Curr. Opin. Investig. Drugs* 3:1249-53; Huang et al. (2002) *Curr. Opin. Investig. Drugs* 3:295-304)); an agent which interferes with the signaling of proinflammatory cytokines such as TNF α and IL-1 (e.g., an IRAK, NIK, IKK, p38 or MAP kinase inhibitor); TPL-2, Mk-2 and NF κ b inhibitors; COX-2 inhibitors (e.g., celecoxib, rofecoxib, etc., and variants thereof); phosphodiesterase inhibitors (such as Rolipram); phospholipase inhibitors (e.g., an inhibitor of cytosolic phospholipase 2 (cPLA2) such as trifluoromethyl ketone analogs (U.S. Patent No. 6,350,892)); inhibitors of vascular endothelial cell growth factor (VEGF); inhibitors of the VEGF receptor; inhibitors of angiogenesis; RAGE and soluble RAGE; estrogen receptor beta (ERB) agonists, ERB-NF κ b antagonists; interferon- β (for example, IFN β -1a and IFN β -1b); copaxone; and corticosteroids.

[0158] Other useful therapeutic agents that may be combined with an IL-21R/IL-21 antagonist include: budenoside; epidermal growth factor; aminosalicylates; 6-mercaptopurine; azathioprine; metronidazole; lipoxigenase inhibitors; mesalamine; olsalazine; balsalazide; antioxidants; thromboxane

inhibitors; growth factors; elastase inhibitors; pyridinyl-imidazole compounds; glucuronide- or dextran-conjugated prodrugs of prednisolone; dexamethasone or budesonide; ICAM-1 antisense phosphorothioate oligodeoxynucleotides (ISIS 2302; Isis Pharmaceuticals, Inc.); soluble complement receptor 1 (TP10; T Cell Sciences, Inc.); slow-release mesalazine; antagonists of platelet activating factor (PAF); ciprofloxacin; lignocaine; cyclosporin A; hydroxychloroquine (PLAQUENIL™); minocycline (MINOCIN™); and anakinra (KINERET™).

[0159] Choosing a particular therapeutic agent for administration in combination with an IL-21/IL-21R antagonist of the invention will largely depend on factors such as the particular subject, the desired target, and chosen length of treatment. Such decisions are well within the skill and knowledge of one skilled in the art.

[0160] Additional examples of therapeutic agents that can be combined with an IL-21R/IL-21 antagonist include one or more of: 6-mercaptopurines (6-MP); azathioprine; sulphasalazine; mesalazine; olsalazine; chloroquine, hydroxychloroquine (PLAQUENIL®); pencillamine; aurothiomalate (intramuscular and oral); azathioprine; colchicine; beta-2 adrenoreceptor agonists (salbutamol, terbutaline, salmeterol); xanthines (theophylline, aminophylline); cromoglycate; nedocromil; ketotifen; ipratropium and oxitropium; mycophenolate mofetil; adenosine agonists; antithrombotic agents; complement inhibitors; and adrenergic agents.

[0161] In one embodiment, an IL-21R/IL-21 antagonist can be used in combination with one or more antibodies directed at other targets involved in regulating immune responses. Nonlimiting examples of agents for treating or preventing immune responses with which an IL-21R/IL-21 antagonist of the invention can be combined include the following: antibodies against other cell surface molecules, including but not limited to CD25 (interleukin-2 receptor- α), CD11a (LFA-1), CD54 (ICAM-1), CD4, CD45, CD28, CTLA4, ICOSL, ICOS, CD80 (B7.1), and/or CD86 (B7.2). In yet another embodiment, an IL-21R/IL-21 antagonist is used in combination with one or more general immunosuppressive agents, such as cyclosporine A or FK506. In another embodiment, an IL-21/IL-21R antagonist is used in combination with a CTLA4 agonist, e.g., (e.g., CTLA4 Ig – abatacept (ORENCIA®)).

[0162] The entire contents of all references, patents, and published patent applications cited throughout this application are hereby incorporated by reference herein.

EXAMPLES

[0163] The following Examples provide illustrative embodiments of the invention and do not in any way limit the invention. One of ordinary skill in the art will recognize that numerous other embodiments are encompassed within the scope of the invention.

Example 1: Materials and Methods

Example 1.1: Mice, Parasite Infections and Antigen Preparation

[0164] Female or male C57BL/6, C57BL/6/Ai-IL-10KO/IL-4KO mice and C57BL/6Ai-IL-10KO/IL-12KO were obtained from Taconic Farms (Germantown, NY) (Hoffmann et al. (1999) *J. Immunol.* 163:927-938). Breeding pairs of IL-21R^{-/-} mice on a C57BL/6 background were obtained from a breeding colony housed at Harvard School of Public Health (Boston, MA) (Kasaian et al. (2002) *Immunity* 16:559-69). All mice were housed under specific pathogen-free conditions at the National Institutes of Health in an American Association for the Accreditation of Laboratory Animal Care-approved facility. The NIAID animal care and use committee approved all experimental procedures. *S. mansoni* eggs were extracted from the livers of infected mice (Biomedical Research Institute, Rockville, MD) as previously described (Wynn et al. (1995) *Nature* 376:594-96). For the induction of synchronous primary pulmonary granulomas, mice were given 5,000 eggs intravenously (i.v.). For the induction of secondary granulomas, mice were sensitized intraperitoneally (i.p.) with 5000 live eggs, and then challenged with 5,000 live eggs i.v. (Wynn et al. (1994) *J. Exp. Med.* 179:551-61). In the infection experiments, mice were infected percutaneously via the tail with 25-30 cercariae of a Puerto-Rican Strain of *S. mansoni* (NMRI) that were obtained from infected *Biomphalaria glabrata* snails (Biomedical Research Institute, Rockville, MD). Soluble egg antigen (SEA) and soluble worm antigenic preparations (SWAP) were from purified and homogenized from *S. mansoni* eggs and adult parasites as previously described (Cheever et al. (1994) *J. Immunol.* 153:753-59). All animals underwent perfusion at the time of

sacrifice so that worm and tissue egg burdens could be determined, as described elsewhere (*id.*). *Nippostrongylus brasiliensis* larvae (L3) were prepared as previously described (Katona et al. (1983) *J. Immunol.* 130:350-56). Mice were inoculated through s.c. injection of 500 L3. On day seven post-inoculation, lung tissue and mediastinal lymph nodes were collected for cytokine analysis.

Example 1.2: Histopathology and Fibrosis

[0165] The sizes of pulmonary and hepatic granulomas were determined on histological sections that were stained with Wright's Giemsa stain (Histopath of America, Clinton, MD). Around 30 granulomas per mouse were included in all analyses. A skilled pathologist evaluated the percentages of eosinophils, mast cells and other types of cells in the same sections. The number of schistosome eggs in the liver and the gut and the collagen content of the liver, as measured by hydroxyproline levels, were determined as previously described (Cheever et al., *supra*). Specifically, hepatic collagen was measured as hydroxyproline by the technique of Bergman and Loxley (Bergman and Loxley (1963) *Analytical Biochem.* 35:1961-65) after hydrolysis of a 200-mg portion of liver in 5 ml of 6N HCl at 110°C for 18 h. The increase in hepatic hydroxyproline was positively related to egg numbers in all experiments and hepatic collagen is reported as the increase above normal liver collagen in micromoles per 10,000 eggs; (infected liver collagen – normal liver collagen)/liver eggs $\times 10^{-4}$ or micromoles per worm pair. At late chronic time points, fibrosis is reported as total liver collagen per liver. The same individual scored all histological features and had no knowledge of the experimental design.

Example 1.3: FACS Analysis

[0166] Whole lungs were harvested and placed in RPMI. Tissues were disrupted by straining through a 70-micron nylon mesh (BD Falcon, San Diego, CA). The single cell suspensions were washed and RBCs were lysed by incubation with ACK lysis solution for 3 min. Lung lymphocytes were labeled with PE-Cy5 labeled anti-CD4 along with Fc Block (both antibodies from BD Pharmingen, San Diego, CA) in FACS buffer for 15 min at 4°C. After washing, the cells were analyzed on a FACS Calibur using FLOWJO™ software (Treestar, Inc., Ashland, OR).

Example 1.4: IL-21 Blocking Experiments with sIL-21R-Fc

[0167] C57BL/6 (10/group) mice were infected percutaneously via the tail with 30-35 *S. mansoni* cercariae. Beginning on week 6 post-infection, mice were treated with either mIL-21R-Fc (Wyeth Research) or Anti-*E. tenella* murine IgG2a control antibody (Wyeth Research). Each mouse received one 200 µg dose via i.p. injection 3X/week for a total of 5 weeks. Mice were sacrificed 12 weeks post-infection and hepatic fibrosis was measured by hydroxyproline assay.

Example 1.5: Lymphocyte Culture and Cytokine Detection Using the Enzyme-linked Immunosorbent Assay (ELISA)

[0168] Spleen and mesenteric lymph nodes (infection model) or lung-associated lymph nodes (pulmonary model) were removed aseptically and single cell suspensions were prepared as previously described (Hesse et al. (2000) *Am. J. Pathol.* 157:945-55). Cultures were incubated at 37°C in a humidified atmosphere of 5% CO₂. Cells were stimulated with SEA (20 µg/ml), SWAP (50 µg/ml), concanavalin A (Con A; 1 µg/ml), or medium alone. Supernatant fluids were harvested at 72 hours and assayed for cytokine production. IFN-γ, IL-5 and IL-10 were measured by sandwich ELISA using paired antibodies (BD Pharmingen, San Diego, CA) as previously described (*id.*). Cytokine levels were calculated with standard curves constructed using recombinant murine cytokines (BD Pharmingen, San Diego, CA). IL-13 levels were measured using murine IL-13 ELISA kits (R&D Systems, Minneapolis, MN) according to the manufacturer's protocol. TGF-β1 levels were quantified using mouse TGF-β1 DUOSET® ELISA development system (R&D Systems, Minneapolis, MN) according to manufacture's protocol. To avoid bovine-derived TGF-β1 contamination, cells were washed 3X in PBS and cultured in media containing 0.5% mouse serum.

Example 1.6: RNA Isolation and Purification and Real-Time Polymerase Chain Reaction

[0169] Total RNA was extracted from lung and liver tissue samples placed individually in 1 ml TRIZOL™ reagent (Invitrogen, Carlsbad, CA). The sample was homogenized using a tissue polytron (Omni International Inc., Marietta, GA) and total RNA was extracted according to the recommendations of the

manufacturer and further purified using RNEASY™ Mini Kit from Qiagen (Qiagen Sciences, Germantown, MD). Individual sample RNA (1 µg) was reverse-transcribed using SUPERScript II™ (Invitrogen, Carlsbad, CA) and a mixture of oligo (dT) and random primers. Real-time polymerase chain reaction (RT-PCR) was performed on an ABI PRISM™ 7900 sequence detection system (Applied Biosystems, Foster City, CA). Relative quantities of mRNA for several genes was determined using SYBR™ Green PCR Master Mix (Applied Biosystems, Foster City, CA) and by the comparative threshold cycle method as described by Applied Biosystems for the ABI PRISM™ 7700/7900 sequence detection systems (Applied Biosystems, Foster City, CA). In this method, mRNA levels for each sample were normalized to hypoxanthine guanine phosphoribosyl transferase mRNA levels and then expressed as a relative increase or decrease compared with levels in uninfected controls. Primers were designed using PRIMER EXPRESS™ software (Applied Biosystems, Foster City, CA). Primers for IL-13, IL-4, IL-10, HPRT (Hesse et al. (2001) *J. Immunol.* 167:6533-44), IL-13Rα2 (Chiaramonte et al. (2003) *J. Exp. Med.* 197:687-701), Ym1, FIZZ1 and acidic chitinase (AMCase) (Sandler et al., *supra*) were published previously, and include:

[0170] IL-21

5' GCCAG ATCGC CTCCT GATTA 3' (sense) (SEQ ID NO:28);

5' CATGC TCACA GTGCC CCTTT 3' (antisense) (SEQ ID NO:29);

[0171] IL-21R

5' CTCCC CCCTT GAACG TGA CT 3' (sense) (SEQ ID NO:30);

5' TTGCC CCTCA GCACG TAGTT 3' (antisense) (SEQ ID NO:31);

[0172] IFN-γ

5' AGAGC CAGAT TATCT CTTTC TACCT CAG 3' (sense) (SEQ ID NO:32);

5' CCTTT TTCGC CTTGC TGTTG 3' (antisense) (SEQ ID NO:33).

Example 1.7: Serum Antibody Isotype Analysis and Bone Marrow-Derived Macrophages

[0173] Total IgE was measured using the BD OPTeia™ mouse IgE ELISA Set (BD Biosciences Pharmingen, San Diego, CA) according to the manufacturer's protocol. SEA-specific IgG1 and IgG2b isotype-specific antibody (Ab) titers were evaluated by indirect ELISA. IMMULON™ 4 plates (Thermo Labsystems Inc., Beverly MA) were coated with 10 µg/ml SEA (100 µl/well) diluted in PBS, and serum samples were analyzed using serial two-fold dilutions. Biotin-Rabbit Anti-mouse IgG1 (Zymed, San Francisco, CA) was used at a 1:1000 dilution. This was followed by peroxidase-labeled streptavidin (KPL, Gaithersburg, MD) substrate enzyme at a 1:1000 dilution. Second-step horseradish peroxidase-conjugated rabbit anti-mouse IgG2b (Zymed, San Francisco, CA) Ab was used at a 1:1000 dilution. The absorbance in the wells was read at 405 nm using a VMAX™ Kinetic Microplate Reader (Molecular Devices) after adding 100 µl one-component ABTS Peroxidase Substrate (KPL, Gaithersburg, MD).

[0174] Bone marrow was recovered from female C57BL/6 mice and cultured in Petri dishes (100 X 15 mm) containing supplemented DMEM media (L929-conditioned medium) for a period of 6 days. After six days, cells were harvested and seeded at a concentration of 0.5×10^6 cells/well in 24 well plates containing supplemented DMEM media (10% FBS, 2 mM L-glutamine, 100 U/mL penicillin, and 100 µg/mL streptomycin). Cells were stimulated with IL-4, IL-13, and IL-21 (R&D, Minneapolis, MN) for a period of 20 hours. In some assays, cells were pretreated with IL-21. Cells were lysed and RNA was purified using the RNA Cleanup procedure with the RNEASY™ kits (Qiagen Sciences, Germantown, MD).

Example 1.8: Arginase Activity Assay

[0175] Bone marrow-derived macrophages were plated at 6×10^5 per well in 96 well tissue culture plates and stimulated with combinations of IL-4, IL-13, and IL-21. IL-21 was added 6 hours prior to IL-4 or IL-13 stimulation. Following stimulation, cells were washed with PBS and lysed with 0.1% Triton X-100 containing protease inhibitor (Roche, Nutley, NJ). Lysates were transferred into a 96 well PCR plate and incubated with 10 mM MnCl₂ and 50mM Tris HCl (pH 7.5) to activate enzyme for 10min at 55°C. After enzyme activation, 25 µl of

lysate was removed and added to 25 μ l 1M arginine (pH 9.7) in a new PCR plate and incubated for 20 hours at 37°C. 5 μ l of each sample was added in duplicate to a 96 well ELISA plate along with 5 μ l of each standard, diluted in the same assay conditions, starting at 100 mg/dL. The urea determination reagent from QUANTICHRON™ Urea Assay Kit (BioAssay Systems, Hayward, CA) was used according to the manufacture's protocol.

Example 1.9: Statistics

[0176] Hepatic fibrosis (adjusted for egg number) decreases with increasing intensity of infection (worm pairs). Therefore, these variables were compared by analysis of covariance, using the logarithm of total liver eggs as the covariate and the logarithm of hydroxyproline content per egg. Variables that did not change with infection intensity were compared by one-way ANOVA or Student's *t* test (Cheever et al., *supra*). Changes in cytokine mRNA expression and granuloma size were evaluated using ANOVA. Differences were considered significant when $p < 0.05^*$, $p < 0.01^{**}$, or $p < 0.001^{***}$.

Example 2: Regulation of IL-21 and IL-21R during Type-1- and Type-2-Polarized Responses

[0177] To investigate the regulation and function of the IL-21 receptor *in vivo*, several different experimental systems of T_H2-dependent inflammation were examined, including models of pulmonary and hepatic inflammation, as well as an experimental model of nematode infection (Pearce et al., *supra*; Wynn et al. (1994), *supra*). In each case, the immune responses of wild type (WT) animals were compared with IL-21R-deficient mice (Hoffman et al., *supra*; Kasaian et al. (2002), *supra*).

[0178] In comparison to IL-21 (Wurster et al. (2002), *supra*; Mehta et al. (2004) *Immunol. Rev.* 202:84-95), little is known about the regulation and function of the IL-21 receptor. To determine whether IL-21 and its receptor are regulated during a pathological T_H2 response *in vivo*, the *S. mansoni* model of granuloma formation was used. In this model, T_H2 cytokines are known to play a prominent role in lesion formation (Pearce and MacDonald, *supra*). Initial studies were designed to determine whether IL-21 and IL-21R mRNA expression were linked with polarized T_H2 cytokine responses *in vivo*. This was achieved by using mice

that develop highly exaggerated T_H1 (IL-4^{-/-}/IL-10^{-/-}) or T_H2 (IL-12^{-/-}/IL-10^{-/-}) cytokine responses following exposure to *S. mansoni* eggs. In IL-4^{-/-}/IL-10^{-/-} “ T_H1 ” mice, IFN- γ mRNA expression increased in the lung 75-fold over baseline by day 4 post-challenge and remained approximately 50-fold above background through day 14 (FIG. 16A). IL-13 mRNA was not detectable in these mice at any time point, confirming the establishment of a highly polarized T_H1 inflammatory response. In contrast, the IL-12^{-/-}/IL-10^{-/-} “ T_H2 ” mice displayed a 200 to 250-fold increase in IL-13 mRNA at all time points post-challenge, with little to no change in IFN- γ . In contrast to the T_H1 / T_H2 cytokines, which displayed a highly polarized pattern of expression, IL-21 was not associated with a polarized phenotype (FIG. 16B). In both groups, IL-21 mRNA levels increased at least 50-fold over baseline following challenge with schistosome eggs, although the increase observed in the T_H1 -polarized mice was on average 3- to 4-fold greater than the T_H2 polarized animals (FIG. 16B; lower panel). IL-21R was also not specifically associated with a T_H1 or T_H2 immune response. However, in contrast to IL-21, which was more pronounced in T_H1 skewed animals, the maximal response for the IL-21R was observed in the T_H2 -polarized mice (FIG. 16B; upper panel).

Example 3: Type-2 Cytokine Production is Reduced in the Lungs of IL-21R-deficient Mice During a Primary Response to Schistosome Eggs

[0179] Given the significant elevation in IL-21R expression in mice challenged with schistosome eggs (FIG. 16), the next series of experiments examined whether IL-21R signaling was influencing the development of the T_H2 response. In these experiments, naïve WT and IL-21R^{-/-} mice were injected intravenously with live schistosome eggs and the production of T_H2 cytokines and T_H2 -regulated genes were monitored in the lung, spleen, and draining lymph nodes over the following 14 days. In WT mice, IL-21R mRNA expression increased rapidly following egg exposure and remained elevated through day 14 (FIG. 17A). IL-21 showed a similar profile with peak expression occurring on day 7 and then declining slightly thereafter. Notably, there was a consistent and highly significant decrease in IL-21 expression on days 7 and 14 in the IL-21R^{-/-} mice, suggesting that IL-21R was positively influencing the expression of its own ligand. Consistent with previous observations (Wynn et al. (1993) *J. Immunol.*

151:1430-40; Vella and Pearce (1992) *J. Immunol.* 148:2283-88), expression of the T_H2-associated cytokines IL-4 and IL-13 rose gradually in the granulomatous tissues of WT mice, with 5- to 15-fold increases detectable by day 14. In contrast, there was a marked and significant decrease in IL-4 and IL-13 mRNA expression in the IL-21R^{-/-} tissues. Although little change in IFN- γ and IL-10 mRNA was detected in WT mice between day 4 and 14 post-challenge, production of IFN- γ and IL-10 also decreased slightly in the IL-21R^{-/-} animals. Thus, the reduced T_H2 response observed in the IL-21R^{-/-} mice was not associated with increased T_H1 cytokine production. The decrease in T_H2 cytokines was also specific to the granulomatous tissues, since significant T_H2 cytokine production was observed in lymph node and splenocyte cultures following *in vitro* stimulation with soluble egg antigen (SEA) or mitogen (FIG. 17B). In fact, SEA consistently stimulated stronger IL-5, IL-10, and IL-13 responses in the lymph node cultures prepared from IL-21R^{-/-} mice. Nevertheless, consistent with the reduced T_H2 response in the lung, a more rapid resolution of granuloma formation was observed in the IL-21R^{-/-} animals (FIG. 17C). In addition, there was a marked decrease in several genes associated with Stat6-activation or “alternatively-activated macrophages” (AAM ϕ) (Nair et al. (2005) *Infect. Immun.* 73:385-94; Zhu et al. (2004) *Science* 304:1678-82; Chiaramonte et al. (2003), *supra*; Gordon, S. (2003) *Nat. Rev. Immunol.* 3:23-35), providing further evidence of an overall reduction in the T_H2 effector response in IL-21R^{-/-} mice (FIG. 17D).

Example 4: TH2 Response is Reduced in IL-21R^{-/-} Mice Following *N. brasiliensis* Infection

[0180] To determine if the reduced T_H2 effector response was specific to *S. mansoni* pulmonary granuloma formation, WT and IL-21R^{-/-} mice were infected with the intestinal nematode *N. brasiliensis*. Infection is established by inoculating third stage larvae (L3) under the skin. As the parasites mature, they migrate from the site of inoculation and enter the lungs via the circulatory system. Once inside the lungs, the parasites trigger a vigorous and highly polarized T_H2 response (Urban et al. (1993) *J. Immunol.* 151:7086-94), which was confirmed by analyzing the expression of several T_H2-associated genes in the lung (FIG. 18A) and lung-associated lymph nodes (FIG. 18B). The lungs and

lymph nodes of WT mice displayed marked increases in IL-4, IL-13, AMCase, FIZZ1/RELM1 α , and Ym1 mRNA expression following *N. brasiliensis* infection (FIG. 18A and 18B). However, in agreement with the pulmonary granuloma model, significantly reduced levels of IL-4, IL-13, and AMCase were observed, as well as slightly reduced levels of Ym1 and FIZZ1 mRNA in the lungs of the IL-21R^{-/-} mice (FIG. 18A). The draining lymph nodes displayed a similar reduction, although the decreases in Ym1 and FIZZ1 were more significant in the lymph nodes (FIG. 18B). The only other major difference between the two tissues was the AMCase mRNA response, which appeared to be restricted to the lung. Together, these data confirm an important role for the IL-21R in T_H2 response development *in vivo*. Notably however, despite developing a markedly attenuated T_H2 response, the *N. brasiliensis* infected IL-21R^{-/-} mice displayed no significant delay in adult worm expulsion (not shown).

Example 5: Type-2 Cytokine-driven Inflammation is Diminished in the Lungs of IL-21R^{-/-} Mice

[0181] The next series of experiments were designed to determine whether the IL-21R modulates the development of secondary T_H2 responses. For these experiments, WT and IL-21R^{-/-} mice were sensitized with *S. mansoni* eggs and challenged intravenously 2 weeks later. As expected, the sensitized mice developed a robust granulomatous response that was 4- to 5-times greater (FIG. 19C) than the primary challenge animals (FIG. 17C). As observed in the primary model, there was a significant increase in IL-21 and IL-21R mRNA in the lungs following egg exposure, although the IL-21 response peaked much earlier during the secondary challenge. IL-21R was only modestly increased when compared with IL-21 although it remained significantly elevated at both time points, while IL-21 mRNA levels declined after reaching a peak on day 4 (FIG. 19A). Thus, there was evidence of tighter regulation of the ligand in the tissues. There was also a remarkable decrease in IL-21 expression in the IL-21R^{-/-} mice, confirming a potent feedback mechanism between the receptor and its ligand. Among the T_H2-associated cytokines, IL-13 was the most robust response, displaying a 50- to 100-fold increase over baseline in WT mice. However, it was reduced to 10- to 20-fold above background in the IL-21R^{-/-} mice, demonstrating that the IL-21R is required for maximum development of the secondary T_H2 response. Again, the

reduction in T_H2 cytokine expression in IL-21R^{-/-} mice was not accompanied by a significant increase in IFN- γ . In fact, IFN- γ mRNA expression decreased in the lungs of the IL-21R^{-/-} mice. Nevertheless, the knockouts displayed a modest but consistent increase in IFN- γ production in the lymph nodes and spleen, suggesting a greater inhibition of the T_H2 cytokines overall (FIG. 19B). Consistent with the primary egg challenge model, the reduction in T_H2 and T_H1 cytokine production was more pronounced in the granulomatous tissues (FIG. 19A), although the SEA-induced T_H2 response was also partially reduced in the spleen (FIG. 19B). The significant reduction in secondary granulomatous inflammation was consistent with the development of a weaker T_H2 response in the lung (FIG. 19C). In addition, there was a marked decrease in FIZZ1, Ym1, and AMCase expression (FIG. 19D), further confirming a significant impairment of secondary T_H2 effector responses in the IL-21R^{-/-} mice.

Example 6: IgG Antibodies, Granuloma Formation, and Type-2 Cytokines Are Substantially Reduced in Infected IL-21R-deficient Mice

[0182] Next, to determine if IL-21 signaling is required for the maintenance of a chronic T_H2 -dominated response, animals were exposed percutaneously to *S. mansoni* cercariae, and their pathological reactions and immune responses at both acute and chronic time points post-infection were analyzed. As observed in the pulmonary granuloma studies, there was a marked upregulation in IL-21R and IL-21 mRNA expression in the livers of infected WT mice. In contrast, IL-21 mRNA was almost undetectable in the IL-21R^{-/-} mice even after chronic infection (FIG. 20A). At the acute stage post-infection, the IL-21R^{-/-} mice also manifested a marked reduction in T_H2 cytokine mRNA expression (FIG. 20A). However, the changes were again restricted to the granulomatous tissues because the lymph node and splenocyte responses of both groups were similar following *in vitro* stimulation with parasite antigens (FIG. 20B). The only consistent difference noted in the *in vitro* assays was a 2- to 3-fold decrease in IL-5 and IL-10 production in the splenocyte cultures. The IL-21R^{-/-} mice also developed significantly smaller granulomas at the acute stage post-infection (FIG. 20C), which was consistent with the reduced IL-4 and IL-13 mRNA responses in the liver (FIG. 20A). However, this was not accompanied by any obvious change in the percentage of eosinophils in the granulomas (FIG. 20C). A more detailed

microscopic analysis of the lesions confirmed that there was no detectable change in the overall composition of the granulomas (FIG. 21A). Experiments were also undertaken to determine whether IL-21R-deficiency was specifically affecting the recruitment of CD4⁺ T cells to the granulomatous tissues. To address this issue, the pulmonary granuloma model was used in order to synchronize the recruitment of inflammatory cells. However, consistent with the microscopic evaluations of liver granulomas (FIG. 21A), the percentage of CD4⁺ T cells in the lungs was similar in WT and IL-21R^{-/-} mice both before and after egg exposure (FIG. 21B). Thus, changes in CD4⁺ T cell recruitment or expansion are unlikely to explain the decreased Th1/Th2 cytokine responses observed in the tissues. Instead, they appear to result from a more general reduction in the overall inflammatory response. Importantly, both groups effectively downmodulated their granulomatous responses by week 12 post-infection (Pearce and MacDonald, *supra*). Consequently, there was no significant difference in granuloma size at the chronic time point (FIG. 20C). Minimal impairment in the T_H2 cytokine response was observed in the chronically infected knockout mice (FIG. 20A). The marked reduction in FIZZ1 and Ym1 observed at the acute stage had also diminished in the chronically infected IL-21R^{-/-} animals (FIG. 20D). Nevertheless, expression of AMCase remained remarkably low on week 12, suggesting a sustained diminution of at least a subset of the T_H2-driven responses in chronically infected IL-21R^{-/-} mice.

[0183] The IL-21R^{-/-} mice were also examined for changes in serum antibody levels (FIG. 22). Consistent with their suppressed cytokine responses (FIG. 20A), the IL-21R^{-/-} mice displayed a marked reduction in parasite specific IgG₁ (T_H2-associated antibody) and IgG_{2b} (T_H1-associated antibody) titers, which was maintained at the chronic time point (FIG. 22B). Interestingly however, this was not accompanied by any significant change in IgE (FIG. 22C), suggesting a selective impairment in only a subset of serum antibody isotypes. Exogenous IL-21 has been shown to inhibit IgE production (Suto et al. (2002) *Blood* 100:4565-73), which may explain the slight elevation of IgE in the chronically infected IL-21R^{-/-} mice. Importantly, the overall reduction in type-2 responsiveness in the IL-21R^{-/-} mice was not attributed to differences in parasite burden since similar numbers of eggs and paired adult parasites were found in the tissues of both groups at all time points (FIG. 22A).

Example 7: IL-21R-deficiency Slows the Progression of Hepatic Fibrosis

[0184] Because T_H2 cytokines are believed to play a major role in tissue fibrogenesis (Wynn (2004), *supra*), the development and progression of hepatic fibrosis in *S. mansoni*-infected IL-21R^{-/-} mice was next examined. Liver hydroxyproline levels were assayed at various time points post-infection as a direct measure of tissue collagen content. As expected, marked hepatic fibrosis was observed in the infected WT mice (FIG. 22D). In contrast, the IL-21R^{-/-} displayed significantly less fibrosis at both the acute and chronic time points. Notably, by week 29 post-infection the IL-21R^{-/-} mice exhibited more than a 50% decrease in total liver collagen content compared to WT mice (FIG. 22E), thus confirming an important and indispensable role for the IL-21R in the progression of T_H2 -dependent fibrosis.

[0185] Experiments were undertaken to examine whether an IL-21 inhibitor could slow the progression of fibrosis in infected WT mice. For these experiments, groups of C57BL/6 mice were treated with sIL-21R-Fc or control protein for a total of 5 weeks, starting on week 6 post-infection, around the time when eggs are first detected in the liver. Although both groups had similar worm and tissue egg burdens (data not shown), mice receiving the IL-21 blocker displayed over a 50% reduction in hepatic fibrosis at the termination of the experiment (FIG. 22F). IL-4 and IL-13 mRNA expression also decreased in the liver, and granuloma size was reduced approximately 15% (data not shown). Thus, these data compliment the experiments performed with IL-21R^{-/-} mice.

Example 8: IL-21 Signaling Promotes the Development of Alternatively Activated Macrophages

[0186] Because Arg-1, FIZZ1, and TGF- β 1 have been linked with the development of fibrosis, and the expression of several T_H2 /Stat6-regulated genes were reduced in the diseased tissues of IL-21R^{-/-} mice (FIGS. 17-20) (see also, Gordon, *supra*; Nair et al., *supra*), experiments were undertaken to determine whether Arg-1, FIZZ1, and TGF- β 1 were directly modulated in macrophages following stimulation with IL-21. Arg-1 and FIZZ1 are also well-known markers of alternatively activated macrophages (AAM ϕ) (Gordon, *supra*). For these studies, bone marrow-derived macrophage cultures (BMM ϕ) were generated and then stimulated with various combinations of IL-4, IL-13, and IL-21. As

expected, IL-4 and IL-13 both increased Arg-1 and FIZZ1 mRNA expression, with an additive effect observed when the two stimuli were used in combination (FIG. 23A). Notably however, although IL-21 had no effect on either gene when used alone, cultures that were pretreated with IL-21 displayed highly significant increases in Arg-1 and FIZZ1 mRNA expression when subsequently stimulated with IL-4 and IL-13 (FIG. 23A). The same combination also significantly increased the function of arginase in the cells (FIG. 23B). In contrast, IL-21 had no effect on the levels of total or active TGF- β 1 in the culture supernatants (FIG. 24). Unexpectedly, IL-21 treatment alone significantly increased the expression IL-4R α and IL-13R α 1 (FIG. 23C). In contrast, IL-4 and IL-13 had no effect when used alone (FIG. 23C) and there was no additional effect when the three stimuli were used in combination (not shown).

[0187] Because the IL-13R α 2 can also influence IL-13-dependent signaling (Chiaromonte et al. (2003), *supra*; Mentink-Kane et al. *Proc. Natl. Acad. Sci. U.S.A.* 101:586-90; Wood et al. (2003) *J. Exp. Med.* 197:703-09), experiments were undertaken to examine whether IL-21 was regulating the production of the IL-13R α 2. Not surprisingly, because the IL-13R α 2 is primarily produced by nonhematopoietic cells like fibroblasts and smooth muscle (Chiaromonte et al. (2003), *supra*; Jakubzick et al. (2003) *Am. J. Pathol.* 162:1475-86; Zheng et al. (2003) *J. Allergy Clin. Immunol.* 111:720-28; Morimoto et al. (2006) *J. Immunol.* 176:342-48), there was no evidence of decoy receptor regulation in the BMM ϕ cultures (data not shown). However, when the regulation of the decoy receptor *in vivo* was examined, IL-21 downregulated IL-13R α 2 mRNA expression in the lungs of i.v. egg-challenged mice and significantly decreased the levels of the soluble IL-13R α 2 in their serum (FIG. 23D). When viewed together, these data suggest that IL-21 contributes to the development of alternatively activated macrophages by upregulating the type-2 IL-4 receptor (signaling receptor) in macrophages and by simultaneously decreasing the levels of the soluble IL-13R α 2 (decoy receptor) in the serum. Both mechanisms likely contributed to the increased activation of Arg-1 and FIZZ1 in the IL-4/IL-13-stimulated macrophages. As such, they provide an additional mechanistic explanation for the impaired T_H2 responses and T_H2-dependent fibrosis in the helminth-infected IL-21R^{-/-} mice.

Example 9: Discussion

[0188] IL-21 was recently characterized as a T_H2 cytokine that can inhibit the differentiation of naïve T_H cells into IFN- γ -producing T_H1 cells (Wurster et al. (2002), *supra*). Because the immune response in schistosomiasis evolves from an early IFN- γ to a sustained and dominant T_H2 response (Pearce and MacDonald, *supra*), the influence of IL-21R signaling on the development of helminth-induced T_H2 responses was examined. Infection of WT mice with *S. mansoni* increased IL-21 and IL-21R expression in the liver, confirming an association of IL-21 signaling with helminth-induced type-2 immunity. However, in the lung, schistosome eggs induced significant IL-21 expression during both T_H1 and T_H2 polarized responses. In fact, IL-21 expression increased most when mice were polarized to a T_H1 response. These data suggested that IL-21 exhibits a less restricted pattern of expression than that of the other T_H2-associated cytokines. The receptor for IL-21 also failed to display a T_H1/T_H2-specific pattern. However, the IL-21 receptor was induced nearly 4-fold more in the lungs of T_H2 versus T_H1 polarized mice, which provided one of the first indications that IL-21R signaling might be involved in the regulation of T_H2-mediated inflammation.

[0189] To determine whether type-2 effector responses were compromised in the absence of the IL-21R, the expression of several genes that are induced preferentially under T_H2-polarizing conditions was examined. These genes included AMCase, Ym1, and FIZZ1, all of which are thought to play important and nonredundant roles in the regulation of T_H2-mediated inflammation (Zhu et al., *supra*; Chiaramonte et al. (2003), *supra*; Nair et al., *supra*; Mentink-Kane et al., *supra*; Guo et al. (2000) *J. Biol. Chem.* 275:8032-37). Although some variation was observed during a primary, secondary, or chronic immune response, in each case the IL-21R^{-/-} mice displayed highly significant decreases in these T_H2-associated genes. Ym1 and AMCase are members of a family of proteins that share homology with chitinases of lower organisms (Nair et al., *supra*). Although their exact function in host immune reactions remains uncertain, they are thought to play important roles in eosinophil chemotaxis, tissue remodeling and fibrosis. Indeed, a recent study showed that AMCase neutralization could ameliorate allergen-driven inflammation and airway

hyperresponsiveness, thus confirming the participation of mammalian chitinases in T_H2 immunity (Zhu et al, *supra*). FIZZ1 is also associated with tissue fibrogenesis (Mentink-Kane et al., *supra*; Liu et al. (2004) *J. Immunol.* 173:3425-31). Consequently, a major function of the IL-21R may be to regulate the mechanisms of wound healing and fibrosis. Therefore, in addition to its participation in helminth-induced immune responses, the IL-21R may be involved in the regulation of a variety of T_H2-mediated inflammatory disorders.

[0190] In schistosomiasis, IL-21R-deficiency had a profound effect on the progression of the disease. Although infection intensities were the same in WT and IL-21R^{-/-} mice, the egg-induced inflammatory response decreased significantly in the absence of the IL-21R. There was also a marked reduction in secondary granuloma formation and a faster resolution of primary granulomas in the lung. Together, these data illustrate an indispensable role for the IL-21R in granulomatous inflammation. Previous studies showed that IL-4 and IL-13 are essential for lesion formation (Pearce and MacDonald; *supra*), thus the IL-21R is believed to be either directly or indirectly affecting the activity of these cytokines. These studies suggested that IL-21 was not acting alone since extremely high levels of IL-21 were observed in IL-4/IL-10 double knockout mice, yet granuloma formation was almost completely ablated in these T_H2-deficient animals (Hoffmann et al. (2000) *J. Immunol.* 164:6406-16; Sandler et al. (2003) *J. Immunol.* 171:3655-67). Thus, IL-21 appears to collaborate with IL-4 and IL-13 to induce a maximal response. The data disclosed herein show there is no detectable change in the cellular composition of the granulomas in IL-21R^{-/-} mice and no specific impairment in CD4⁺ T cell recruitment. Together, these findings suggested that the IL-21R regulates the development of parasite-induced pathology by modulating the overall intensity of the T_H2 effector response.

[0191] IL-21 is not thought to regulate IL-4-induced T_H2 cell differentiation directly (Suto et al., *supra*; Wurster et al. (2002), *supra*). Instead, it was hypothesized in a recent paper that IL-21 might amplify T_H2-driven responses by downregulating the expansion of IFN- γ -producing T_H1 cells (Wurster et al. (2002), *supra*). As such, it is theorized that the IFN- γ response in schistosome-infected mice might increase in the absence of the IL-21 receptor. Although a small increase was observed in lung-associated lymph nodes *in vitro*, IFN- γ

production was consistently reduced in the granulomatous tissues. Thus, the studies did not show that endogenous IL-21R played a substantial role in the inhibition of IFN- γ production during helminth infection. However, the IL-21R^{-/-} mice simultaneously generated weaker T_H1 and T_H2 cytokine responses in the tissues. The significant reduction in IgG_{2b} (T_H1-associated) and IgG₁ (T_H2-associated) antibody titers at all times post-infection supports this conclusion. Th2 cytokines were also decreased at the mRNA level in both the lungs and lymph nodes following *N. brasiliensis* infection. Indeed, all of the direct *ex vivo* data confirmed a marked reduction in T_H2 cytokine expression and function within the affected tissues. Nevertheless, there was no consistent reduction in T_H2 cytokine production by isolated lymphocytes following antigen restimulation, which suggests the IL-21R^{-/-} mice are capable of generating significant T_H2 responses, at least *in vitro*. Thus, the data disclosed herein suggest the IL-21R is selectively augmenting T_H2 responses in the tissues. In addition to promoting the T_H2 response, the IL-21R also increased IL-21 production. Thus, the IL-21R appears to operate in an autocrine fashion to drive T_H2 cytokine expression and type-2 effector functions *in vivo*.

[0192] To further elucidate the mechanisms involved, experiments were undertaken to determine whether IL-21 was directly modulating macrophage function, because the *in vivo* data showed a marked reduction in several genes that have been associated with the “alternatively-activated” phenotype (Gordon, *supra*; Mantovani et al. (2005) *Immunity* 23:344-46). Macrophages and fibroblasts exhibiting an alternatively activated phenotype are major cellular constituents of schistosome granulomas and functional studies suggested they are critically involved in the progression of the disease (Hesse et al. (2001), *supra*). Indeed, an important study by Brombacher et al. showed that mice that are completely deficient in alternatively activated macrophages develop lethal egg-induced pathology following infection with *S. mansoni* (Herbert et al. (2004) *Immunity* 20:623-35). In addition, because macrophage-derived TGF- β 1 has been implicated in the mechanism of IL-13-mediated fibrosis (Lee et al. (2001) *J. Exp. Med.* 194:809-21; Fichtner-Feigl et al. (2006) *Nat. Med.* 12:99-106), experiments were undertaken to determine whether IL-21 was modulating TGF- β 1 production in macrophages. To investigate these issues, Arg-1 and

FIZZ1 mRNA, arginase activity, and TGF- β 1 protein responses were measured in bone marrow-derived macrophages following stimulation with various combinations of IL-21, IL-4, and IL-13. Arg-1 and FIZZ1 are IL-4R α /Stat6-dependent genes (Liu et al., *supra*; Hesse et al. (2001), *supra*; Munder et al. (1998) *J. Immunol.* 160:5347-54); therefore, they serve as functional markers of alternative macrophage activation. Importantly, the findings suggested that when macrophages were exposed to IL-21, they became much more sensitive to the Arg-1- and FIZZ1-inducing activities of IL-4 and IL-13. Arginase activity assessed by the production of urea also increased significantly, confirming IL-21 as an important stimulus for the development of highly functional alternatively activated macrophages. In contrast, IL-21 had no effect on the production of TGF- β 1 by macrophages. Thus, the pro-fibrotic cytokine TGF- β 1 does appear to be involved, which is consistent with previous studies that have investigated the role of TGF- β 1 in schistosomiasis (Kaviratne et al. (2004) *J. Immunol.* 173:4020-29). Instead, IL-21 significantly increased IL-4R α and IL-13R α 1 expression in BMM ϕ s and decreased the production of the soluble IL-13 decoy receptor *in vivo*, which likely explains their heightened sensitivity to IL-4 and IL-13. As such, these data compliment the *in vivo* studies with IL-21R $^{-/-}$ mice and suggest that an important function of IL-21R signaling is to enhance the development of AAM ϕ , which have been implicated in the mechanism of fibrosis (Hesse et al. (2001), *supra*; Hesse et al. (2000), *supra*). Moreover, because AAM ϕ have been shown to amplify CD4 $^{+}$ T $_H$ 2 cell differentiation (Bonecchi et al. (1998) *Blood* 92:2668-71), these data may also explain the overall reduction in helminth-induced T $_H$ 2 activity in the IL-21R $^{-/-}$ mice.

[0193] In human schistosomiasis, the development of fibrotic liver pathology is the principle cause of chronic morbidity and mortality (Pearce and MacDonald, *supra*; Wynn et al. (2004) *Immunol. Rev.* 201:156-67). Because the T $_H$ 2 cytokine response is known to play an important role in collagen deposition (Wynn et al. (2004), *supra*), a final series of experiments examined the influence of the IL-21R on the progression of hepatic fibrosis. Notably, development of fibrosis decreased significantly in the IL-21R $^{-/-}$ mice, with the knockout animals displaying over a 50% reduction in hepatic fibrosis by week 29 post-infection. Importantly, similar findings were also generated when infected WT mice were

treated with sIL-21R-Fc. Thus, the IL-21 receptor was revealed as a potential new target for anti-fibrotic therapy. In conclusion, these studies illustrate an essential role for the IL-21R in the progression of T_H2 cytokine-mediated disease. As such, the IL-21R should be added to list of important receptors that regulate type-2 immunity and macrophage polarization.

Example 10: Prophetic Treatments

[0194] A nonlimiting set of prophetic treatment examples follows.

[0195] A subject diagnosed with liver cirrhosis is administered an IL-21R fusion protein to reduce the accumulation of fibrotic tissue in the liver. The IL-21R fusion protein includes amino acids 1-235 of SEQ ID NO:2 fused at its C-terminus via a linker (corresponding to amino acids 236-243 of SEQ ID NO:17) to a human immunoglobulin G1 (IgG1) Fc-mutated sequence (corresponding to amino acids 244-467 of SEQ ID NO:17).

[0196] A subject diagnosed with an infection with schistosoma is administered a soluble IL-21R fragment to reduce the accumulation of fibrotic tissue. The fragment contains amino acids 20-538 of SEQ ID NO:2.

[0197] Following surgery, a subject is administered an IL-21R antibody to reduce the accumulation of fibrosis due to surgical incision during the wound healing process.

[0198] A subject diagnosed with liver cirrhosis is administered an IL-21 antibody to reduce the accumulation of fibrotic tissue in the liver.

WHAT IS CLAIMED IS:

1. A method for treating, ameliorating, or preventing fibrosis or a fibrosis-associated disorder in a subject comprising administering to the subject a therapeutically effective amount of an agent that reduces the level of IL-21 and/or IL-21R in the subject.
2. The method of claim 1, wherein the agent is an IL-21/IL-21R antagonist selected from the group consisting of an anti-IL-21R antibody, an anti-IL-21 antibody, an antigen-binding fragment of an anti-IL-21R antibody, an antigen-binding fragment of an anti-IL-21 antibody, and a soluble fragment of an IL-21R.
3. The method of claim 2, wherein the agent is a soluble fragment of an IL-21R, and wherein the soluble fragment of the IL-21R comprises an amino acid sequence that is at least 90% identical to an amino acid sequence selected from the group consisting of amino acids 1-538 of SEQ ID NO:2, amino acids 20-538 of SEQ ID NO:2, amino acids 1-235 of SEQ ID NO:2, amino acids 20-235 of SEQ ID NO:2, amino acids 1-236 of SEQ ID NO:2, amino acids 20-236 of SEQ ID NO:2, amino acids 1-529 of SEQ ID NO:5, amino acids 20-529 of SEQ ID NO:5, amino acids 1-236 of SEQ ID NO:5, and amino acid 20-236 of SEQ ID NO:5.
4. The method of claim 3, wherein the soluble fragment of the IL-21R binds to an IL-21 polypeptide.
5. The method of claim 1, wherein the agent is a soluble fragment of an IL-21R, and wherein the soluble fragment of the IL-21R comprises an amino acid sequence that is substantially identical to the amino acid sequence set forth in SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, or SEQ ID NO:27.
6. The method of claim 5, wherein the amino acid sequence of the soluble fragment of the IL-21R comprises an amino acid sequence that is substantially identical to the amino acid sequence set forth in SEQ ID NO:11 or SEQ ID NO:13.

7. The method of claim 1, wherein the agent is a soluble fragment of an IL-21R, and wherein the soluble fragment of the IL-21R is encoded by a nucleotide sequence that is substantially identical to the nucleic acid sequence set forth in SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, or SEQ ID NO:26.
8. The method of claim 7, wherein the soluble fragment of the IL-21R is encoded by a nucleotide sequence that is substantially identical to the nucleic acid sequence set forth in SEQ ID NO:12 or SEQ ID NO:16.
9. The method of claim 2, wherein the agent is a soluble fragment of an IL-21R, and wherein the soluble fragment of the IL-21R comprises an extracellular domain of IL-21R and an immunoglobulin Fc fragment.
10. The method of claim 9, wherein the amino acid sequence of the extracellular domain of the IL-21R comprises an amino acid sequence that is at least 90% identical to amino acids 1-235 of SEQ ID NO:2 or amino acids 20-235 of SEQ ID NO:2.
11. The method of claim 9, wherein the immunoglobulin Fc fragment has an altered function.
12. The method of claim 11, wherein the immunoglobulin Fc fragment has the amino acid sequence of amino acids 244-467 of SEQ ID NO:17.
13. The method of claim 1, wherein the fibrosis or fibrosis-associated disorder affects the liver, epidermis, endodermis, muscle, tendon, cartilage, heart, pancreas, lung, uterus, nervous system, testis, ovary, adrenal gland, artery, vein, colon, small intestine, biliary tract, or stomach.
14. The method of claim 13, wherein the fibrosis or fibrosis-associated disorder affects the liver, epidermis, endodermis, or lung.
15. The method of claim 14, wherein the fibrosis or fibrosis-associated disorder is interstitial lung fibrosis.

16. The method of claim 13, wherein the fibrosis or fibrosis-associated disorder is the result of an infection with schistosoma.

17. The method of claim 1, wherein the fibrosis or fibrosis-associated disorder is the result of wound healing.

18. The method of claim 17, wherein the wound healing results from a surgical incision.

19. The method of claim 1, further comprising administering to the subject at least one additional therapeutic agent.

20. The method of claim 19, wherein the at least one additional therapeutic agent is selected from the group consisting of cytokine inhibitors, growth factor inhibitors, immunosuppressants, anti-inflammatory agents, metabolic inhibitors, enzyme inhibitors, cytotoxic agents, and cytostatic agents.

21. The method of claim 19, wherein the at least one additional therapeutic agent is selected from the group consisting of TNF antagonists, anti-TNF agents, IL-12 antagonists, IL-15 antagonists, IL-17 antagonists, IL-18 antagonists, IL-22 antagonists, T cell-depleting agents, B cell-depleting agents, cyclosporin, FK506, CCI-779, etanercept, infliximab, rituximab, adalimumab, prednisolone, azathioprine, gold, sulphasalazine, hydroxychloroquine, minocycline, anakinra, abatacept, methotrexate, leflunomide, rapamycin, rapamycin analogs, Cox-2 inhibitors, cPLA2 inhibitors, NSAIDs, p38 inhibitors, antagonists of B7.1, B7.2, ICOSL, ICOS and/or CD28, and agonists of CTLA4.

22. The method of claim 1, wherein the subject is a human.

23. A method for identifying a compound for treating, ameliorating or preventing fibrosis or a fibrosis-associated disorder in a subject, comprising: (a) measuring the level of IL-21 and/or IL-21R in a cell or sample of interest; (b) contacting the cell or sample of interest with a compound; and (c) measuring the level of IL-21 and/or IL-21R in the cell or sample of interest following contact with the compound,

wherein a lower level of IL-21 and/or IL-21R in the contacted cell or sample of interest, in comparison to the level of IL-21 and/or IL-21R in a noncontacted cell or sample of interest, identifies the compound as a compound useful for treating, ameliorating, or preventing fibrosis or a fibrosis-associated condition in a subject.

24. A method for identifying a compound for treating, ameliorating or preventing fibrosis or a fibrosis-associated disorder in a subject, comprising: (a) measuring the level of IL-21 and/or IL-21R in a cell or sample of interest; (b) contacting the cell or sample of interest with a compound; (c) measuring the level of IL-21 and/or IL-21R in the cell or sample of interest following contact with the compound; and (d) comparing the level of IL-21 and/or IL-21R in the contacted cell or sample of interest with a reference level of IL-21 and/or IL-21R, wherein a lower level of IL-21 and/or IL-21R in the contacted cell or sample of interest, in comparison to the reference level of IL-21 and/or IL-21R, identifies the compound as a compound useful for treating, ameliorating, or preventing fibrosis or a fibrosis-associated condition in a subject.

25. A method for monitoring the progress of fibrosis or a fibrosis-associated condition in a subject, comprising: (a) measuring the level of IL-21 and/or IL-21R in a cell or sample of interest from the subject at a first time point; and (b) measuring the level of IL-21 and/or IL-21R in a cell or sample of interest from the subject at a second time point, wherein a lower level of IL-21 and/or IL-21R in the cell or sample of interest from the subject at the second time point, in comparison to the level of IL-21 and/or IL-21R in the cell or sample of interest from the subject at the first time point, provides an indication that the fibrosis or fibrosis-associated condition has decreased in severity.

26. A method for prognosing fibrosis or a fibrosis-associated condition in a subject, comprising: (a) measuring the level of IL-21 and/or IL-21R in a cell or sample of interest from the subject at a first time point; and (b) measuring the level of IL-21 and/or IL-21R in a cell or sample of interest from the subject at a second time point, wherein a lower level of IL-21 and/or IL-21R in the cell or sample of interest from the subject at the second time point, in comparison to the level of IL-21 and/or IL-21R in the cell or sample of interest from the subject at the first

time point, indicates a decreased likelihood that the subject will develop fibrosis or the fibrosis-associated condition or a decreased likelihood that the fibrosis or fibrosis-associated condition will worsen in the subject.

27. A method for prognosing fibrosis or a fibrosis-associated condition in a subject, comprising: (a) measuring the level of IL-21 and/or IL-21R in a cell or sample of interest from the subject; and (b) comparing the level of IL-21 and/or IL-21R in the cell or sample of interest from the subject to a reference level of IL-21 and/or IL-21R, wherein a lower level of IL-21 and/or IL-21R in the cell or sample of interest from the subject, in comparison to the reference level of IL-21 and/or IL-21R, indicates a decreased likelihood that the subject will develop fibrosis or the fibrosis-associated condition or a decreased likelihood that the fibrosis or fibrosis-associated condition will worsen in the subject.

28. A method for diagnosing fibrosis or a fibrosis-associated condition in a subject, comprising: (a) measuring the level of IL-21 and/or IL-21R in a cell or sample of interest from the subject, and (b) comparing the level of IL-21 and/or IL-21R in the cell or sample of interest from the subject with a reference level of IL-21 and/or IL-21R, wherein a higher level of IL-21 and/or IL-21R in the cell or sample of interest from the subject, in comparison to the reference level of IL-21 and/or IL-21R, indicates the presence of fibrosis or the fibrosis-associated condition in the subject.

Fig. 1

1 GTCGACGCGG CGGTACCAGC TGTCTGCCCC CTTCTCCTGT GGTGTGCCTC
51 ACGGTCACTT GCTTGTCTGA CCGCAAAGTCT GCCCATCCCT GGGGCAGCCA
101 ACTGGCCTCA GCCCGTGCCC CAGGCGTGCC CTGTCTCTGT CTGGCTGCCC
151 CAGCCCTACT GTCTTCCTCT GTGTAGGCTC TGCCCAGATG CCCGGCTGGT
201 CCTCAGCCTC AGGACTATCT CAGCAGTGAC TCCCCTGATT CTGGACTTGC
251 ACCTGACTGA ACTCCTGCCC ACCTCAAACC TTCACCTCCC ACCACCACCA
301 CTCCGAGTCC CGCTGTGACT CCCACGCCCC GGAGACCACC CAAGTGCCCC
351 AGCCTAAAGA ATGGCTTTCT GAGAAAGACC CTGAAGGAGT AGGTCTGGGA
401 CACAGCATGC CCCGGGGCCC AGTGGCTGCC TTACTCCTGC TGATTCTCCA
451 TGGAGCTTGG AGCTGCCTGG ACCTCACTTG CTACACTGAC TACCTCTGGA
501 CCATCACCTG TGTCTGGAG ACACGGAGCC CCAACCCAG CATACTCAGT
551 CTCACCTGGC AAGATGAATA TGAGGAACTT CAGGACCAAG AGACCTTCTG
601 CAGCCTACAC AGGTCTGGCC ACAACACCAC ACATATATGG TACACGTGCC
651 ATATGCGCTT GTCTCAATTC CTGTCCGATG AAGTTTTCAT TGTCAATGTG
701 ACGGACCAGT CTGGCAACAA CTCCCAAGAG TGTGGCAGCT TTGTCCTGGC
751 TGAGAGCATC AAACCAGCTC CCCCCTTGAA CGTGA CTGTG GCCTTCTCAG
801 GACGCTATGA TATCTCCTGG GACTCAGCTT ATGACGAACC CTCCA ACTAC
851 GTGCTGAGGG GCAAGCTACA ATATGAGCTG CAGTATCGGA ACCTCAGAGA
901 CCCCTATGCT GTGAGGCCGG TGACCAAGCT GATCTCAGTG GACTCAAGAA
951 ACGTCTCTCT TCTCCCTGAA GAGTTCCACA AAGATTCTAG CTACCAGCTG
1001 CAGGTGCGGG CAGCGCCTCA GCCAGGCACT TCATT CAGGG GGACCTGGAG
1051 TGAGTGGAGT GACCCCGTCA TCTTTCAGAC CCAGGCTGGG GAGCCCGAGG
1101 CAGGCTGGGA CCCTCACATG CTGCTGCTCC TGGCTGTCTT GATCATTTGTC
1151 CTGGTTTTCA TGGGTCTGAA GATCCACCTG CCTTGGAGGC TATGGAAAAA
1201 GATATGGGCA CCAGTGCCCC CCCCTGAGAG TTTCTTCCAG CCCCTGTACA
1251 GGGAGCACAG CGGGAAC TTC AAGAAATGGG TTAATACCCC TTTCACGGCC
1301 TCCAGCATAG AGTTGGTGCC ACAGAGTTCC ACAACAACAT CAGCCTTACA
1351 TCTGTCA TTG TATCCAGCCA AGGAGAAGAA GTTCCCGGGG CTGCCGGGTC
1401 TGGAAGAGCA ACTGGAGTGT GATGGAATGT CTGAGCCTGG TCACTGGTGC

Fig. 1 (continued)

1451 ATAATCCCCCT TGGCAGCTGG CCAAGCGGTC TCAGCCTACA GTGAGGAGAG
1501 AGACCGGCCA TATGGTCTGG TGTCCATTGA CACAGTGA CT GTGGGAGATG
1551 CAGAGGGCCT GTGTGTCTGG CCCTGTAGCT GTGAGGATGA TGGCTATCCA
1601 GCCATGAACC TGGATGCTGG CCGAGAGTCT GGCCCTAATT CAGAGGATCT
1651 GCTCTTGGTC ACAGACCCTG CTTTCTGTG TTGCGGCTGT GTCTCAGGTA
1701 GTGGTCTCAG GCTTGGAGGC TCCCCAGGCA GCCTACTGGA CAGGTTGAGG
1751 CTGTCA TTG CAAAGGAAGG GGA CTGGACA GCAGACCCAA CCTGGAGAAC
1801 TGGGTCCCCA GGAGGGGGCT CTGAGAGTGA AGCAGGTTCC CCCCCTGGTC
1851 TGGACATGGA CACATTTGAC AGTGGCTTTG CAGGTT CAGA CTGTGGCAGC
1901 CCCGTGGAGA CTGATGAAGG ACCCCCTCGA AGCTATCTCC GCCAGTGGGT
1951 GGT CAGGACC CCTCCACCTG TGGACAGTGG AGCCCAGAGC AGCTAGCATA
2001 TAATAACCAG CTATAGTGAG AAGAGGCCTC TGAGCCTGGC ATTTACAGTG
2051 TGAACATGTA GGGGTGTGTG TGTGTGTGTG TGTGTGTGTG TGTGTGTGTG
2101 TGTGTGTGTG TGTGTGTGTG TGTCTTGGGT TGTGTGTTAG CACATCCATG
2151 TTGGGATTTG GTCTGTTGCT ATGTATTGTA ATGCTAAATT CTCTACCCAA
2201 AGTTC TAGGC CTACGAGTGA ATTCTCATGT TTACAAACTT GCTGTGTAAA
2251 CCTTGTT CCT TAATTTAATA CCATTGGTTA AATAAAATTG GCTGCAACCA
2301 ATTACTGGAG GGATTAGAGG TAGGGGGCTT TTGAGTTACC TGT TTGGAGA
2351 TGGAGAAGGA GAGAGGAGAG ACCAAGAGGA GAAGGAGGAA GGAGAGGAGA
2401 GGAGAGGAGA GGAGAGGAGA GGAGAGGAGA GGAGAGGAGA GGAGAGGAGA
2451 GGCTGCCGTG AGGGGAGAGG GACCATGAGC CTGTGGCCAG GAGAAACAGC
2501 AAGTATCTGG GGTACACTGG TGAGGAGGTG GCCAGGCCAG CAGTTAGAAG
2551 AGTAGATTAG GGGTGACCTC CAGTATTTGT CAAAGCCAAT TAAAATAACA
2601 AAAAAAAAAA AAAAGCGGCC GCTCTAGA

Fig. 2A

1 MPRGPVAALL LLILHGAWSC LD LTCYTDYL WTITCVLETR SPNPSILSLT
51 WQDEYEELQD QETFCSLHRS GHNTTHIWYT CHMRLSQFLS DEVFIVNVD
101 QSGNNSQECG SFVLAESIKP APPLNVTVAF SGRYDISWDS AYDEPSNYVL
151 RGKLQYELQY RNL RDPYAVR PVTKLISVDS RNVSL LPEEF HKDSSYQLQV
201 RAAPQPGTSF RGT**WSEWS**DP VIFQTQAGEP EAGWDPHMLL LLAVLIIVLV
251 FMGLKIHLPW RLWKKIWAPV PTPESFFQPL YREHSGNFKK WVNTPF TASS
301 IELVPQSSTT TSALHLSLYP AKEKKFPGLP GLEEQL ECDG MSEP GHWCII
351 PLAAGQAVSA YSEERDRPYG LVSIDTVTVG DAEGLCVWPC SCEDDGYPAM
401 NLDAGRESGP NSEDLLL VTD PAFLSCGCVS GSGLRLGGSP GSLLDRLRLS
451 FAKEGDWTAD PTWRTGSPGG GSESEAGSPP GLDMDTFDSG FAGSDCGSPV
501 ETDEGP PRSY LRQWVVRTPP PVDSGAQSS




Figure 2B

Fig. 3

```

huMU-1      .....NNGTCGACTGGAGGCCCAGCTGCCCGTCATCA 32
murMU-1      CAGCCCTACTGTCTTCCTCTGTGTAGGCTCTGCCCAGATGCCCGGC.... 196
huMU-1      GAGTGACAGGTCTTATGACAGCCTGATTGGTGA CTGGGCTGGGTGTGGA 82
murMU-1      TGGTCCTCAGCCTCAGGACTATCTCAGCAGTGA CTG.CCCTGATTCTGGA 245
huMU-1      TTCTCACCCCAGGCCTCTGCCTGCTTTCTCAGACCCTCATCT...GTCAC 129
murMU-1      CTTGCACCTGACTGAACTCCTGCCACCTCAAACCTTACCTCCCACCAC 295
huMU-1      CCCACAGCTGAACCCAGCTG.....CCACCCCAGAAGCCCATCAGACT 173
murMU-1      CACCACTCCGAGTCCCGCTGTGACTCCACGCCCAGGAGACCACCCAAGT 345
huMU-1      GCCCCCAGCACAACGGAATGGATTTCTGAGAAAGAAGCCGAAACAGAAGGC 223
murMU-1      G.CCCCAGCCTAAAGAATGGCTTTCTGAGAAAGACCCTGAAGGAGTAGGT 394
huMU-1      CCGTGGGAGTCAGCATGCCGCGTGGCTGGGCCGCCCCCTTGCTCCTGCTG 273
murMU-1      C..TGGGACACAGCATGCCCGGGGCCAGTGGCTGCCTTACTCCTGCTG 442
huMU-1      CTGCTCCAGGGAGGCTGGGGCTGCCCGACCTCGTCTGCTACACCGATTA 323
murMU-1      ATTCTCCATGGAGCTTGGAGCTGCCCTGGACCTCACTTGCTACACTGACTA 492
huMU-1      CCTCCAGACGGTCATCTGCATCCTGGAAATGTGGAACCTCCACCCAGCA 373
murMU-1      CCTCTGGACCATCACCTGTGTCTCTGGAGACACGGAGCCCCAACCCAGCA 542
huMU-1      CGCTCACCCCTTACCTGGCAAGACCAGTATGAAGAGCTGAAGGACGAGGCC 423
murMU-1      TACTCAGTCTCACCTGGCAAGATGAATATGAGGAACTTCAGGACCAAGAG 592
huMU-1      ACCTCCTGCAGCCTCCACAGGTGGGCCCAATGCCACGCATGCCACCTA 473
murMU-1      ACCTTCTGCAGCCTACACAGGTCTGGCCACAACACCACACATATATGGTA 642
huMU-1      CACCTGCCACATGGATGTATTCCACTTCATGGCCGACGACATTTTCAGTG 523
murMU-1      CACGTGCCATATGCGCTTGTCTCAATTCTGTCCGATGAAGTTTTCATTG 692
huMU-1      TCAACATCACAGACCAGTCTGGCAACTACTCCAGGAGTGTGGCAGCTTT 573
murMU-1      TCAATGTGACGGACCAGTCTGGCAACAAC TCCAAGAGTGTGGCAGCTTT 742
huMU-1      CTCCTGGCTGAGAGCATCAAGCCGGCTCCCCCTTTCAACGTGACTGTGAC 623
murMU-1      GTCCTGGCTGAGAGCATCAAACCAGCTCCCCCTTGAACGTGACTGTGGC 792
huMU-1      CTTCTCAGGACAGTATAATATCTCCTGGCGCTCAGATTACGAAGACCCTG 673
murMU-1      CTTCTCAGGACGCTATGATATCTCCTGGGACTCAGCTTATGACGAACCT 842
huMU-1      CCTTCTACATGCTGAAGGGCAAGCTTCA GTATGAGCTGCAGTACAGGAAC 723
murMU-1      CCAACTACGTGCTGAGGGGCAAGCTACA ATATGAGCTGCAGTATCGGAAC 892

```

Fig. 3 (continued)

huMU-1	CGGGGAGACCCCTGGGCTGTGAGTCCGAGGAGAAAGCTGATCTCAGTGGA	773
murMU-1	CTCAGAGACCCCTATGCTGTGAGGCCGGTGACCAAGCTGATCTCAGTGGA	942
huMU-1	CTCAAGAAAGTGTCTCCCTCCTCCCCCTGGAGTTCCGCAAAGACTCGAGCT	823
murMU-1	CTCAAGAAACGTCTCTCTTCTCCCTGAAGAGTTCCACAAAGATTCTAGCT	992
huMU-1	ATGAGCTGCAGGTGCGGGCAGGGCCCATGCCTGGCTCCTCCTACCAGGGG	873
murMU-1	ACCAGCTGCAGGTGCGGGCAGCGCCTCAGCCAGGCAC TTCATT CAGGGGG	1042
huMU-1	ACCTGGAGTGAATGGAGTGACCCGGTCATCTTTTCAGACCCAGTCAGAGGA	923
murMU-1	ACCTGGAGTGAATGGAGTGACCCCGTCATCTTTTCAGACCCAGGCTGGGGA	1092
huMU-1	GTTAAAGGAAGGCTGGAACCCCTCACCTGCTGCTTCTCCTCCTGCTTGTC	973
murMU-1	GCCCCAGGCAGGCTGGGACCCCTCATATGCTG...CTGCTCCTGGCTGTCT	1139
huMU-1	TAGTCTTCATTCCTGCCTTCTGGAGCCTGAAGACCCATCCATTGTGGAGG	1023
murMU-1	TGATCATTTGCTCCTGGTTTTCATGGGTCTGAAGATCCACCTGCCTTGAGG	1189
huMU-1	CTATGGAAGAAGATATGGG...CCGTCCCAGCCCTGAGCGGTTCTTCAT	1070
murMU-1	CTATGGAAGAAAGATATGGGACACAGTGCCACCCCTGAGAGTTTCTTCCA	1239
huMU-1	GCCCCCTGTACAAGGGCTGCAGCGGAGACTTCAAGAAATGGGTGGGTGCAC	1120
murMU-1	GCCCCCTGTACAGGGAGCACAGCGGGAAC TCAAGAAATGGGTAAATACCC	1289
huMU-1	CCTTCACTGGCTCCAGCCTGGAGCTGGGACCCCTGGAGCC CAGAGGTGCC	1170
murMU-1	CTTTTACGGCCTCCAGCATAGAGTTGGTGCCACAGAGTTCCACAACAACA	1339
huMU-1	TCCACCCTGGAGGTGTACAGCTGCCACCCACCACGGAGCCCGGCCAAGAG	1220
murMU-1	TCAGCCTTACATCTGT.....CATTGTATCCAGCCAAGGA	1374
huMU-1	GCTGCAGCTCACGGAGCTACAAGAACCAGCAGAGCTGGTGGAGTCTGACG	1270
murMU-1	GAAGAAGTTCCCGGGGCTGCCGGGTCTGGAAGAGCAACTGGAGTGTGATG	1424
huMU-1	GTGTGCCCAAGCCAGCTTCTGG.....CCGACAGCCAGAACTCG	1311
murMU-1	GAATGTCTGAGCCTGGTCACTGGTGCATAATCCCCTTGGCAGCTGGCCAA	1474
huMU-1	GGGGGCTCAGCTTACAGTGAGGAGAGGGATCGGCCATACGGCCTGGTGTC	1361
murMU-1	GCGGTCTCAGCCTACAGTGAGGAGAGAGACCGGCCATATGGTCTGGTGTC	1524
huMU-1	CATTGACACAGTGACTGTGCTAGATGCAGAGGGGCCATGCACCTGGCCCT	1411
murMU-1	CATTGACACAGTGACTGTGGGAGATGCAGAGGGCCTGTGTGTCTGGCCCT	1574
huMU-1	GCAGCTGTGAGGATGACGGCTACCCAGCCCTGGACCTGGATGCTGGCCTG	1461
murMU-1	GTAGCTGTGAGGATGATGGCTATCCAGCCATGAACCTGGATGCTGGCCGA	1624
huMU-1	GAGCCCAGCC CAGGCCTAGAGGACCCACTCTTGATGCAGGGACCACAGT	1511
murMU-1	GAGTCTGGCCCTAATT CAGAGGATCTGCTCTTGGT CACAGACCCTGCTTT	1674

Fig. 3 (continued)

huMU-1	CCTGTCCTGTGGCTGTGTCTCAGCTGGCAGCCCTGGGCTAGGAGGGCCCC	1561
murMU-1	TCTGTCTTGCGGCTGTGTCTCAGGTAGTGGTCTCAGGCTTGGAGGCTCCC	1724
huMU-1	TGGGAAGCCTCCTGGACAGACTAAAGCCACCCCTTGCAATGGGGAGGAC	1611
murMU-1	CAGGCAGCCTACTGGACAGGTTGAGGCTGTCATTTGCAAAGGAAGGGGAC	1774
huMU-1	TGGGCTGGGGGACTGCCCTGGGGTGGCCGGTCACCTGGAGGGGTCTCAGA	1661
murMU-1	TGGACAGCAGACCCAACCTGGAGAACTGGGTCCCCAGGAGGGGGCTCTGA	1824
huMU-1	GAGTGAGGCGGGCTCACCCCTGGCCGGCCTGGATATGGACACGTTTGACA	1711
murMU-1	GAGTGAAGCAGGTTCCCCC...CTGGTCTGGACATGGACACATTTGACA	1871
huMU-1	GTGGCTTTGTGGGCTCTGACTGCAGCAGCCCTGTGGAGTGTGACTTCACC	1761
murMU-1	GTGGCTTTGCAGGTTGAGACTGTGGCAGCCCCGTGGAGACT.....	1912
huMU-1	AGCCCCGGGGACGAAGGACCCCCCGGAGCTACCTCCGCCAGTGGGTGGT	1811
murMU-1GATGAAGGACCCCCCTCGAAGCTATCTCCGCCAGTGGGTGGT	1953
huMU-1	CATTCTCCGCCACTTTGAGCCCTGGACCCAGGCCAGCTAATGAGGCT	1861
murMU-1	CAGGACCCCTCCACCTGTGGACAGTGGAGCCAGAGCAGCTA.....	1995
huMU-1	GACTGGATGTCCAGAGCTGGCCAGGCCACTGGGCCCTGAGCCAGAGACAA	1911
murMU-1	.GCATATAATAACCAGCTATAGTGAGAAGAGGCCTCTGAGCC.....	2036
huMU-1	TGGGCCCTTTGAGCCTGATGTTTACAGTGTCTGTGTGTGTGTGTGCATATG	2011
murMU-1	TGGCATTTACAGTGTGAACATGTAGGGGTGTGTGTGTGTGTGTGTGTGTG	2086
huMU-1	TGTGTGTGTGCATATGCATGTGTGTGTGTGTGTGTGTGTGTCTTACTGGACTCA	2061
murMU-1	TG	2135
huMU-1	CGGAGCTCACCCATGTGCACAAGTGTGCACAGTAAACGTGTTTGTGGTCA	2111
murMU-1	GTTAGCACATCCATGTTGGGATTG.....GTCTGTTGCTA	2171
huMU-1	ACAGATGACAACAGCCGTCCTCCCTCCTAGGGTCTTGTGTTGCAAGTTGG	2161
murMU-1	TGTATTGTAATGCTAAATTCTCTACCCAAAGTTCTAGGCCTACGAGTGAA	2221
huMU-1	TCCACAGCATCTCCGGGGCTTTGTGGGATCAGGGCATTGCCTGTGACTGA	2211
murMU-1	TTCTCATGTTTACAAACTTGCTGTGTAAACCTTG...TTCCTTAATTAA	2268
huMU-1	GGCGGAGCCCAGCCCTCCAGCGTCTGCCTCCAGGAGCTGCAAGAAGTCCA	2261
murMU-1	TACCATTGGTTAAATAAAATTGGCTGCAACCAATTACTGGAGGGATTAGA	2318
huMU-1	TATTG.....TTCCTTATCACCTGCCAACAGGAAGCGAAAGGGGATGGAG	2306
murMU-1	GGTAGGGGGCTTTTGTAGTTACCTGTTTGGAGATGGAGAAGGAGAGAGGAG	2368

Fig. 3 (continued)

```

      .       .       .       .
huMU-1  TGAGCCCATGGTGTGACCTCGGGAATGGCAATTTTTTGGGCGGCCCTGGAC 2356
      || |      || ||      || |      |      || |      |
murMU-1  AGACCAAGAGGAGAAGGAGGAAGGAGAGGAGAGGAGAGGAGAGGAGAGGA 2418

      .       .       .       .
huMU-1  GAAGGTCTGAATCCCGACTCTGATACCTTCTGGCTGTGCTACCTGAGCCA 2406
      || |      ||      ||      ||      |      |||      ||
murMU-1  GAGGAGAGGAGAGGAGA.GGAGAGGAGAGGAGAGGCTGCCGTGAGGGGAG 2467

      .       .       .       .
huMU-1  AGTCGCCTCCCCCTCTCTGGGCTAGAGTTTCCTTATCCAGACAGTGGGGAA 2456
      ||      ||      || ||| | |||      |      |      |
murMU-1  AGGGACCATGAGCCTGTGGCCAGGAGAAACAGCA.....AGTA 2505

      .       .       .       .
huMU-1  GGCATGACACACCTGGGGGAAATTGGCGATGTCACCCGTGTACGGTACGC 2506
      |      |||| | | | | |||| | | ||| | || |      |
murMU-1  TCTGGGGTACACTGGTGTGAGGAGGTGGCCAGGCCAGC..AGTTAGAAGAGT 2553

      .       .       .       .
huMU-1  AGCCCAGAGCAGACCCTCAATAAACGTCAGCTTCCTTCAAAAAAAAAAAAA 2556
      ||      || | |||| | | || |||| | | |||| | | ||||
murMU-1  AGATTAGGGGTGACCTCCAGTATTTGTCAAAGCCAATTAAAATAACAAAA 2603

```

Human MU-1	MPRGWAAPLLLLLLLQGGWGCPDLVCYTDYLTQTVICILEMWNLHPSTLTLT	50
MurineMU-1	MPRGPVAAALLLLILHGAWSCLDLTCYTDYLTWTITCVLETRSPNPSILSLT	50
Human MU-1	WQDQYEELKDEATSCSLHRSAHNATHATYTCHEMDVVFHFMAADDIFSVNITD	100
MurineMU-1	WQDEYEELQDQETFCSLHRSGHNTTHIWTCTHMRLSQFLSDEVEFIVNVT	100
Human MU-1	QSGNYSQECGSFLLAESIKPAPPFNVTFTVTFSGQYNISWRSDYEDPAFYML	150
MurineMU-1	QSGNNSQECGSFVLAESIKPAPPLNVTFAFSGRYDISWDSAYDEPSNYVL	150
Human MU-1	KGKLYELQYRNREGDPWAVSPRRKLISVDSRSVSLPPEFRKDDSSYELQV	200
MurineMU-1	RGKLYELQYRNLRDPYAVRPVTKLISVDSRNVSLPPEEFHKDDSSYQLQV	200
Human MU-1	RAGPMPGSSYQGTWSEWSDPVIFQTQSEELKEGWNPHLLLLLLLVIVFIP	250
MurineMU-1	RAAPQPGTSTFRGTWSEWSDPVIFQTQAGEPEAGWDPHMLLLLAVALIIVL	249
Human MU-1	AFWSLKTHPLWRLWKKIWA.VPSPERFFMPLYKGCSDGFKKWVGAPFTGS	299
MurineMU-1	VFMGLKIHLPWRLWKKIWAPVPTPESFFQPLYREHSGNFKKWVNTPTFTAS	299
Human MU-1	SLELGPWSPEVPSTLEVYSCHPPRSPAKRLQLTELQEPALVESDGVPKP	349
MurineMU-1	SIELVPQSSSTTTTAL.....HLSLYPAKEKKFPGLPGLLEEQLCEDGMSEP	344
Human MU-1	SFW...PTAQNSGGSAYSEERDRPYGLVSDITVTVLDAEGPCTWPCSCD	396
MurineMU-1	GHWCIIPLAAGQAVSAYSEERDRPYGLVSDITVTVGDAEGLCVWPCSCD	394
Human MU-1	DGYPALDLDAGLEPSPGLEDPLLDAGTTFVLSCGCVSAGSPGLGGPLGSL	446
MurineMU-1	DGYPAMNLDAGRESGPNSDLLLLVTDPAFLSCGCVSGSGLRLGGSPGSL	444
Human MU-1	DRLKPPLADGEDWAGGLPWGGRSPGGVSESEAGSPLAGLDMDTFDSGFVG	496
MurineMU-1	DRLRLSFAKEGDWTADPTWRTGSPGGGSESEAGSP.PGLDMDTFDSGFAG	493
Human MU-1	SDCSSPVECDFTSPGDEGPPRSYLQWVV.IPPPLSSPGPQAS*	539
MurineMU-1	SDCGSPVET.....DEGPPRSYLQWVVRTPPFVDS.GAQSS.	529

Fig. 5

	1		50
humu	~~~MPRGWAA	PLLLLLL..LQ	GGWG..... CPDLVCYTDY LQTVICILEM
mousemu	~~~MPRGPVA	ALLLLI..LH	GAWG..... CLDLTCYTDY LWTITCVLET
humil2rbc	MAAPALSWRL	PLLILLPLA	TSWASAAVNG TSQFTCFYNS RANISCVWSQ
	51		100
humu	WN..LHPSTL	TLTWQDQYEE	LKDEATSCSL HRSAHNATHA TYTCHM....
mousemu	RS..PNPSIL	SLTWQDEYEE	LQDQETFCSL HRSGHNTTHI WYTCHM....
humil2rbc	DGALQDTSCQ	VHAWPDR...	.RRWNQTCELLPVSQA SWACNLILGA
	101		150
humu	.DVFHFMAADD	IFSVNITDQS	GN..YSQECG SFLLAESIKP APPFNVTVTF
mousemu	.RLSQFLSDE	VFIVNVTDQS	GN..NSQECG SFVLAESIKP APPLNVTVAF
humil2rbc	PDSQKLTVD	IVTLRVLCRE	GVRWRVMAIQ DFKPFENLRL MAPISLQVVH
	151		200
humu	..SGQYNISW	RSDYEDPAFY	MLKGKLQYEL QYRNRGDPWA VSPRRKLISV
mousemu	..SGRYDISW	DSAYDEPSNY	VLRGKLQYEL QYRNLRDPYA VRPVTKLISV
humil2rbc	VETHRCNISW	E..ISQASHY	FER.HLEFEA RTLSPGHTWE EAP...LLTL
	201		250
humu	DSRSVSLPL	EFKRDSSYEL	QVRAGPMPGS SYQGTWSEWS DPVIFQTQS.
mousemu	DSRNVSLPE	EFHKDSSYQL	QVRAAPQPGT SFRGTWSEWS DPVIFQTOA.
humil2rbc	KQKQEWICLE	TLTPDTQYEF	QVRVKPLQGE F..TTWSPWS QPLAFRTKPA
	251		300
humu	..EELKEGWN	PHLLLLL...	LLVIVFIPAF WSLKTHPLWR LWKKIWA.VP
mousemu	..GEPEAGWD	PHMLLLL...	AVLIIVL.VF MGLKIHLPWR LWKKIWAPVP
humil2rbc	ALGKDTIPWL	GHLVLVLSGA	FGFIILVYLL INCRNTGPW. LKKVLKCNTP
	301		350
humu	SPERFFMPLY	KGCSGDFKKW	VGAPFTGSSL ELGPWSPEVP STLEVYSCHP
mousemu	TPESFFQPLY	REHSGNFKKW	VNTPFTASSI ELVPQSSTTT SAL.....HL
humil2rbc	DPSKFFSQLS	SEHGGDVQKW	LSSPFPSSSF SPGGLAPEIS PLEVLERDKV
	351		400
humu	PRSPAKRLQL	TELQEPA..E	LVESDGVPKP SFW...PTAQ NSGGSAYSEE
mousemu	SLYPAKEKKF	PGLPGLE..E	QLECDGMSEP GHWCIIPLAA GOAVSAYSEE
humil2rbc	TQLLLQQDKV	PEPASLSSNH	SLTSCFTNQG YFFFHLPDAL EIEACQVYFT
	401		450
humu	RDRPYGLVSI	DTVTVLDAEG	PC...TWPCS CEDDGYPALD LDAGLEPSPG
mousemu	RDRPYGLVSI	DTVTVGDAEG	LC...VWPCS CEDDGYPAMN LDAGRESGPN
humil2rbc	YD.PYSEEDP	DEGVAGAPTG	SSPQPLQPLS GEDDAYCTF.PS
	451		500
humu	LEDPLLDAGT	TVLSCGCVSA	GSPGLGGPLG SLLDRLKPPL AD..GEDWAG
mousemu	SEDLLLVTD	AFLSCGCVSG	SGLRLGGSPG SLLDRLRLSF AK..EGDWT
humil2rbc	RDDLLLF.S.P	SLL..GGPSP	PSTAPGGS.G AGEERMPPSL QERVPRDWD
	501		550
humu	GLPWGGRSPG	GVSESEAGSP	LAGLDMDTFD SGFVGSDCSS PVECDFTSPG
mousemu	DPTWRTGSPG	GGSESEAGSP	.PGLDMDTFD SGFAGSDCGS PVET.....
humil2rbc	Q.PLGPPTPG	VPDLVDFQPP	P...ELVLRE AGEEVDPAG. PRE.GVSFPW
	551		588
humu	DEGPFRSYLR	QWVVI PPPLS	SPGPQAS*~ ~~~~~
mousemu	DEGPFRSYLR	QWVVRTPPPV	DSGAQSS~ ~~~~~
humil2rbc	SRPPGQGEFR	ALNARLPLNT	DAYLSLQELQ GQDPHTLV

Fig. 6

Signaling through MU-1




Fig. 7A

atg	aaa	ttc	tta	gtc	aac	ggt	gcc	ctt	ggt	ttt	atg	gtc	gtg	tac	att	48
Met	Lys	Phe	Leu	Val	Asn	Val	Ala	Leu	Val	Phe	Met	Val	Val	Tyr	Ile	
1				5				10						15		
tct	tac	atc	tat	gcc	ggc	agc	gga	cac	cac	cat	cat	cac	cac	ggt	agc	96
Ser	Tyr	Ile	Tyr	Ala	Gly	Ser	Gly	His	His	His	His	His	His	Gly	Ser	
			20					25					30			
ggc	gac	tat	aaa	gac	gat	gac	gat	aag	ggt	tcc	gga	tgc	ccc	gac	ctc	144
Gly	Asp	Tyr	Lys	Asp	Asp	Asp	Asp	Lys	Gly	Ser	Gly	Cys	Pro	Asp	Leu	
			35					40				45				
gtc	tgc	tac	acc	gat	tac	ctc	cag	acg	gtc	atc	tgc	atc	ctg	gaa	atg	192
Val	Cys	Tyr	Thr	Asp	Tyr	Leu	Gln	Thr	Val	Ile	Cys	Ile	Leu	Glu	Met	
	50					55					60					
tgg	aac	ctc	cac	ccc	agc	acg	ctc	acc	ctt	acc	tgg	caa	gac	cag	tat	240
Trp	Asn	Leu	His	Pro	Ser	Thr	Leu	Thr	Leu	Thr	Trp	Gln	Asp	Gln	Tyr	
65					70				75						80	
gaa	gag	ctg	aag	gac	gag	gcc	acc	tcc	tgc	agc	ctc	cac	agg	tcg	gcc	288
Glu	Glu	Leu	Lys	Asp	Glu	Ala	Thr	Ser	Cys	Ser	Leu	His	Arg	Ser	Ala	
				85					90					95		
cac	aat	gcc	acg	cat	gcc	acc	tac	acc	tgc	cac	atg	gat	gta	ttc	cac	336
His	Asn	Ala	Thr	His	Ala	Thr	Tyr	Thr	Cys	His	Met	Asp	Val	Phe	His	
			100					105					110			
ttc	atg	gcc	gac	gac	att	ttc	agt	gtc	aac	atc	aca	gac	cag	tct	ggc	384
Phe	Met		Asp	Asp	Ile	Phe	Ser	Val	Asn	Ile	Thr	Asp	Gln	Ser	Gly	
		115					120					125				
aac	tac	tcc	cag	gag	tgt	ggc	agc	ttt	ctc	ctg	gct	gag	agc	atc	aag	432
Asn	Tyr	Ser	Gln	Glu	Cys	Gly	Ser	Phe	Leu	Leu	Ala	Glu	Ser	Ile	Lys	
	130					135					140					
ccg	gct	ccc	cct	ttc	aac	gtg	act	gtg	acc	ttc	tca	gga	cag	tat	aat	480
Pro	Ala	Pro	Pro	Phe	Asn	Val	Thr	Val	Thr	Phe	Ser	Gly	Gln	Tyr	Asn	
145					150				155						160	
atc	tcc	tgg	cgc	tca	gat	tac	gaa	gac	cct	gcc	ttc	tac	atg	ctg	aag	528
Ile	Ser	Trp	Arg	Ser	Asp	Tyr	Glu	Asp	Pro	Ala	Phe	Tyr	Met	Leu	Lys	
				165					170					175		
ggc	aag	ctt	cag	tat	gag	ctg	cag	tac	agg	aac	cgg	gga	gac	ccc	tgg	576
Gly	Lys	Leu	Gln	Tyr	Glu	Leu	Gln	Tyr	Arg	Asn	Arg	Gly	Asp	Pro	Trp	
			180					185					190			
gct	gtg	agt	ccg	agg	aga	aag	ctg	atc	tca	gtg	gac	tca	aga	agt	gtc	624
Ala	Val	Ser	Pro	Arg	Arg	Lys	Leu	Ile	Ser	Val	Asp	Ser	Arg	Ser	Val	
		195					200					205				
tcc	ctc	ctc	ccc	ctg	gag	ttc	cgc	aaa	gac	tcg	agc	tat	gag	ctg	cag	672
Ser	Leu	Leu	Pro	Leu	Glu	Phe	Arg	Lys	Asp	Ser	Ser	Tyr	Glu	Leu	Gln	
	210					215					220					

Fig. 8A

gcggccgcac cacc	atg	ccg	cgt	ggc	tgg	gcc	gcc	ccc	ttg	ctc	ctg	ctg	50
	Met	Pro	Arg	Gly	Trp	Ala	Ala	Pro	Leu	Leu	Leu	Leu	
	1				5					10			
ctg ctc cag gga ggc tgg ggc tgc ccc gac ctc gtc tgc tac acc gat	98												
Leu Leu Gln Gly Gly Trp Gly Cys Pro Asp Leu Val Cys Tyr Thr Asp													
	15						20			25			
tac ctc cag acg gtc atc tgc atc ctg gaa atg tgg aac ctc cac ccc	146												
Tyr Leu Gln Thr Val Ile Cys Ile Leu Glu Met Trp Asn Leu His Pro													
	30					35			40				
agc acg ctc acc ctt acc tgg caa gac cag tat gaa gag ctg aag gac	194												
Ser Thr Leu Thr Leu Thr Trp Gln Asp Gln Tyr Glu Glu Leu Lys Asp													
	45				50			55				60	
gag gcc acc tcc tgc agc ctc cac agg tgc gcc cac aat gcc acg cat	242												
Glu Ala Thr Ser Cys Ser Leu His Arg Ser Ala His Asn Ala Thr His													
				65				70				75	
gcc acc tac acc tgc cac atg gat gta ttc cac ttc atg gcc gac gac	290												
Ala Thr Tyr Thr Cys His Met Asp Val Phe His Phe Met Ala Asp Asp													
			80				85				90		
att ttc agt gtc aac atc aca gac cag tct ggc aac tac tcc cag gag	338												
Ile Phe Ser Val Asn Ile Thr Asp Gln Ser Gly Asn Tyr Ser Gln Glu													
		95					100				105		
tgt ggc agc ttt ctc ctg gct gag agc atc aag ccg gct ccc cct ttc	386												
Cys Gly Ser Phe Leu Leu Ala Glu Ser Ile Lys Pro Ala Pro Pro Phe													
	110					115				120			
aac gtg act gtg acc ttc tca gga cag tat aat atc tcc tgg cgc tca	434												
Asn Val Thr Val Thr Phe Ser Gly Gln Tyr Asn Ile Ser Trp Arg Ser													
	125				130			135				140	
gat tac gaa gac cct gcc ttc tac atg ctg aag ggc aag ctt cag tat	482												
Asp Tyr Glu Asp Pro Ala Phe Tyr Met Leu Lys Gly Lys Leu Gln Tyr													
				145				150				155	
gag ctg cag tac agg aac cgg gga gac ccc tgg gct gtg agt ccg agg	530												
Glu Leu Gln Tyr Arg Asn Arg Gly Asp Pro Trp Ala Val Ser Pro Arg													
			160				165				170		
aga aag ctg atc tca gtg gac tca aga agt gtc tcc ctc ctc ccc ctg	578												
Arg Lys Leu Ile Ser Val Asp Ser Arg Ser Val Ser Leu Leu Pro Leu													
		175				180				185			
gag ttc cgc aaa gac tgc agc tat gag ctg cag gtg cgg gca ggg ccc	626												
Glu Phe Arg Lys Asp Ser Ser Tyr Glu Leu Gln Val Arg Ala Gly Pro													
	190					195			200				
atg cct ggc tcc tcc tac cag ggg acc tgg agt gaa tgg agt gac ccg	674												
Met Pro Gly Ser Ser Tyr Gln Gly Thr Trp Ser Glu Trp Ser Asp Pro													
	205				210			215				220	

Fig. 8B

gtc	atc	ttt	cag	acc	cag	tca	gag	gag	tta	aag	gaa	ggc	tgg	aac	ggc	722
Val	Ile	Phe	Gln	Thr	Gln	Ser	Glu	Glu	Leu	Lys	Glu	Gly	Trp	Asn	Gly	
				225					230					235		
tcc	ggc	tct	aga	gac	aaa	act	cac	aca	tgc	cca	ccg	tgc	cca	gca	cct	770
Ser	Gly	Ser	Arg	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	
			240					245					250			
gaa	ctc	ctg	ggg	gga	ccg	tca	gtc	ttc	ctc	ttc	ccc	cca	aaa	ccc	aag	818
Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	
		255					260					265				
gac	acc	ctc	atg	atc	tcc	cgg	acc	cct	gag	gtc	aca	tgc	gtg	gtg	gtg	866
Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	
	270					275					280					
gac	gtg	agc	cac	gaa	gac	cct	gag	gtc	aag	ttc	aac	tgg	tac	gtg	gac	914
Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	
285					290				295						300	
ggc	gtg	gag	gtg	cat	aat	gcc	aag	aca	aag	ccg	cgg	gag	gag	cag	tac	962
Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	
				305					310					315		
aac	agc	acg	tac	cgt	gtg	gtc	agc	gtc	ctc	acc	gtc	ctg	cac	cag	gac	1010
Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	
			320					325					330			
tgg	ctg	aat	ggc	aag	gag	tac	aag	tgc	aag	gtc	tcc	aac	aaa	gcc	ctc	1058
Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	
		335					340					345				
cca	gtc	ccc	atc	gag	aaa	acc	atc	tcc	aaa	gcc	aaa	ggg	cag	ccc	cga	1106
Pro	Val	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	
	350					355					360					
gaa	cca	cag	gtg	tac	acc	ctg	ccc	cca	tcc	cgg	gag	gag	atg	acc	aag	1154
Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	
365					370				375						380	
aac	cag	gtc	agc	ctg	acc	tgc	ctg	gtc	aaa	ggc	ttc	tat	ccc	agc	gac	1202
Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	
				385					390					395		
atc	gcc	gtg	gag	tgg	gag	agc	aat	ggg	cag	ccg	gag	aac	aac	tac	aag	1250
Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	
			400					405					410			
acc	acg	cct	ccc	gtg	ctg	gac	tcc	gac	ggc	tcc	ttc	ttc	ctc	tat	agc	1298
Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	
		415					420					425				
aag	ctc	acc	gtg	gac	aag	agc	agg	tgg	cag	cag	ggg	aac	gtc	ttc	tca	1346
Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	
	430					435					440					

Fig. 8C

[illegible]

Fig. 9A

gcggccgcac	cacc	atg Met 1	ccg Pro	cgt Arg	ggc Gly	tgg Trp 5	gcc Ala	gcc Ala	ccc Pro	ttg Leu 10	ctc Leu 10	ctg Leu	ctg Leu	50		
ctg Leu	ctc Leu	cag Gln 15	gga Gly	ggc Gly	tgg Trp	ggc Gly	tgc Cys 20	ccc Pro	gac Asp	ctc Leu	gtc Val	tgc Cys 25	tac Tyr	acc Thr	gat Asp	98
tac Tyr	ctc Leu 30	cag Gln	acg Thr	gtc Val	atc Ile	tgc Cys 35	atc Ile	ctg Leu	gaa Glu	atg Met	tgg Trp 40	aac Asn	ctc Leu	cac His	ccc Pro	146
agc Ser 45	acg Thr	ctc Leu	acc Thr	ctt Leu	acc Thr 50	tgg Trp	caa Gln	gac Asp	cag Gln	tat Tyr 55	gaa Glu	gag Glu	ctg Leu	aag Lys	gac Asp 60	194
gag Glu	gcc Ala	acc Thr	tcc Ser	tgc Cys 65	agc Ser	ctc Leu	cac His	agg Arg	tcg Ser 70	gcc Ala	cac His	aat Asn	gcc Ala	acg Thr 75	cat His	242
gcc Ala	acc Thr	tac Tyr	acc Thr 80	tgc Cys	cac His	atg Met	gat Asp	gta Val 85	ttc Phe	cac His	ttc Phe	atg Met	gcc Ala 90	gac Asp	gac Asp	290
att Ile	ttc Phe	agt Ser 95	gtc Val	aac Asn	atc Ile	aca Thr	gac Asp 100	cag Gln	tct Ser	ggc Gly	aac Asn	tac Tyr 105	tcc Ser	cag Gln	gag Glu	338
tgt Cys	ggc Gly 110	agc Ser	ttt Phe	ctc Leu	ctg Leu	gct Ala 115	gag Glu	agc Ser	atc Ile	aag Lys	ccg Pro 120	gct Ala	ccc Pro	cct Pro	ttc Phe	386
aac Asn 125	gtg Val	act Thr	gtg Val	acc Thr	ttc Phe 130	tca Ser	gga Gly	cag Gln	tat Tyr	aat Asn 135	atc Ile	tcc Ser	tgg Trp	cgc Arg	tca Ser 140	434
gat Asp	tac Tyr	gaa Glu	gac Asp	cct Pro 145	gcc Ala	ttc Phe	tac Tyr	atg Met	ctg Leu 150	aag Lys	ggc Gly	aag Lys	ctt Leu	cag Gln 155	tat Tyr	482
gag Glu	ctg Leu	cag Gln	tac Tyr 160	agg Arg	aac Asn	cgg Arg	gga Gly	gac Asp 165	ccc Pro	tgg Trp	gct Ala	gtg Val	agt Ser 170	ccg Pro	agg Arg	530
aga Arg	aag Lys	ctg Leu 175	atc Ile	tca Ser	gtg Val	gac Asp	tca Ser 180	aga Arg	agt Ser	gtc Val	tcc Ser	ctc Leu 185	ctc Leu	ccc Pro	ctg Leu	578
gag Glu	ttc Phe 190	cgc Arg	aaa Lys	gac Asp	tcg Ser	agc Ser 195	tat Tyr	gag Glu	ctg Leu	cag Gln	gtg Val 200	cgg Arg	gca Ala	ggg Gly	ccc Pro	626
atg Met 205	cct Pro	ggc Gly	tcc Ser	tcc Ser	tac Tyr 210	cag Gln	ggg Gly	acc Thr	tgg Trp	agt Ser 215	gaa Glu	tgg Trp	agt Ser	gac Asp	ccg Pro 220	674

Fig. 9B

gtc	atc	ttt	cag	acc	cag	tca	gag	gag	tta	aag	gaa	ggc	tgg	aac	ggc	722
Val	Ile	Phe	Gln	Thr	Gln	Ser	Glu	Glu	Leu	Lys	Glu	Gly	Trp	Asn	Gly	
			225						230					235		
tcc	ggc	tct	aga	gac	aaa	act	cac	aca	tgc	cca	ccg	tgc	cca	gca	cct	770
Ser	Gly	Ser	Arg	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	
			240					245					250			
gaa	ctc	ctg	ggg	gga	ccg	tca	gtc	ttc	ctc	ttc	ccc	cca	aaa	ccc	aag	818
Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	
		255					260					265				
gac	acc	ctc	atg	atc	tcc	cgg	acc	cct	gag	gtc	aca	tgc	gtg	gtg	gtg	866
Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	
	270					275					280					
gac	gtg	agc	cac	gaa	gac	cct	gag	gtc	aag	ttc	aac	tgg	tac	gtg	gac	914
Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	
285					290					295					300	
ggc	gtg	gag	gtg	cat	aat	gcc	aag	aca	aag	ccg	cgg	gag	gag	cag	tac	962
Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	
				305					310					315		
aac	agc	acg	tac	cgt	gtg	gtc	agc	gtc	ctc	acc	gtc	ctg	cac	cag	gac	1010
Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	
			320					325					330			
tgg	ctg	aat	ggc	aag	gag	tac	aag	tgc	aag	gtc	tcc	aac	aaa	gcc	ctc	1068
Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	
		335					340					345				
cca	gtc	ccc	atc	gag	aaa	acc	atc	tcc	aaa	gcc	aaa	ggg	cag	ccc	cga	1106
Pro	Val	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	
	350					355					360					
gaa	cca	cag	gtg	tac	acc	ctg	ccc	cca	tcc	cgg	gag	gag	atg	acc	aag	1154
Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	
365					370					375					380	
aac	cag	gtc	agc	ctg	acc	tgc	ctg	gtc	aaa	ggc	ttc	tat	ccc	agc	gac	1202
Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	
				385					390					395		
atc	gcc	gtg	gag	tgg	gag	agc	aat	ggg	cag	ccg	gag	aac	aac	tac	aag	1250
Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	
			400					405					410			
acc	acg	cct	ccc	gtg	ctg	gac	tcc	gac	ggc	tcc	ttc	ttc	ctc	tat	agc	1298
Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	
		415					420					425				
aag	ctc	acc	gtg	gac	aag	agc	agg	tgg	cag	cag	ggg	aac	gtc	ttc	tca	1346
Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	
	430					435					440					

Fig. 9C

tgc tcc gtg atg cat gag gct ctg cac aac cac tac acg cag aag agc	1394
Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser	
445 450 455 460	
ctc tcc ctg tcc ccg ggt aaa tca gga atg gca tca atg aca gga ggt	1442
Leu Ser Leu Ser Pro Gly Lys Ser Gly Met Ala Ser Met Thr Gly Gly	
465 470 475	
caa caa atg ggt tct gga tct cat cat cat cat cat cat tct gga ggt	1490
Gln Gln Met Gly Ser Gly Ser His His His His His His Ser Gly Gly	
480 485 490	
tgagaattc	1499

Fig. 10A

gcggccgcac cacc atg ccg cgt ggc tgg gcc gcc ccc ttg ctc ctg ctg	50
Met Pro Arg Gly Trp Ala Ala Pro Leu Leu Leu	
1 5 10	
ctg ctc cag gga ggc tgg ggc tgc ccc gac ctc gtc tgc tac acc gat	98
Leu Leu Gln Gly Gly Trp Gly Cys Pro Asp Leu Val Cys Tyr Thr Asp	
15 20 25	
tac ctc cag acg gtc atc tgc atc ctg gaa atg tgg aac ctc cac ccc	146
Tyr Leu Gln Thr Val Ile Cys Ile Leu Glu Met Trp Asn Leu His Pro	
30 35 40	
agc acg ctc acc ctt acc tgg caa gac cag tat gaa gag ctg aag gac	194
Ser Thr Leu Thr Leu Thr Trp Gln Asp Gln Tyr Glu Glu Leu Lys Asp	
45 50 55 60	
gag gcc acc tcc tgc agc ctc cac agg tgc gcc cac aat gcc acg cat	242
Glu Ala Thr Ser Cys Ser Leu His Arg Ser Ala His Asn Ala Thr His	
65 70 75	
gcc acc tac acc tgc cac atg gat gta ttc cac ttc atg gcc gac gac	290
Ala Thr Tyr Thr Cys His Met Asp Val Phe His Phe Met Ala Asp Asp	
80 85 90	
att ttc agt gtc aac atc aca gac cag tct ggc aac tac tcc cag gag	338
Ile Phe Ser Val Asn Ile Thr Asp Gln Ser Gly Asn Tyr Ser Gln Glu	
95 100 105	
tgt ggc agc ttt ctc ctg gct gag agc atc aag ccg gct ccc cct ttc	386
Cys Gly Ser Phe Leu Leu Ala Glu Ser Ile Lys Pro Ala Pro Pro Phe	
110 115 120	
aac gtg act gtg acc ttc tca gga cag tat aat atc tcc tgg cgc tca	434
Asn Val Thr Val Thr Phe Ser Gly Gln Tyr Asn Ile Ser Trp Arg Ser	
125 130 135 140	
gat tac gaa gac cct gcc ttc tac atg ctg aag ggc aag ctt cag tat	482
Asp Tyr Glu Asp Pro Ala Phe Tyr Met Leu Lys Gly Lys Leu Gln Tyr	
145 150 155	
gag ctg cag tac agg aac cgg gga gac ccc tgg gct gtg agt ccg agg	530
Glu Leu Gln Tyr Arg Asn Arg Gly Asp Pro Trp Ala Val Ser Pro Arg	
160 165 170	
aga aag ctg atc tca gtg gac tca aga agt gtc tcc ctc ctc ccc ctg	578
Arg Lys Leu Ile Ser Val Asp Ser Arg Ser Val Ser Leu Leu Pro Leu	
175 180 185	
gag ttc cgc aaa gac tgc agc tat gag ctg cag gtg cgg gca ggg ccc	626
Glu Phe Arg Lys Asp Ser Ser Tyr Glu Leu Gln Val Arg Ala Gly Pro	
190 195 200	
atg cct ggc tcc tcc tac cag ggg acc tgg agt gaa tgg agt gac ccg	674
Met Pro Gly Ser Ser Tyr Gln Gly Thr Trp Ser Glu Trp Ser Asp Pro	
205 210 215 220	

Fig. 10B

gtc atc ttt cag acc cag tca gag gag tta aag gaa ggc tgg aac ggc	722
Val Ile Phe Gln Thr 225	
tcc ggc tct aga gac aaa act cac aca tgc cca ccg tgc cca gca cct	770
Ser Gly Ser Arg Asp Lys Thr His Thr 245	
gaa gcc ctg ggg gca ccg tca gtc ttc ctc ttc ccc cca aaa ccc aag	818
Glu Ala Leu Gly Ala Pro Ser Val 260	
gac acc ctc atg atc tcc cgg acc cct gag gtc aca tgc gtg gtg gtg	866
Asp Thr 270	
gac gtg agc cac gaa gac cct gag gtc aag ttc aac tgg tac gtg gac	914
Asp Val 285	
ggc gtg gag gtg cat aat gcc aag aca aag ccg cgg gag gag cag tac	962
Gly Val 305	
aac agc acg tac cgt gtg gtc agc gtc ctc acc gtc ctg cac cag gac	1010
Asn Ser Thr Tyr Arg Val Val Ser Val 325	
tgg ctg aat ggc aag gag tac aag tgc aag gtc tcc aac aaa gcc ctc	1058
Trp Leu Asn 335	
cca gcc ccc atc gag aaa acc atc tcc aaa gcc aaa ggg cag ccc cga	1106
Pro Ala Pro Ile Glu Lys Thr 355	
gaa cca cag gtg tac acc ctg ccc cca tcc cgg gag gag atg acc aag	1154
Glu Pro Gln Val Tyr Thr 370	
aac cag gtc agc ctg acc tgc ctg gtc aaa ggc ttc tat ccc agc gac	1202
Asn Gln Val Ser Leu Thr Cys Leu Val 390	
atc gcc gtg gag tgg gag agc aat ggg cag ccg gag aac aac tac aag	1250
Ile Ala Val Glu Trp Glu Ser Asn Gly 405	
acc acg cct ccc gtg ctg gac tcc gac ggc tcc ttc ttc ctc tat agc	1298
Thr Thr 415	
aag ctc acc gtg gac aag agc agg tgg cag cag ggg aac gtc ttc tca	1346
Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser	
430 435 440	

tgc	tcc	gtg	atg	cat	gag	gct	ctg	cac	aac	cac	tac	acg	cag	aag	agc	1394
Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	
445					450					455					460	
ctc	tcc	ctg	tcc	ccg	ggg	aaa	tgagtgaatt.	c								1426
Leu	Ser	Leu	Ser	Pro	Gly	Lys										
				465												

Fig. 11A

atg Met 1	ccg Pro	cgt Arg	ggc Gly	tgg Trp 5	gcc Ala	gcc Ala	ccc Pro	ttg Leu 10	ctc Leu 10	ctg Leu	ctg Leu	ctg Leu	ctc Leu	cag Gln 15	gga Gly	48
ggc Gly	tgg Trp	ggc Gly	tgc Cys 20	ccc Pro	gac Asp	ctc Leu	gtc Val	tgc Cys 25	tac Tyr	acc Thr	gat Asp	tac Tyr	ctc Leu 30	cag Gln	acg Thr	96
gtc Val	atc Ile	tgc Cys 35	atc Ile	ctg Leu	gaa Glu	atg Met 40	tgg Trp 40	aac Asn	ctc Leu	cac His	ccc Pro	agc Ser 45	acg Thr	ctc Leu	acc Thr	144
ctt Leu	acc Thr 50	tgg Trp	caa Gln	gac Asp	cag Gln	tat Tyr 55	gaa Glu	gag Glu	ctg Leu	aag Lys	gac Asp 60	gag Glu	gcc Ala	acc Thr	tcc Ser	192
tgc Cys 65	agc Ser	ctc Leu	cac His	agg Arg	tcg Ser 70	gcc Ala	cac His	aat Asn	gcc Ala	acg Thr 75	cat His	gcc Ala	acc Thr	tac Tyr	acc Thr 80	240
tgc Cys	cac His	atg Met	gat Asp	gta Val 85	ttc Phe	cac His	ttc Phe	atg Met	gcc Ala 90	gac Asp	gac Asp	att Ile	ttc Phe	agt Ser 95	gtc Val	288
aac Asn	atc Ile	aca Thr	gac Asp 100	cag Gln	tct Ser	ggc Gly	aac Asn	tac Tyr 105	tcc Ser	cag Gln	gag Glu	tgt Cys	ggc Gly 110	agc Ser	ttt Phe	336
ctc Leu	ctg Leu	gct Ala 115	gag Glu	agc Ser	atc Ile	aag Lys	ccg Pro 120	gct Ala	ccc Pro	cct Pro	ttc Phe	aac Asn 125	gtg Val	act Thr	gtg Val	384
acc Thr 130	ttc Phe	tca Ser	gga Gly	cag Gln	tat Tyr	aat Asn 135	atc Ile	tcc Ser	tgg Trp	cgc Arg	tca Ser 140	gat Asp	tac Tyr	gaa Glu	gac Asp	432
cct Pro 145	gcc Ala	ttc Phe	tac Tyr	atg Met	ctg Leu 150	aag Lys	ggc Gly	aag Lys	ctt Leu	cag Gln 155	tat Tyr	gag Glu	ctg Leu	cag Gln	tac Tyr 160	480
agg Arg	aac Asn	cgg Arg	gga Gly	gac Asp 165	ccc Pro	tgg Trp	gct Ala	gtg Val	agt Ser 170	ccg Pro	agg Arg	aga Arg	aag Lys	ctg Leu 175	atc Ile	528
tca Ser	gtg Val	gac Asp	tca Ser 180	aga Arg	agt Ser	gtc Val	tcc Ser	ctc Leu 185	ctc Leu	ccc Pro	ctg Leu	gag Glu	ttc Phe 190	cgc Arg	aaa Lys	576
gac Asp	tcg Ser	agc Ser 195	tat Tyr	gag Glu	ctg Leu	cag Gln	gtg Val 200	cgg Arg	gca Ala	ggg Gly	ccc Pro	atg Met 205	cct Pro	ggc Gly	tcc Ser	624
tcc Ser	tac Tyr 210	cag Gln	ggg Gly	acc Thr	tgg Trp	agt Ser 215	gaa Glu	tgg Trp	agt Ser	gac Asp	ccg Pro 220	gtc Val	atc Ile	ttt Phe	cag Gln	672

Fig. 11B

[illegible]

Fig. 12A

atg Met 1	ccg Pro	cgt Arg	ggc Gly 5	tgg Trp	gcc Ala	gcc Ala	ccc Pro	ttg Leu 10	ctc Leu 10	ctg Leu	ctg Leu	ctc Leu	cag Gln 15	gga Gly	48	
ggc Gly	tgg Trp	ggc Gly	tgc Cys 20	ccc Pro	gac Asp	ctc Leu	gtc Val	tgc Cys 25	tac Tyr	acc Thr	gat Asp	tac Tyr	ctc Leu 30	cag Gln	acg Thr	96
gtc Val	atc Ile	tgc Cys 35	atc Ile	ctg Leu	gaa Glu	atg Met	tgg Trp 40	aac Asn	ctc Leu	cac His	ccc Pro	agc Ser 45	acg Thr	ctc Leu	acc Thr	144
ctt Leu	acc Thr 50	tgg Trp	caa Gln	gac Asp	cag Gln	tat Tyr 55	gaa Glu	gag Glu	ctg Leu	aag Lys	gac Asp 60	gag Glu	gcc Ala	acc Thr	tcc Ser	192
tgc Cys 65	agc Ser	ctc Leu	cac His	agg Arg	tcg Ser 70	gcc Ala	cac His	aat Asn	gcc Ala	acg Thr 75	cat His	gcc Ala	acc Thr	tac Tyr	acc Thr 80	240
tgc Cys	cac His	atg Met	gat Asp	gta Val 85	ttc Phe	cac His	ttc Phe	atg Met	gcc Ala 90	gac Asp	gac Asp	att Ile	ttc Phe	agt Ser 95	gtc Val	288
aac Asn	atc Ile	aca Thr	gac Asp 100	cag Gln	tct Ser	ggc Gly	aac Asn	tac Tyr 105	tcc Ser	cag Gln	gag Glu	tgt Cys	ggc Gly 110	agc Ser	ttt Phe	336
ctc Leu	ctg Leu	gct Ala 115	gag Glu	agc Ser	atc Ile	aag Lys	ccg Pro 120	gct Ala	ccc Pro	cct Pro	ttc Phe	aac Asn 125	gtg Val	act Thr	gtg Val	384
acc Thr	ttc Phe 130	tca Ser	gga Gly	cag Gln	tat Tyr	aat Asn 135	atc Ile	tcc Ser	tgg Trp	cgc Arg	tca Ser 140	gat Asp	tac Tyr	gaa Glu	gac Asp	432
cct Pro 145	gcc Ala	ttc Phe	tac Tyr	atg Met	ctg Leu 150	aag Lys	ggc Gly	aag Lys	ctt Leu	cag Gln 155	tat Tyr	gag Glu	ctg Leu	cag Gln	tac Tyr 160	480
agg Arg	aac Asn	cgg Arg	gga Gly 165	gac Asp	ccc Pro	tgg Trp	gct Ala	gtg Val	agt Ser 170	ccg Pro	agg Arg	aga Arg	aag Lys	ctg Leu 175	atc Ile	528
tca Ser	gtg Val	gac Asp 180	tca Ser	aga Arg	agt Ser	gtc Val	tcc Ser	ctc Leu 185	ctc Leu	ccc Pro	ctg Leu	gag Glu	ttc Phe 190	cgc Arg	aaa Lys	576
gac Asp	tcg Ser	agc Ser 195	tat Tyr	gag Glu	ctg Leu	cag Gln	gtg Val 200	cgg Arg	gca Ala	ggg Gly	ccc Pro	atg Met 205	cct Pro	ggc Gly	tcc Ser	624
tcc Ser	tac Tyr 210	cag Gln	ggg Gly	acc Thr	tgg Trp	agt Ser 215	gaa Glu	tgg Trp	agt Ser	gac Asp	ccg Pro 220	gtc Val	atc Ile	ttt Phe	cag Gln	672

Fig. 12B

```

acc cag tca gag gag tta aag gaa ggc tgg aac gat gac gat gac aag
Thr Gln Ser Glu Glu Leu Lys Glu Gly Trp Asn Asp Asp Asp Asp Lys
225                               230                               235                               240

ggc tcc ggc gac aaa act cac aca tgc cca ccg tgc cca gca cct gaa
Gly Ser Gly Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu
                               245                               250                               255

gcc ctg ggg gca ccg tca gtc ttc ctc ttc ccc cca aaa ccc aag gac
Ala Leu Gly Ala Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp
                               260                               265                               270

acc ctc atg atc tcc cgg acc cct gag gtc aca tgc gtg gtg gtg gac
Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp
                               275                               280                               285

gtg agc cac gaa gac cct gag gtc aag ttc aac tgg tac gtg gac ggc
Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly
290                               295                               300

gtg gag gtg cat aat gcc aag aca aag ccg cgg gag gag cag tac aac
Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn
305                               310                               315                               320

agc acg tac cgt gtg gtc agc gtc ctc acc gtc ctg cac cag gac tgg
Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp
                               325                               330                               335

ctg aat ggc aag gag tac aag tgc aag gtc tcc aac aaa gcc ctc cca
Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro
                               340                               345                               350

gcc ccc atc gag aaa acc atc tcc aaa gcc aaa ggg cag ccc cga gaa
Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu
                               355                               360                               365

cca cag gtg tac acc ctg ccc cca tcc ccg gag gag atg acc aag aac
Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn
370                               375                               380

cag gtc agc ctg acc tgc ctg gtc aaa ggc ttc tat ccc agc gac atc
Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile
385                               390                               395                               400

gcc gtg gag tgg gag agc aat ggg cag ccg gag aac aac tac aag acc
Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr
405                               410                               415

acg cct ccc gtg ctg gac tcc gac ggc tcc ttc ttc ctc tat agc aag
Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys
420                               425                               430

ctc acc gtg gac aag agc agg tgg cag cag ggg aac gtc ttc tca tgc
Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys
435                               440                               445

```


Fig. 13A

atg ccc cgg ggc cca gtg gct gcc tta ctc ctg ctg att ctc cat gga	48
Met Pro Arg Gly Pro Val Ala Ala Leu Leu Leu Ile Leu His Gly	
1 5 10 15	
gct tgg agc tgc ctg gac ctc act tgc tac act gac tac ctc tgg acc	96
Ala Trp Ser Cys Leu Asp Leu Thr Cys Tyr Thr Asp Tyr Leu Trp Thr	
20 25 30	
atc acc tgt gtc ctg gag aca cgg agc ccc aac ccc agc ata ctc agt	144
Ile Thr Cys Val Leu Glu Thr Arg Ser Pro Asn Pro Ser Ile Leu Ser	
35 40 45	
ctc acc tgg caa gat gaa tat gag gaa ctt cag gac caa gag acc ttc	192
Leu Thr Trp Gln Asp Glu Tyr Glu Glu Leu Gln Asp Gln Glu Thr Phe	
50 55 60	
tgc agc cta cac agg tct ggc cac aac acc aca cat ata tgg tac acg	240
Cys Ser Leu His Arg Ser Gly His Asn Thr Thr His Ile Trp Tyr Thr	
65 70 75 80	
tgc cat atg cgc ttg tct caa ttc ctg tcc gat gaa gtt ttc att gtc	288
Cys His Met Arg Leu Ser Gln Phe Leu Ser Asp Glu Val Phe Ile Val	
85 90 95	
aat gtg acg gac cag tct ggc aac aac tcc caa gag tgt ggc agc ttt	336
Asn Val Thr Asp Gln Ser Gly Asn Asn Ser Gln Glu Cys Gly Ser Phe	
100 105 110	
gtc ctg gct gag agc atc aaa cca gct ccc ccc ttg aac gtg act gtg	384
Val Leu Ala Glu Ser Ile Lys Pro Ala Pro Pro Leu Asn Val Thr Val	
115 120 125	
gcc ttc tca gga cgc tat gat atc tcc tgg gac tca gct tat gac gaa	432
Ala Phe Ser Gly Arg Tyr Asp Ile Ser Trp Asp Ser Ala Tyr Asp Glu	
130 135 140	
ccc tcc aac tac gtg ctg agg ggc aag cta caa tat gag ctg cag tat	480
Pro Ser Asn Tyr Val Leu Arg Gly Lys Leu Gln Tyr Glu Leu Gln Tyr	
145 150 155 160	
cgg aac ctc aga gac ccc tat gct gtg agg ccg gtg acc aag ctg atc	528
Arg Asn Leu Arg Asp Pro Tyr Ala Val Arg Pro Val Thr Lys Leu Ile	
165 170 175	
tca gtg gac tca aga aac gtc tct ctt ctc cct gaa gag ttc cac aaa	576
Ser Val Asp Ser Arg Asn Val Ser Leu Leu Pro Glu Glu Phe His Lys	
180 185 190	
gat tct agc tac cag ctg cag gtg cgg gca gcg cct cag cca ggc act	624
Asp Ser Ser Tyr Gln Leu Gln Val Arg Ala Ala Pro Gln Pro Gly Thr	
195 200 205	
tca ttc agg ggg acc tgg agt gag tgg agt gac ccc gtc atc ttt cag	672
Ser Phe Arg Gly Thr Trp Ser Glu Trp Ser Asp Pro Val Ile Phe Gln	
210 215 220	

Fig. 13B

acc	cag	gct	ggg	gag	ccc	gag	gca	ggc	tgg	gac	ggc	tcc	ggc	tct	aga	720
Thr	Gln	Ala	Gly	Glu	Pro	Glu	Ala	Gly	Trp	Asp	Gly	Ser	Gly	Ser	Arg	
225					230					235					240	
gagccccgcg	gaccgacaat	caagccctgt	cctccatgca	aatgcccagg	taagtcacta	780										
gaccagagct	ccactcccgg	gagaatggta	agtgtataa	acatccctgc	actagaggat	840										
aagccatgta	cagatccatt	tccatctctc	ctcatcagca	cctaacctcg	aggggtggacc	900										
atccgtcttc	atcttccctc	caaagatcaa	ggatgtactc	atgatctccc	tgagcccat	960										
agtcacatgt	gtgggtgggtg	atgtgagcga	ggatgacca	gatgtccaga	tcagctgggtt	1020										
tgtgaacaac	gtggaagtac	acacagctca	gacacaaacc	catagagagg	attacaacag	1080										
tactctccgg	gtggtcagtg	ccctcccat	ccagcaccag	gactggatga	gtggcaaggc	1140										
tttcgcatgc	gccgtcaaca	aaaagacct	cccagcgcgc	atcgagagaa	ccatctcaaa	1200										
acccaaaggt	gagagctgca	gcctgactgc	atgggggctg	ggatgggcat	aaggataaag	1260										
gtctgtgtgg	acagccttct	gcttcagcca	tgacctttgt	gtatgtttct	accctcacag	1320										
ggtcagtaag	agctccacag	gtatatgtct	tgctccacc	agaagaagag	atgactaaga	1380										
aacaggtcac	tctgacctgc	atgggtcacag	acttcatgcc	tgaagacatt	tacgtggagt	1440										
ggaccaacaa	cgggaaaaca	gagctaaact	acaagaacac	tgaaccagtc	ctggactctg	1500										
atggttctta	cttcatgtac	agcaagctga	gagtggaaaa	gaagaactgg	gtggaaagaa	1560										
atagctactc	ctgttcagtg	gtccacgagg	gtctgcacaa	tcaccacacg	actaagagct	1620										
tctcccggac	tccgggtaaa	tgagctcagc	accacaaaaa	ctctcaggtc	caaagagaca	1680										
cccacactca	tctccatgct	tcccttgat	aaataaagca	cccagcaatg	cctgggacca	1740										
tgtaatagga	attc					1754										

Fig. 14A

ctgcaggtcg	acaccacc	atg	ccc	cgg	ggc	cca	gtg	gct	gcc	tta	ctc	ctg	51
		Met	Pro	Arg	Gly	Pro	Val	Ala	Ala	Leu	Leu	Leu	
		1				5				10			
ctg	att	ctc	cat	gga	gct	tgg	agc	tgc	ctg	gac	ctc	act	99
Leu	Ile	Leu	His	Gly	Ala	Trp	Ser	Cys	Leu	Asp	Leu	Thr	
			15					20				25	
gac	tac	ctc	tgg	acc	atc	acc	tgt	gtc	ctg	gag	aca	cgg	147
Asp	Tyr	Leu	Trp	Thr	Ile	Thr	Cys	Val	Leu	Glu	Thr	Arg	
		30					35				40	Ser	
ccc	agc	ata	ctc	agt	ctc	acc	tgg	caa	gat	gaa	tat	gag	195
Pro	Ser	Ile	Leu	Ser	Leu	Thr	Trp	Gln	Asp	Glu	Tyr	Glu	
	45					50				55		Glu	
gac	caa	gag	acc	ttc	tgc	agc	cta	cac	agg	tct	ggc	cac	243
Asp	Gln	Glu	Thr	Phe	Cys	Ser	Leu	His	Arg	Ser	Gly	His	
60					65					70		Asn	
cat	ata	tgg	tac	acg	tgc	cat	atg	cgc	ttg	tct	caa	ttc	291
His	Ile	Trp	Tyr	Thr	Cys	His	Met	Arg	Leu	Ser	Gln	Phe	
				80					85			Leu	
gaa	gtt	ttc	att	gtc	aat	gtg	acg	gac	cag	tct	ggc	aac	339
Glu	Val	Phe	Ile	Val	Asn	Val	Thr	Asp	Gln	Ser	Gly	Asn	
			95					100				Asn	
gag	tgt	ggc	agc	ttt	gtc	ctg	gct	gag	agc	atc	aaa	cca	387
Glu	Cys	Gly	Ser	Phe	Val	Leu	Ala	Glu	Ser	Ile	Lys	Pro	
		110					115					Ala	
ttg	aac	gtg	act	gtg	gcc	ttc	tca	gga	cgc	tat	gat	atc	435
Leu	Asn	Val	Thr	Val	Ala	Phe	Ser	Gly	Arg	Tyr	Asp	Ile	
	125					130					135	Ser	
tca	gct	tat	gac	gaa	ccc	tcc	aac	tac	gtg	ctg	agg	ggc	483
Ser	Ala	Tyr	Asp	Glu	Pro	Ser	Asn	Tyr	Val	Leu	Arg	Gly	
140					145					150		Lys	
tat	gag	ctg	cag	tat	cgg	aac	ctc	aga	gac	ccc	tat	gct	531
Tyr	Glu	Leu	Gln	Tyr	Arg	Asn	Leu	Arg	Asp	Pro	Tyr	Ala	
				160					165			Val	
gtg	acc	aag	ctg	atc	tca	gtg	gac	tca	aga	aac	gtc	tct	579
Val	Thr	Lys	Leu	Ile	Ser	Val	Asp	Ser	Arg	Asn	Val	Ser	
			175				180					Leu	
gaa	gag	ttc	cac	aaa	gat	tct	agc	tac	cag	ctg	cag	gtg	627
Glu	Glu	Phe	His	Lys	Asp	Ser	Ser	Tyr	Gln	Leu	Gln	Val	
		190					195					Arg	
cct	cag	cca	ggc	act	tca	ttc	agg	ggg	acc	tgg	agt	gag	675
Pro	Gln	Pro	Gly	Thr	Ser	Phe	Arg	Gly	Thr	Trp	Ser	Glu	
	205					210						Trp	

Fig. 14B

ccc gtc atc ttt cag acc cag gct ggg gag ccc gag gca ggc tgg gac	723
Pro Val Ile Phe Gln Thr Gln Ala Gly Glu Pro Glu Ala Gly Trp Asp	
220 225 230 235	
ggc agc gga cac cac cat cat cac cac ggt agc ggc gac tat aaa gac	771
Gly Ser Gly His His His His His Gly Ser Gly Asp Tyr Lys Asp	
240 245 250	
gat gac gat aag tagtgagaat tc	795
Asp Asp Asp Lys	
255	

atg Met 1	aaa Lys	ttc Phe	tta Leu	gtc Val 5	aac Asn	gtt Val	gcc Ala	ctt Leu	gtt Val 10	ttt Phe	atg Met	gtc Val	gtg Val	tac Tyr 15	att Ile	48
tct Ser	tac Tyr	atc Ile	tat Tyr 20	gcc Ala	ggc Gly	agc Ser	gga Gly	cac His 25	cac His	cat His	cat His	cac His	cac His 30	ggg Gly	agc Ser	96
ggc Gly	gac Asp	tat Tyr 35	aaa Lys	gac Asp	gat Asp	gac Asp	gat Asp 40	aag Lys	ggg Gly	tcc Ser	gga Gly	tgc Cys 45	ctg Leu	gac Asp	ctc Leu	144
act Thr	tgc Cys 50	tac Tyr	act Thr	gac Asp	tac Tyr	ctc Leu 55	tgg Trp	acc Thr	atc Ile	acc Thr	tgt Cys 60	gtc Val	ctg Leu	gag Glu	aca Thr	192
cgg Arg 65	agc Ser	ccc Pro	aac Asn	ccc Pro	agc Ser 70	ata Ile	ctc Leu	agt Ser	ctc Leu	acc Thr 75	tgg Trp	caa Gln	gat Asp	gaa Glu	tat Tyr 80	240
gag Glu	gaa Glu	ctt Leu	cag Gln	gac Asp 85	caa Gln	gag Glu	acc Thr	ttc Phe	tgc Cys 90	agc Ser	cta Leu	cac His	agg Arg	tct Ser 95	ggc Gly	288
cac His	aac Asn	acc Thr	aca Thr 100	cat His	ata Ile	tgg Trp	tac Tyr	acg Thr 105	tgc Cys	cat His	atg Met	cgc Arg	ttg Leu 110	tct Ser	caa Gln	336
ttc Phe	ctg Leu	tcc Ser 115	gat Asp	gaa Glu	gtt Val	ttc Phe	att Ile 120	gtc Val	aat Asn	gtg Val	acg Thr	gac Asp 125	cag Gln	tct Ser	ggc Gly	384
aac Asn	aac Asn 130	tcc Ser	caa Gln	gag Glu	tgt Cys	ggc Gly 135	agc Ser	ttt Phe	gtc Val	ctg Leu	gct Ala 140	gag Glu	agc Ser	atc Ile	aaa Lys	432
cca Pro 145	gct Ala	ccc Pro	ccc Pro	ttg Leu	aac Asn 150	gtg Val	act Thr	gtg Val	gcc Ala	ttc Phe 155	tca Ser	gga Gly	cgc Arg	tat Tyr	gat Asp 160	480
atc Ile	tcc Ser	tgg Trp	gac Asp	tca Ser 165	gct Ala	tat Tyr	gac Asp	gaa Glu	ccc Pro 170	tcc Ser	aac Asn	tac Tyr	gtg Val	ctg Leu 175	agg Arg	528
ggc Gly	aag Lys	cta Leu	caa Gln 180	tat Tyr	gag Glu	ctg Leu	cag Gln	tat Tyr 185	cgg Arg	aac Asn	ctc Leu	aga Arg	gac Asp 190	ccc Pro	tat Tyr	576
gct Ala	gtg Val	agg Arg 195	ccg Pro	gtg Val	acc Thr	aag Lys	ctg Leu 200	atc Ile	tca Ser	gtg Val	gac Asp	tca Ser 205	aga Arg	aac Asn	gtc Val	624
tct Ser	ctt Leu 210	ctc Leu	cct Pro	gaa Glu	gag Glu	ttc Phe 215	cac His	aaa Lys	gat Asp	tct Ser	agc Ser 220	tac Tyr	cag Gln	ctg Leu	cag Gln	672

gtg	cgg	gca	gcg	ctt	cag	cca	ggc	act	tca	ttc	agg	ggg	acc	tgg	agt	720
Val	Arg	Ala	Ala	Pro	Gln	Pro	Gly	Thr	Ser	Phe	Arg	Gly	Thr	Trp	Ser	
225					230					235					240	
gag	tgg	agt	gac	ccc	gtc	atc	ttt	cag	acc	cag	gct	ggg	gag	ccc	gag	768
Glu	Trp	Ser	Asp	Pro	Val	Ile	Phe	Gln	Thr	Gln	Ala	Gly	Glu	Pro	Glu	
				245					250					255		
gca	ggc	tgg	gac	tagtgagaat tc										792		
Ala	Gly	Trp	Asp													
			260													

Fig. 16A




Fig. 16B




Fig. 17A




Fig. 17B




Fig. 17C




Fig. 17D




Fig. 18A




Fig. 18B




Fig. 19A




Fig. 19B




Fig. 19C




Fig. 19D




Fig. 20A




Fig. 20B






Fig. 20C**Fig. 20D**

Fig. 21A




Fig. 21B




Fig. 22A

Week postinfection	Group	Worm Pairs	Total Worms	Eggs/worm pair (in thousands)
9	WT (<i>n</i> =29)	6.53 ± 0.63	15.53 ± 1.55	7.29 ± 0.49
	KO (<i>n</i> =26)	5.00 ± 0.70	12.30 ± 1.74	9.63 ± 1.16
12	WT (<i>n</i> =19)	3.74 ± 0.60	10.44 ± 2.08	14.75 ± 1.91
	KO (<i>n</i> =19)	4.05 ± 0.62	9.89 ± 1.77	14.50 ± 1.95



Fig. 22B**Fig. 22C**

Fig. 22D




Fig. 22E




Fig. 22F









Fig. 23A**Fig. 23B****Fig. 23C****Fig. 23D**

Fig. 24



SEQUENCE LISTING

<110> Wyeth
The Government of the United States of America as
Represented by the Secretary of the Department of Health and
Human Services
The President and Fellows of Harvard College

<120> METHODS FOR TREATING AND PREVENTING FIBROSIS

<130> 01997.043800

<150> 60/671,374

<151> 2005-04-14

<160> 33

<170> PatentIn version 3.3

<210> 1

<211> 2665

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (236)..(1849)

<400> 1

gtcgactgga ggcccagctg cccgtcatca gagtgacagg tcttatgaca gcctgattgg	60
tgactcgggc tgggtgtgga ttctcacccc aggcctctgc ctgctttctc agaccctcat	120
ctgtcacccc cacgctgaac ccagctgcca cccccagaag cccatcagac tgcccccagc	180
acacggaatg gatttctgag aaagaagccg aaacagaagg cccgtgggag tcagc atg	238
Met	
1	
ccg cgt ggc tgg gcc gcc ccc ttg ctc ctg ctg ctg ctc cag gga ggc	286
Pro Arg Gly Trp Ala Ala Pro Leu Leu Leu Leu Leu Gln Gly Gly	
5 10 15	
tgg ggc tgc ccc gac ctc gtc tgc tac acc gat tac ctc cag acg gtc	334
Trp Gly Cys Pro Asp Leu Val Cys Tyr Thr Asp Tyr Leu Gln Thr Val	
20 25 30	
atc tgc atc ctg gaa atg tgg aac ctc cac ccc agc acg ctc acc ctt	382
Ile Cys Ile Leu Glu Met Trp Asn Leu His Pro Ser Thr Leu Thr Leu	
35 40 45	
acc tgg caa gac cag tat gaa gag ctg aag gac gag gcc acc tcc tgc	430
Thr Trp Gln Asp Gln Tyr Glu Glu Leu Lys Asp Glu Ala Thr Ser Cys	
50 55 60 65	
agc ctc cac agg tcg gcc cac aat gcc acg cat gcc acc tac acc tgc	478
Ser Leu His Arg Ser Ala His Asn Ala Thr His Ala Thr Tyr Thr Cys	
70 75 80	
cac atg gat gta ttc cac ttc atg gcc gac gac att ttc agt gtc aac	526
His Met Asp Val Phe His Phe Met Ala Asp Asp Ile Phe Ser Val Asn	
85 90 95	

atc Ile	aca Thr	gac Asp 100	cag Gln	tct Ser	ggc Gly	aac Asn	tac Tyr 105	tcc Ser	cag Gln	gag Glu	tgt Cys	ggc Gly 110	agc Ser	ttt Phe	ctc Leu	574
ctg Leu	gct Ala 115	gag Glu	agc Ser	atc Ile	aag Lys	ccg Pro 120	gct Ala	ccc Pro	cct Pro	ttc Phe	aac Asn 125	gtg Val	act Thr	gtg Val	acc Thr	622
ttc Phe 130	tca Ser	gga Gly	cag Gln	tat Tyr	aat Asn 135	atc Ile	tcc Ser	tgg Trp	cgc Arg	tca Ser 140	gat Asp	tac Tyr	gaa Glu	gac Asp	cct Pro 145	670
gcc Ala	ttc Phe	tac Tyr	atg Met	ctg Leu 150	aag Lys	ggc Gly	aag Lys	ctt Leu	cag Gln 155	tat Tyr	gag Glu	ctg Leu	cag Gln	tac Tyr 160	agg Arg	718
aac Asn	cgg Arg	gga Gly	gac Asp 165	ccc Pro	tgg Trp	gct Ala	gtg Val	agt Ser 170	ccg Pro	agg Arg	aga Arg	aag Lys	ctg Leu 175	atc Ile	tca Ser	766
gtg Val	gac Asp	tca Ser 180	aga Arg	agt Ser	gtc Val	tcc Ser	ctc Leu 185	ctc Leu	ccc Pro	ctg Leu	gag Glu	ttc Phe 190	cgc Arg	aaa Lys	gac Asp	814
tcg Ser	agc Ser 195	tat Tyr	gag Glu	ctg Leu	cag Gln	gtg Val 200	cgg Arg	gca Ala	ggg Gly	ccc Pro	atg Met 205	cct Pro	ggc Gly	tcc Ser	tcc Ser	862
tac Tyr 210	cag Gln	ggg Gly	acc Thr	tgg Trp	agt Ser 215	gaa Glu	tgg Trp	agt Ser	gac Asp	ccg Pro 220	gtc Val	atc Ile	ttt Phe	cag Gln	acc Thr 225	910
cag Gln	tca Ser	gag Glu	gag Glu	tta Leu 230	aag Lys	gaa Glu	ggc Gly	tgg Trp	aac Asn 235	cct Pro	cac His	ctg Leu	ctg Leu	ctt Leu 240	ctc Leu	958
ctc Leu	ctg Leu	ctt Leu	gtc Val 245	ata Ile	gtc Val	ttc Phe	att Ile	cct Pro 250	gcc Ala	ttc Phe	tgg Trp	agc Ser	ctg Leu 255	aag Lys	acc Thr	1006
cat His	cca Pro	ttg Leu 260	tgg Trp	agg Arg	cta Leu	tgg Trp	aag Lys 265	aag Lys	ata Ile	tgg Trp	gcc Ala	gtc Val 270	ccc Pro	agc Ser	cct Pro	1054
gag Glu	cgg Arg 275	ttc Phe	ttc Phe	atg Met	ccc Pro	ctg Leu 280	tac Tyr	aag Lys	ggc Gly	tgc Cys	agc Ser 285	gga Gly	gac Asp	ttc Phe	aag Lys	1102
aaa Lys 290	tgg Trp	gtg Val	ggt Gly	gca Ala	ccc Pro 295	ttc Phe	act Thr	ggc Gly	tcc Ser	agc Ser 300	ctg Leu	gag Glu	ctg Leu	gga Gly	ccc Pro 305	1150
tgg Trp	agc Ser	cca Pro	gag Glu	gtg Val 310	ccc Pro	tcc Ser	acc Thr	ctg Leu	gag Glu 315	gtg Val	tac Tyr	agc Ser	tgc Cys	cac His 320	cca Pro	1198
cca Pro	cgg Arg	agc Ser	ccg Pro 325	gcc Ala	aag Lys	agg Arg	ctg Leu	cag Gln 330	ctc Leu	acg Thr	gag Glu	cta Leu	caa Gln 335	gaa Glu	cca Pro	1246
gca Ala	gag Glu	ctg Leu	gtg Val	gag Glu	tct Ser	gac Asp	ggg Gly	gtg Val	ccc Pro	aag Lys	ccc Pro	agc Ser	ttc Phe	tgg Trp	ccg Pro	1294

340	345	350	
aca gcc cag aac tcg ggg ggc tca gct tac agt gag gag agg gat cgg Thr Ala Gln Asn Ser Gly Gly Ser Ala Tyr Ser Glu Glu Arg Asp Arg 355 360 365			1342
cca tac ggc ctg gtg tcc att gac aca gtg act gtg cta gat gca gag Pro Tyr Gly Leu Val Ser Ile Asp Thr Val Thr Val Leu Asp Ala Glu 370 375 380 385			1390
ggg cca tgc acc tgg ccc tgc agc tgt gag gat gac ggc tac cca gcc Gly Pro Cys Thr Trp Pro Cys Ser Cys Glu Asp Asp Gly Tyr Pro Ala 390 400			1438
ctg gac ctg gat gct ggc ctg gag ccc agc cca ggc cta gag gac cca Leu Asp Leu Asp Ala Gly Leu Glu Pro Ser Pro Gly Leu Glu Asp Pro 405 410 415			1486
ctc ttg gat gca ggg acc aca gtc ctg tcc tgt ggc tgt gtc tca gct Leu Leu Asp Ala Gly Thr Thr Val Leu Ser Cys Gly Cys Val Ser Ala 420 425 430			1534
ggc agc cct ggc cta gga ggg ccc ctg gga agc ctc ctg gac aga cta Gly Ser Pro Gly Leu Gly Gly Pro Leu Gly Ser Leu Leu Asp Arg Leu 435 440 445			1582
aag cca ccc ctt gca gat ggg gag gac tgg gct ggg gga ctg ccc tgg Lys Pro Pro Leu Ala Asp Gly Glu Asp Trp Ala Gly Gly Leu Pro Trp 450 455 460 465			1630
ggt ggc cgg tca cct gga ggg gtc tca gag agt gag gcg ggc tca ccc Gly Gly Arg Ser Pro Gly Gly Val Ser Glu Ser Glu Ala Gly Ser Pro 470 475 480			1678
ctg gcc ggc ctg gat atg gac acg ttt gac agt ggc ttt gtg ggc tct Leu Ala Gly Leu Asp Met Asp Thr Phe Asp Ser Gly Phe Val Gly Ser 485 490 495			1726
gac tgc agc agc cct gtg gag tgt gac ttc acc agc ccc ggg gac gaa Asp Cys Ser Ser Pro Val Glu Cys Asp Phe Thr Ser Pro Gly Asp Glu 500 505 510			1774
gga ccc ccc cgg agc tac ctc cgc cag tgg gtg gtc att cct ccg cca Gly Pro Pro Arg Ser Tyr Leu Arg Gln Trp Val Val Ile Pro Pro Pro 515 520 525			1822
ctt tcg agc cct gga ccc cag gcc agc taatgaggct gactggatgt Leu Ser Ser Pro Gly Pro Gln Ala Ser 530 535			1869
ccagagctgg ccaggccact gggccctgag ccagagacaa ggtcacctgg gctgtgatgt			1929
gaagacacct gcagcctttg gtctcctgga tgggcctttg agcctgatgt ttacagtgtc			1989
tgtgtgtgtg tgtgcatatg tgtgtgtgtg catatgcatg tgtgtgtgtg tgtgtgtcctt			2049
aggtgctgag tggcatgtcc acgtgtgtgt gtgattgcac gtgcctgtgg gcctgggata			2109
atgccccatgg tactccatgc attcacctgc cctgtgcatg tctggactca cggagctcac			2169
ccatgtgcac aagtgtgcac agtaaactgt tttgtgggtca acagatgaca acagccgtcc			2229
tccctcctag ggtcttgtgt tgcaagttgg tccacagcat ctccggggct ttgtgggatc			2289

```

agggcattgc ctgtgactga ggcggagccc agccctccag cgtctgcctc caggagctgc 2349
aagaagtcca tattgttctt tatcacctgc caacaggaag cgaaagggga tggagtgagc 2409
ccatggtgac ctcggaatg gcaatTTTTT gggcggtccc tggacgaagg tctgaatccc 2469
gactctgata ctttctggct gtgctacctg agccaagtcg cctcccctct ctgggctaga 2529
gtttccttat ccagacagtg gggaaggcat gacacacctg ggggaaattg gcgatgtcac 2589
ccgtgtacgg tacgcagccc agagcagacc ctcaataaac gtcagcttcc ttcaaaaaaa 2649
aaaaaaaaaa tctaga 2665

```

```

<210> 2
<211> 538
<212> PRT
<213> Homo sapiens

```

```

<400> 2

```

```

Met Pro Arg Gly Trp Ala Ala Pro Leu Leu Leu Leu Leu Leu Gln Gly
1          5          10          15

Gly Trp Gly Cys Pro Asp Leu Val Cys Tyr Thr Asp Tyr Leu Gln Thr
20          25          30

Val Ile Cys Ile Leu Glu Met Trp Asn Leu His Pro Ser Thr Leu Thr
35          40          45

Leu Thr Trp Gln Asp Gln Tyr Glu Glu Leu Lys Asp Glu Ala Thr Ser
50          55          60

Cys Ser Leu His Arg Ser Ala His Asn Ala Thr His Ala Thr Tyr Thr
65          70          75          80

Cys His Met Asp Val Phe His Phe Met Ala Asp Asp Ile Phe Ser Val
85          90          95

Asn Ile Thr Asp Gln Ser Gly Asn Tyr Ser Gln Glu Cys Gly Ser Phe
100          105          110

Leu Leu Ala Glu Ser Ile Lys Pro Ala Pro Pro Phe Asn Val Thr Val
115          120          125

Thr Phe Ser Gly Gln Tyr Asn Ile Ser Trp Arg Ser Asp Tyr Glu Asp
130          135          140

Pro Ala Phe Tyr Met Leu Lys Gly Lys Leu Gln Tyr Glu Leu Gln Tyr
145          150          155          160

```

Arg Asn Arg Gly Asp Pro Trp Ala Val Ser Pro Arg Arg Lys Leu Ile
 165 170 175
 Ser Val Asp Ser Arg Ser Val Ser Leu Leu Pro Leu Glu Phe Arg Lys
 180 185 190
 Asp Ser Ser Tyr Glu Leu Gln Val Arg Ala Gly Pro Met Pro Gly Ser
 195 200 205
 Ser Tyr Gln Gly Thr Trp Ser Glu Trp Ser Asp Pro Val Ile Phe Gln
 210 215 220
 Thr Gln Ser Glu Glu Leu Lys Glu Gly Trp Asn Pro His Leu Leu Leu
 225 230 235 240
 Leu Leu Leu Leu Val Ile Val Phe Ile Pro Ala Phe Trp Ser Leu Lys
 245 250 255
 Thr His Pro Leu Trp Arg Leu Trp Lys Lys Ile Trp Ala Val Pro Ser
 260 265 270
 Pro Glu Arg Phe Phe Met Pro Leu Tyr Lys Gly Cys Ser Gly Asp Phe
 275 280 285
 Lys Lys Trp Val Gly Ala Pro Phe Thr Gly Ser Ser Leu Glu Leu Gly
 290 295 300
 Pro Trp Ser Pro Glu Val Pro Ser Thr Leu Glu Val Tyr Ser Cys His
 305 310 315 320
 Pro Pro Arg Ser Pro Ala Lys Arg Leu Gln Leu Thr Glu Leu Gln Glu
 325 330 335
 Pro Ala Glu Leu Val Glu Ser Asp Gly Val Pro Lys Pro Ser Phe Trp
 340 345 350
 Pro Thr Ala Gln Asn Ser Gly Gly Ser Ala Tyr Ser Glu Glu Arg Asp
 355 360 365
 Arg Pro Tyr Gly Leu Val Ser Ile Asp Thr Val Thr Val Leu Asp Ala
 370 375 380
 Glu Gly Pro Cys Thr Trp Pro Cys Ser Cys Glu Asp Asp Gly Tyr Pro
 385 390 395 400
 Ala Leu Asp Leu Asp Ala Gly Leu Glu Pro Ser Pro Gly Leu Glu Asp
 405 410 415

Pro Leu Leu Asp Ala Gly Thr Thr Val Leu Ser Cys Gly Cys Val Ser
 420 425 430
 Ala Gly Ser Pro Gly Leu Gly Gly Pro Leu Gly Ser Leu Leu Asp Arg
 435 440 445
 Leu Lys Pro Pro Leu Ala Asp Gly Glu Asp Trp Ala Gly Gly Leu Pro
 450 455 460
 Trp Gly Gly Arg Ser Pro Gly Gly Val Ser Glu Ser Glu Ala Gly Ser
 465 470 475 480
 Pro Leu Ala Gly Leu Asp Met Asp Thr Phe Asp Ser Gly Phe Val Gly
 485 490 495
 Ser Asp Cys Ser Ser Pro Val Glu Cys Asp Phe Thr Ser Pro Gly Asp
 500 505 510
 Glu Gly Pro Pro Arg Ser Tyr Leu Arg Gln Trp Val Val Ile Pro Pro
 515 520 525
 Pro Leu Ser Ser Pro Gly Pro Gln Ala Ser
 530 535

<210> 3
 <211> 5
 <212> PRT
 <213> Unknown

<220>
 <223> Chemokine receptor conserved motif

<220>
 <221> misc_feature
 <222> (3)..(3)
 <223> Xaa can be any naturally occurring amino acid

<400> 3

Trp Ser Xaa Trp Ser
 1 5

<210> 4
 <211> 2628
 <212> DNA
 <213> Mus musculus

<220>
 <221> CDS
 <222> (407)..(1993)

<400> 4
gtcgcgcgcg cggtaccagc tgtctgcccc cttctcctgt ggtgtgcctc acggtcactt 60
gcttgtctga ccgcaagtct gcccatccct ggggcagcca actggcctca gcccggtgcc 120
caggcgtgcc ctgtctctgt ctggctgccc cagccctact gtcttcctct gtgtaggctc 180
tgcccagatg cccggctggg cctcagcctc aggactatct cagcagtgac tcccctgatt 240
ctggacttgc acctgactga actcctgccc acctcaaacc ttcacctccc accaccacca 300
ctccgagtcc cgctgtgact cccacgcccc ggagaccacc caagtgcctc agcctaaaga 360
atggccttct gagaaagacc ctgaaggagt aggtctggga cacagc atg ccc cgg 415
Met Pro Arg
1
ggc cca gtg gct gcc tta ctc ctg ctg att ctc cat gga gct tgg agc 463
Gly Pro Val Ala Ala Leu Leu Leu Ile Leu His Gly Ala Trp Ser
5 10 15
tgc ctg gac ctc act tgc tac act gac tac ctc tgg acc atc acc tgt 511
Cys Leu Asp Leu Thr Cys Tyr Thr Asp Tyr Leu Trp Thr Ile Thr Cys
20 25 30 35
gtc ctg gag aca cgg agc ccc aac ccc agc ata ctc agt ctc acc tgg 559
Val Leu Glu Thr Arg Ser Pro Asn Pro Ser Ile Leu Ser Leu Thr Trp
40 45 50
caa gat gaa tat gag gaa ctt cag gac caa gag acc ttc tgc agc cta 607
Gln Asp Glu Tyr Glu Glu Leu Gln Asp Gln Glu Thr Phe Cys Ser Leu
55 60 65
cac agg tct ggc cac aac acc aca cat ata tgg tac acg tgc cat atg 655
His Arg Ser Gly His Asn Thr Thr His Ile Trp Tyr Thr Cys His Met
70 75 80
cgc ttg tct caa ttc ctg tcc gat gaa gtt ttc att gtc aat gtg acg 703
Arg Leu Ser Gln Phe Leu Ser Asp Glu Val Phe Ile Val Asn Val Thr
85 90 95
gac cag tct ggc aac aac tcc caa gag tgt ggc agc ttt gtc ctg gct 751
Asp Gln Ser Gly Asn Asn Ser Gln Glu Cys Gly Ser Phe Val Leu Ala
100 105 110 115
gag agc atc aaa cca gct ccc ccc ttg aac gtg act gtg gcc ttc tca 799
Glu Ser Ile Lys Pro Ala Pro Pro Leu Asn Val Thr Val Ala Phe Ser
120 125 130
gga cgc tat gat atc tcc tgg gac tca gct tat gac gaa ccc tcc aac 847
Gly Arg Tyr Asp Ile Ser Trp Asp Ser Ala Tyr Asp Glu Pro 145
135 140 145
tac gtg ctg agg ggc aag cta caa tat gag ctg cag tat cgg aac ctc 895
Tyr Val Leu Arg Gly Lys Leu Gln Tyr Glu Leu Gln Tyr Arg Asn Leu
150 155 160
aga gac ccc tat gct gtg agg ccg gtg acc aag ctg atc tca gtg gac 943
Arg Asp Pro Tyr Ala Val Arg Pro Val Thr Lys Leu Ile Ser Val Asp
165 170 175
tca aga aac gtc tct ctt ctc cct gaa gag ttc cac aaa gat tct agc 991
Ser Arg Asn Val Ser Leu Leu Pro Glu Glu Phe His Lys Asp Ser Ser

180					185					190					195	
tac	cag	ctg	cag	gtg	cgg	gca	gcg	cct	cag	cca	ggc	act	tca	ttc	agg	1039
Tyr	Gln	Leu	Gln	Val	Arg	Ala	Ala	Pro	Gln	Pro	Gly	Thr	Ser	Phe	Arg	
				200					205					210		
ggg	acc	tgg	agt	gag	tgg	agt	gac	ccc	gtc	atc	ttt	cag	acc	cag	gct	1087
Gly	Thr	Trp	Ser	Glu	Trp	Ser	Asp	Pro	Val	Ile	Phe	Gln	Thr	Gln	Ala	
			215					220					225			
ggg	gag	ccc	gag	gca	ggc	tgg	gac	cct	cac	atg	ctg	ctg	ctc	ctg	gct	1135
Gly	Glu	Pro	Glu	Ala	Gly	Trp	Asp	Pro	His	Met	Leu	Leu	Leu	Leu	Ala	
		230					235					240				
gtc	ttg	atc	att	gtc	ctg	gtt	ttc	atg	ggc	ctg	aag	atc	cac	ctg	cct	1183
Val	Leu	Ile	Ile	Val	Leu	Val	Phe	Met	Gly	Leu	Lys	Ile	His	Leu	Pro	
	245					250					255					
tgg	agg	cta	tgg	aaa	aag	ata	tgg	gca	cca	gtg	ccc	acc	cct	gag	agt	1231
Trp	Arg	Leu	Trp	Lys	Lys	Ile	Trp	Ala	Pro	Val	Pro	Thr	Pro	Glu	Ser	
260					265					270					275	
ttc	ttc	cag	ccc	ctg	tac	agg	gag	cac	agc	ggg	aac	ttc	aag	aaa	tgg	1279
Phe	Phe	Gln	Pro	Leu	Tyr	Arg	Glu	His	Ser	Gly	Asn	Phe	Lys	Lys	Trp	
				280					285					290		
gtt	aat	acc	cct	ttc	acg	gcc	tcc	agc	ata	gag	ttg	gtg	cca	cag	agt	1327
Val	Asn	Thr	Pro	Phe	Thr	Ala	Ser	Ser	Ile	Glu	Leu	Val	Pro	Gln	Ser	
			295					300					305			
tcc	aca	aca	aca	tca	gcc	tta	cat	ctg	tca	ttg	tat	cca	gcc	aag	gag	1375
Ser	Thr	Thr	Thr	Ser	Ala	Leu	His	Leu	Ser	Leu	Tyr	Pro	Ala	Lys	Glu	
			310				315					320				
aag	aag	ttc	ccg	ggg	ctg	ccg	ggc	ctg	gaa	gag	caa	ctg	gag	tgt	gat	1423
Lys	Lys	Phe	Pro	Gly	Leu	Pro	Gly	Leu	Glu	Glu	Gln	Leu	Glu	Cys	Asp	
	325					330					335					
gga	atg	tct	gag	cct	ggc	cac	tgg	tgc	ata	atc	ccc	ttg	gca	gct	ggc	1471
Gly	Met	Ser	Glu	Pro	Gly	His	Trp	Cys	Ile	Ile	Pro	Leu	Ala	Ala	Gly	
340					345					350					355	
caa	gcg	gtc	tca	gcc	tac	agt	gag	gag	aga	gac	cgg	cca	tat	ggc	ctg	1519
Gln	Ala	Val	Ser	Ala	Tyr	Ser	Glu	Glu	Arg	Asp	Arg	Pro	Tyr	Gly	Leu	
				360					365					370		
gtg	tcc	att	gac	aca	gtg	act	gtg	gga	gat	gca	gag	ggc	ctg	tgt	gtc	1567
Val	Ser	Ile	Asp	Thr	Val	Thr	Val	Gly	Asp	Ala	Glu	Gly	Leu	Cys	Val	
			375					380					385			
tgg	ccc	tgt	agc	tgt	gag	gat	gat	ggc	tat	cca	gcc	atg	aac	ctg	gat	1615
Trp	Pro	Cys	Ser	Cys	Glu	Asp	Asp	Gly	Tyr	Pro	Ala	Met	Asn	Leu	Asp	
		390					395					400				
gct	ggc	cga	gag	tct	ggc	cct	aat	tca	gag	gat	ctg	ctc	ttg	gtc	aca	1663
Ala	Gly	Arg	Glu	Ser	Gly	Pro	Asn	Ser	Glu	Asp	Leu	Leu	Leu	Val	Thr	
	405					410					415					
gac	cct	gct	ttt	ctg	tct	tgc	ggc	tgt	gtc	tca	ggc	agt	ggc	ctc	agg	1711
Asp	Pro	Ala	Phe	Leu	Ser	Cys	Gly	Cys	Val	Ser	Gly	Ser	Gly	Leu	Arg	
420					425					430					435	
ctt	gga	ggc	tcc	cca	ggc	agc	cta	ctg	gac	agg	ttg	agg	ctg	tca	ttt	1759

Leu	Gly	Gly	Ser	Pro	Gly	Ser	Leu	Leu	Asp	Arg	Leu	Arg	Leu	Ser	Phe	
				440					445					450		
gca	aag	gaa	ggg	gac	tgg	aca	gca	gac	cca	acc	tgg	aga	act	ggg	tcc	1807
Ala	Lys	Glu	Gly	Asp	Trp	Thr	Ala	Asp	Pro	Thr	Trp	Arg	Thr	Gly	Ser	
			455					460					465			
cca	gga	ggg	ggc	tct	gag	agt	gaa	gca	ggg	tcc	ccc	cct	ggg	ctg	gac	1855
Pro	Gly	Gly	Gly	Ser	Glu	Ser	Glu	Ala	Gly	Ser	Pro	Pro	Gly	Leu	Asp	
		470					475				480					
atg	gac	aca	ttt	gac	agt	ggc	ttt	gca	ggg	tca	gac	tgt	ggc	agc	ccc	1903
Met	Asp	Thr	Phe	Asp	Ser	Gly	Phe	Ala	Gly	Ser	Asp	Cys	Gly	Ser	Pro	
	485					490					495					
gtg	gag	act	gat	gaa	gga	ccc	cct	cga	agc	tat	ctc	cgc	cag	tgg	gtg	1951
Val	Glu	Thr	Asp	Glu	Gly	Pro	Pro	Arg	Ser	Tyr	Leu	Arg	Gln	Trp	Val	
500					505					510					515	
gtc	agg	acc	cct	cca	cct	gtg	gac	agt	gga	gcc	cag	agc	agc			1993
Val	Arg	Thr	Pro	Pro	Pro	Val	Asp	Ser	Gly	Ala	Gln	Ser	Ser			
				520					525							
tagcatataa	taaccagcta	tagtgagaag	aggcctctga	gcctggcatt	tacagtgtga											2053
acatgtaggg	gtgtgtgtgt	gtgtgtgtgt	gtgtgtgtgt	gtgtgtgtgt	gtgtgtgtgt											2113
gtgtgtgtgt	cttgggttgt	gtgttagcac	atccatgttg	ggatttggtc	tggtgctatg											2173
tattgtaatg	ctaaattctc	tacccaaagt	tctaggccta	cgagtgaatt	ctcatgttta											2233
caaacttgct	gtgtaaacct	tggtccttaa	tttaatacca	ttgggttaa	aaaattggct											2293
gcaaccaatt	actggaggga	ttagaggtag	ggggcctttg	agttacctgt	ttggagatgg											2353
agaaggagag	aggagagacc	aagaggagaa	ggaggaagga	gaggagagga	gaggagagga											2413
gaggagagga	gaggagagga	gaggagagga	gaggagaggc	tgccgtgagg	ggagaggggac											2473
catgagcctg	tggccaggag	aaacagcaag	tatctggggg	acactgggtga	ggaggtggcc											2533
aggccagcag	ttagaagagt	agattagggg	tgacctccag	tatttggtcaa	agccaattaa											2593
aataacaaaa	aaaaaaaaaa	agcggccgct	ctaga													2628

<210> 5
 <211> 529
 <212> PRT
 <213> Mus musculus

<400> 5

Met Pro Arg Gly Pro Val Ala Ala Leu Leu Leu Ile Leu His Gly
1 5 10 15

Ala Trp Ser Cys Leu Asp Leu Thr Cys Tyr Thr Asp Tyr Leu Trp Thr
20 25 30

Ile Thr Cys Val Leu Glu Thr Arg Ser Pro Asn Pro Ser Ile Leu Ser
35 40 45

Leu Thr Trp Gln Asp Glu Tyr Glu Glu Leu Gln Asp Gln Glu Thr Phe
 50 55 60
 Cys Ser Leu His Arg Ser Gly His Asn Thr Thr His Ile Trp Tyr Thr
 65 70 75 80
 Cys His Met Arg Leu Ser Gln Phe Leu Ser Asp Glu Val Phe Ile Val
 85 90 95
 Asn Val Thr Asp Gln Ser Gly Asn Asn Ser Gln Glu Cys Gly Ser Phe
 100 105 110
 Val Leu Ala Glu Ser Ile Lys Pro Ala Pro Pro Leu Asn Val Thr Val
 115 120 125
 Ala Phe Ser Gly Arg Tyr Asp Ile Ser Trp Asp Ser Ala Tyr Asp Glu
 130 135 140
 Pro Ser Asn Tyr Val Leu Arg Gly Lys Leu Gln Tyr Glu Leu Gln Tyr
 145 150 155 160
 Arg Asn Leu Arg Asp Pro Tyr Ala Val Arg Pro Val Thr Lys Leu Ile
 165 170 175
 Ser Val Asp Ser Arg Asn Val Ser Leu Leu Pro Glu Glu Phe His Lys
 180 185 190
 Asp Ser Ser Tyr Gln Leu Gln Val Arg Ala Ala Pro Gln Pro Gly Thr
 195 200 205
 Ser Phe Arg Gly Thr Trp Ser Glu Trp Ser Asp Pro Val Ile Phe Gln
 210 215 220
 Thr Gln Ala Gly Glu Pro Glu Ala Gly Trp Asp Pro His Met Leu Leu
 225 230 235 240
 Leu Leu Ala Val Leu Ile Ile Val Leu Val Phe Met Gly Leu Lys Ile
 245 250 255
 His Leu Pro Trp Arg Leu Trp Lys Lys Ile Trp Ala Pro Val Pro Thr
 260 265 270
 Pro Glu Ser Phe Phe Gln Pro Leu Tyr Arg Glu His Ser Gly Asn Phe
 275 280 285
 Lys Lys Trp Val Asn Thr Pro Phe Thr Ala Ser Ser Ile Glu Leu Val

Ser

<210> 6
 <211> 224
 <212> PRT
 <213> Unknown

<220>
 <223> Immunoglobulin Fc Fragment

<400> 6

His Thr Cys Pro Pro Cys Pro Ala Pro Glu Ala Leu Gly Ala Pro Ser
 1 5 10 15

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg
 20 25 30

Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro
 35 40 45

Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala
 50 55 60

Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val
 65 70 75 80

Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr
 85 90 95

Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Val Pro Ile Glu Lys Thr
 100 105 110

Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu
 115 120 125

Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys
 130 135 140

Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser
 145 150 155 160

Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp
 165 170 175

Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser
 180 185 190

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala
 195 200 205

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys

<210> 8
 <211> 162
 <212> PRT
 <213> Homo sapiens

<400> 8

Met Arg Ser Ser Pro Gly Asn Met Glu Arg Ile Val Ile Cys Leu Met
 1 5 10 15

Val Ile Phe Leu Gly Thr Leu Val His Lys Ser Ser Ser Gln Gly Gln
 20 25 30

Asp Arg His Met Ile Arg Met Arg Gln Leu Ile Asp Ile Val Asp Gln
 35 40 45

Leu Lys Asn Tyr Val Asn Asp Leu Val Pro Glu Phe Leu Pro Ala Pro
 50 55 60

Glu Asp Val Glu Thr Asn Cys Glu Trp Ser Ala Phe Ser Cys Phe Gln
 65 70 75 80

Lys Ala Gln Leu Lys Ser Ala Asn Thr Gly Asn Asn Glu Arg Ile Ile
 85 90 95

Asn Val Ser Ile Lys Lys Leu Lys Arg Lys Pro Pro Ser Thr Asn Ala
 100 105 110

Gly Arg Arg Gln Lys His Arg Leu Thr Cys Pro Ser Cys Asp Ser Tyr
 115 120 125

Glu Lys Lys Pro Pro Lys Glu Phe Leu Glu Arg Phe Lys Ser Leu Leu
 130 135 140

Gln Lys Met Ile His Gln His Leu Ser Ser Arg Thr His Gly Ser Glu
 145 150 155 160

Asp Ser

<210> 9
 <211> 16
 <212> PRT
 <213> Apis mellifera

<400> 9

Met Pro Leu Leu Leu Leu Leu Leu Leu Pro Ser Pro Leu His Pro
 1 5 10 15

<210> 10

<211> 786
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)..(780)

<400> 10
 atg aaa ttc tta gtc aac gtt gcc ctt gtt ttt atg gtc gtg tac att 48
 Met Lys Phe Leu Val Asn Val Ala Leu Val Phe Met Val Val Tyr Ile
 1 5 10 15
 tct tac atc tat gcc ggc agc gga cac cac cat cat cac cac ggt agc 96
 Ser Tyr Ile Tyr Ala Gly Ser Gly His His His His His His Gly Ser
 20 25 30
 ggc gac tat aaa gac gat gac gat aag ggt tcc gga tgc ccc gac ctc 144
 Gly Asp Tyr Lys Asp Asp Asp Asp Lys Gly Ser Gly Cys Pro Asp Leu
 35 40 45
 gtc tgc tac acc gat tac ctc cag acg gtc atc tgc atc ctg gaa atg 192
 Val Cys Tyr Thr Asp Tyr Leu Gln Thr Val Ile Cys Ile Leu Glu Met
 50 55 60
 tgg aac ctc cac ccc agc acg ctc acc ctt acc tgg caa gac cag tat 240
 Trp Asn Leu His Pro Ser Thr Leu Thr Leu Thr Trp Gln Asp Gln Tyr
 65 70 75
 gaa gag ctg aag gac gag gcc acc tcc tgc agc ctc cac agg tcg gcc 288
 Glu Glu Leu Lys Asp Glu Ala Thr Ser Cys Ser Leu His Arg Ser Ala
 85 90 95
 cac aat gcc acg cat gcc acc tac acc tgc cac atg gat gta ttc cac 336
 His Asn Ala Thr His Ala Thr Tyr Thr Cys His Met Asp Val Phe His
 100 105 110
 ttc atg gcc gac gac att ttc agt gtc aac atc aca gac cag tct ggc 384
 Phe Met Ala Asp Asp Ile Phe Ser Val Asn Ile Thr Asp Gln Ser Gly
 115 120 125
 aac tac tcc cag gag tgt ggc agc ttt ctc ctg gct gag agc atc aag 432
 Asn Tyr Ser Gln Glu Cys Gly Ser Phe Leu Leu Ala Glu Ser Ile Lys
 130 135 140
 ccg gct ccc cct ttc aac gtg act gtg acc ttc tca gga cag tat aat 480
 Pro Ala Pro Pro Phe Asn Val Thr Val Thr Phe Ser Gly Gln Tyr Asn
 145 150 155 160
 atc tcc tgg cgc tca gat tac gaa gac cct gcc ttc tac atg ctg aag 528
 Ile Ser Trp Arg Ser Asp Tyr Glu Asp Pro Ala Phe Tyr Met Leu Lys
 165 170 175
 ggc aag ctt cag tat gag ctg cag tac agg aac cgg gga gac ccc tgg 576
 Gly Lys Leu Gln Tyr Glu Leu Gln Tyr Arg Asn Arg Gly Asp Pro Trp
 180 185 190
 gct gtg agt ccg agg aga aag ctg atc tca gtg gac tca aga agt gtc 624
 Ala Val Ser Pro Arg Arg Lys Leu Ile Ser Val Asp Arg Ser Val
 195 200 205
 tcc ctc ctc ccc ctg gag ttc cgc aaa gac tcg agc tat gag ctg cag 672

Ser	Leu	Leu	Pro	Leu	Glu	Phe	Arg	Lys	Asp	Ser	Ser	Tyr	Glu	Leu	Gln		
210						215				220							
gtg	cgg	gca	ggg	ccc	atg	cct	ggc	tcc	tcc	tac	cag	ggg	acc	tgg	agt	720	
Val	Arg	Ala	Gly	Pro	Met	Pro	Gly	Ser	Ser	Tyr	Gln	Gly	Thr	Trp	Ser		
225					230					235					240		
gaa	tgg	agt	gac	ccg	gtc	atc	ttt	cag	acc	cag	tca	gag	gag	tta	aag	768	
Glu	Trp	Ser	Asp	Pro	Val	Ile	Phe	Gln	Thr	Gln	Ser	Glu	Glu	Leu	Lys		
				245					250					255			
gaa	ggc	tgg	aac	taatga												786	
Glu	Gly	Trp	Asn														
			260														

<210> 11
 <211> 260
 <212> PRT
 <213> Homo sapiens
 <400> 11

Met	Lys	Phe	Leu	Val	Asn	Val	Ala	Leu	Val	Phe	Met	Val	Val	Tyr	Ile		
1				5					10					15			
Ser	Tyr	Ile	Tyr	Ala	Gly	Ser	Gly	His	His	His	His	His	His	Gly	Ser		
			20					25						30			
Gly	Asp	Tyr	Lys	Asp	Asp	Asp	Asp	Lys	Gly	Ser	Gly	Cys	Pro	Asp	Leu		
		35					40					45					
Val	Cys	Tyr	Thr	Asp	Tyr	Leu	Gln	Thr	Val	Ile	Cys	Ile	Leu	Glu	Met		
	50					55					60						
Trp	Asn	Leu	His	Pro	Ser	Thr	Leu	Thr	Leu	Thr	Trp	Gln	Asp	Gln	Tyr		
65					70					75					80		
Glu	Glu	Leu	Lys	Asp	Glu	Ala	Thr	Ser	Cys	Ser	Leu	His	Arg	Ser	Ala		
				85					90					95			
His	Asn	Ala	Thr	His	Ala	Thr	Tyr	Thr	Cys	His	Met	Asp	Val	Phe	His		
			100					105					110				
Phe	Met	Ala	Asp	Asp	Ile	Phe	Ser	Val	Asn	Ile	Thr	Asp	Gln	Ser	Gly		
		115					120					125					
Asn	Tyr	Ser	Gln	Glu	Cys	Gly	Ser	Phe	Leu	Leu	Ala	Glu	Ser	Ile	Lys		
	130					135					140						
Pro	Ala	Pro	Pro	Phe	Asn	Val	Thr	Val	Thr	Phe	Ser	Gly	Gln	Tyr	Asn		
145					150					155					160		

Ile Ser Trp Arg Ser₁₆₅ Asp Tyr Glu Asp₁₇₀ Pro Ala Phe Tyr Met Leu₁₇₅ Lys
 Gly Lys Leu Gln₁₈₀ Tyr Glu Leu Gln₁₈₅ Tyr Arg Asn Arg Gly Asp₁₉₀ Pro Trp
 Ala Val₁₉₅ Ser Pro Arg Arg Lys Leu₂₀₀ Ile Ser Val Asp₂₀₅ Ser Arg Ser Val
 Ser Leu₂₁₀ Leu Pro Leu Glu Phe₂₁₅ Arg Lys Asp Ser Ser Tyr Glu Leu Gln
 Val Arg Ala Gly Pro Met₂₃₀ Pro Gly Ser Ser Tyr₂₃₅ Gln Gly Thr Trp Ser₂₄₀
 Glu Trp Ser Asp Pro₂₄₅ Val Ile Phe Gln Thr₂₅₀ Gln Ser Glu Glu Leu₂₅₅ Lys
 Glu Gly Trp Asn₂₆₀

<210> 12
 <211> 1472
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (15)..(1415)

<400> 12
 gcggccgcac cacc atg ccg cgt ggc tgg gcc gcc ccc ttg ctc ctg ctg 50
 Met Pro Arg Gly Trp Ala Ala Pro Leu Leu Leu Leu
 1 5 10
 ctg ctc cag gga ggc tgg ggc tgc ccc gac ctc gtc tgc tac acc gat 98
 Leu Leu Gln Gly Gly Trp Gly Cys Pro Asp Leu Val Cys Tyr Thr Asp
 15 20 25
 tac ctc cag acg gtc atc tgc atc ctg gaa atg tgg aac ctc cac ccc 146
 Tyr Leu Gln Thr Val Ile Cys Ile Leu Glu Met Trp Asn Leu His Pro
 30 35 40
 agc acg ctc acc ctt acc tgg caa gac cag tat gaa gag ctg aag gac 194
 Ser Thr Leu Thr Leu Thr Trp Gln Asp Gln Tyr Glu Glu Leu Lys Asp
 45 50 55 60
 gag gcc acc tcc tgc agc ctc cac agg tcg gcc cac aat gcc acg cat 242
 Glu Ala Thr Ser Cys Ser Leu His Arg Ser Ala His Asn Ala Thr His
 65 70
 gcc acc tac acc tgc cac atg gat gta ttc cac ttc atg gcc gac gac 290
 Ala Thr Tyr Thr Cys His Met Asp Val Phe His Phe Met Ala Asp Asp
 80 85 90

att Ile	ttc Phe	agt Ser 95	gtc Val	aac Asn	atc Ile	aca Thr	gac Asp 100	cag Gln	tct Ser	ggc Gly	aac Asn	tac Tyr 105	tcc Ser	cag Gln	gag Glu	338
tgt Cys	ggc Gly 110	agc Ser	ttt Phe	ctc Leu	ctg Leu	gct Ala 115	gag Glu	agc Ser	atc Ile	aag Lys	ccg Pro 120	gct Ala	ccc Pro	cct Pro	ttc Phe	386
aac Asn 125	gtg Val	act Thr	gtg Val	acc Thr	ttc Phe 130	tca Ser	gga Gly	cag Gln	tat Tyr	aat Asn 135	atc Ile	tcc Ser	tgg Trp	cgc Arg	tca Ser 140	434
gat Asp	tac Tyr	gaa Glu	gac Asp	cct Pro 145	gcc Ala	ttc Phe	tac Tyr	atg Met	ctg Leu 150	aag Lys	ggc Gly	aag Lys	ctt Leu	cag Gln 155	tat Tyr	482
gag Glu	ctg Leu	cag Gln	tac Tyr 160	agg Arg	aac Asn	cgg Arg	gga Gly	gac Asp 165	ccc Pro	tgg Trp	gct Ala	gtg Val	agt Ser 170	ccg Pro	agg Arg	530
aga Arg	aag Lys	ctg Leu 175	atc Ile	tca Ser	gtg Val	gac Asp	tca Ser 180	aga Arg	agt Ser	gtc Val	tcc Ser	ctc Leu 185	ctc Leu	ccc Pro	ctg Leu	578
gag Glu 190	ttc Phe	cgc Arg	aaa Lys	gac Asp	tcg Ser	agc Ser 195	tat Tyr	gag Glu	ctg Leu	cag Gln	gtg Val 200	cgg Arg	gca Ala	ggg Gly	ccc Pro	626
atg Met 205	cct Pro	ggc Gly	tcc Ser	tcc Ser	tac Tyr 210	cag Gln	ggg Gly	acc Thr	tgg Trp	agt Ser 215	gaa Glu	tgg Trp	agt Ser	gac Asp	ccg Pro 220	674
gtc Val	atc Ile	ttt Phe	cag Gln	acc Thr 225	cag Gln	tca Ser	gag Glu	gag Glu	tta Leu 230	aag Lys	gaa Glu	ggc Gly	tgg Trp	aac Asn 235	ggc Gly	722
tcc Ser	ggc Gly	tct Ser	aga Arg 240	gac Asp	aaa Lys	act Thr	cac His	aca Thr 245	tgc Cys	cca Pro	ccg Pro	tgc Cys	cca Pro 250	gca Ala	cct Pro	770
gaa Glu	ctc Leu	ctg Leu 255	ggg Gly	gga Gly	ccg Pro	tca Ser	gtc Val 260	ttc Phe	ctc Leu	ttc Phe	ccc Pro	cca Pro 265	aaa Lys	ccc Pro	aag Lys	818
gac Asp 270	acc Thr	ctc Leu	atg Met	atc Ile	tcc Ser	cgg Arg 275	acc Thr	cct Pro	gag Glu	gtc Val	aca Thr 280	tgc Cys	gtg Val	gtg Val	gtg Val	866
gac Asp 285	gtg Val	agc Ser	cac His	gaa Glu	gac Asp 290	cct Pro	gag Glu	gtc Val	aag Lys	ttc Phe 295	aac Asn	tgg Trp	tac Tyr	gtg Val	gac Asp 300	914
ggc Gly	gtg Val	gag Glu	gtg Val	cat His 305	aat Asn	gcc Ala	aag Lys	aca Thr 310	aag Lys	ccg Pro	cgg Arg	gag Glu	gag Glu	cag Gln 315	tac Tyr	962
aac Asn	agc Ser	acg Thr	tac Tyr 320	cgt Arg	gtg Val	gtc Val	agc Ser	gtc Val 325	ctc Leu	acc Thr	gtc Val	ctg Leu	cac His 330	cag Gln	gac Asp	1010
tgg Trp	ctg Leu	aat Asn 335	ggc Gly	aag Lys	gag Glu	tac Tyr	aag Lys 340	tgc Cys	aag Lys	gtc Val	tcc Ser	aac Asn 345	aaa Lys	gcc Ala	ctc Leu	1058

cca gtc ccc atc gag aaa acc atc tcc aaa gcc aaa ggg cag ccc cga 1106
 Pro Val Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg
 350 355 360

gaa cca cag gtg tac acc ctg ccc cca tcc cgg gag gag atg acc aag 1154
 Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys
 365 370 375 380

aac cag gtc agc ctg acc tgc ctg gtc aaa ggc ttc tat ccc agc gac 1202
 Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp
 385 390 395

atc gcc gtg gag tgg gag agc aat ggg cag ccg gag aac aac tac aag 1250
 Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys
 400 405 410

acc acg cct ccc gtg ctg gac tcc gac ggc tcc ttc ttc ctc tat agc 1298
 Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser
 415 420 425

aag ctc acc gtg gac aag agc agg tgg cag cag ggg aac gtc ttc tca 1346
 Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser
 430 435 440

tgc tcc gtg atg cat gag gct ctg cac aac cac tac acg cag aag agc 1394
 Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser
 445 450 455 460

ctc tcc ctg tcc ccg ggt aaa tgagtgaatt cacacgcaga agagcctctc 1445
 Leu Ser Leu Ser Pro Gly Lys

465

cctgtcccccg ggtaaagtgaattc 1472

<210> 13
 <211> 467
 <212> PRT
 <213> Homo sapiens

<400> 13

Met Pro Arg Gly Trp Ala Ala Pro Leu Leu Leu Leu Leu Gln Gly
 1 5 10 15

Gly Trp Gly Cys Pro Asp Leu Val Cys Tyr Thr Asp Tyr Leu Gln Thr
 20 25 30

Val Ile Cys Ile Leu Glu Met Trp Asn Leu His Pro Ser Thr Leu Thr
 35 40 45

Leu Thr Trp Gln Asp Gln Tyr Glu Glu Leu Lys Asp Glu Ala Thr Ser
 50 55 60

Cys Ser Leu His Arg Ser Ala His Asn Ala Thr His Ala Thr Tyr Thr
 65 70 75 80

Cys His Met Asp Val Phe His Phe Met Ala Asp Asp Ile Phe Ser Val
 85 90 95

Asn Ile Thr Asp Gln Ser Gly Asn Tyr Ser Gln Glu Cys Gly Ser Phe
 100 105 110

Leu Leu Ala Glu Ser Ile Lys Pro Ala Pro Pro Phe Asn Val Thr Val
 115 120 125

Thr Phe Ser Gly Gln Tyr Asn Ile Ser Trp Arg Ser Asp Tyr Glu Asp
 130 135 140

Pro Ala Phe Tyr Met Leu Lys Gly Lys Leu Gln Tyr Glu Leu Gln Tyr
 145 150 155 160

Arg Asn Arg Gly Asp Pro Trp Ala Val Ser Pro Arg Arg Lys Leu Ile
 165 170 175

Ser Val Asp Ser Arg Ser Val Ser Leu Leu Pro Leu Glu Phe Arg Lys
 180 185 190

Asp Ser Ser Tyr Glu Leu Gln Val Arg Ala Gly Pro Met Pro Gly Ser
 195 200 205

Ser Tyr Gln Gly Thr Trp Ser Glu Trp Ser Asp Pro Val Ile Phe Gln
 210 215 220

Thr Gln Ser Glu Glu Leu Lys Glu Gly Trp Asn Gly Ser Gly Ser Arg
 225 230 235 240

Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly
 245 250 255

Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met
 260 265 270

Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His
 275 280 285

Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val
 290 295 300

His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr
 305 310 315 320

Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly
 325 330 335

Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Val Pro Ile
 340 345 350
 Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val
 355 360 365
 Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser
 370 375 380
 Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu
 385 390 395 400
 Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro
 405 410 415
 Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val
 420 425 430
 Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met
 435 440 445
 His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser
 450 455 460
 Pro Gly Lys
 465

<210> 14
 <211> 1499
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (15)..(1493)

<400> 14
 gcggccgcac cacc atg ccg cgt ggc tgg gcc gcc ccc ttg ctc ctg ctg 50
 Met Pro Arg Gly Trp Ala Ala Pro Leu Leu Leu
 1 5 10
 ctg ctc cag gga ggc tgg ggc tgc ccc gac ctc gtc tgc tac acc gat 98
 Leu Leu Gln Gly Gly Trp Gly Cys Pro Asp Leu Val Cys Tyr Thr Asp
 15 20 25
 tac ctc cag acg gtc atc tgc atc ctg gaa atg tgg aac ctc cac ccc 146
 Tyr Leu Gln Thr Val Ile Cys Ile Leu Glu Met Trp Asn Leu His Pro
 30 35 40
 agc acg ctc acc ctt acc tgg caa gac cag tat gaa gag ctg aag gac 194
 Ser Thr Leu Thr Leu Thr Trp Gln Asp Gln Tyr Glu Glu Leu Lys Asp
 45 50 55 60

gag Glu	gcc Ala	acc Thr	tcc Ser	tgc Cys 65	agc Ser	ctc Leu	cac His	agg Arg	tcg Ser 70	gcc Ala	cac His	aat Asn	gcc Ala	acg Thr 75	cat His	242
gcc Ala	acc Thr	tac Tyr	acc Thr 80	tgc Cys	cac His	atg Met	gat Asp	gta Val 85	ttc Phe	cac His	ttc Phe	atg Met	gcc Ala 90	gac Asp	gac Asp	290
att Ile	ttc Phe	agt Ser 95	gtc Val	aac Asn	atc Ile	aca Thr	gac Asp 100	cag Gln	tct Ser	ggc Gly	aac Asn	tac Tyr 105	tcc Ser	cag Gln	gag Glu	338
tgt Cys	ggc Gly 110	agc Ser	ttt Phe	ctc Leu	ctg Leu	gct Ala 115	gag Glu	agc Ser	atc Ile	aag Lys	ccg Pro 120	gct Ala	ccc Pro	cct Pro	ttc Phe	386
aac Asn 125	gtg Val	act Thr	gtg Val	acc Thr	ttc Phe 130	tca Ser	gga Gly	cag Gln	tat Tyr	aat Asn 135	atc Ile	tcc Ser	tgg Trp	cgc Arg	tca Ser 140	434
gat Asp	tac Tyr	gaa Glu	gac Asp	cct Pro 145	gcc Ala	ttc Phe	tac Tyr	atg Met	ctg Leu 150	aag Lys	ggc Gly	aag Lys	ctt Leu	cag Gln 155	tat Tyr	482
gag Glu	ctg Leu	cag Gln	tac Tyr 160	agg Arg	aac Asn	cgg Arg	gga Gly	gac Asp 165	ccc Pro	tgg Trp	gct Ala	gtg Val 170	agt Ser 170	ccg Pro	agg Arg	530
aga Arg	aag Lys	ctg Leu 175	atc Ile	tca Ser	gtg Val	gac Asp	tca Ser 180	aga Arg	agt Ser	gtc Val	tcc Ser	ctc Leu 185	ctc Leu	ccc Pro	ctg Leu	578
gag Glu 190	ttc Phe	cgc Arg	aaa Lys	gac Asp	tcg Ser	agc Ser 195	tat Tyr	gag Glu	ctg Leu	cag Gln	gtg Val 200	cgg Arg	gca Ala	ggg Gly	ccc Pro	626
atg Met 205	cct Pro	ggc Gly	tcc Ser	tcc Ser	tac Tyr 210	cag Gln	ggg Gly	acc Thr	tgg Trp	agt Ser 215	gaa Glu	tgg Trp	agt Ser	gac Asp	ccg Pro 220	674
gtc Val	atc Ile	ttt Phe	cag Gln 225	acc Thr	cag Gln	tca Ser	gag Glu	gag Glu	tta Leu 230	aag Lys	gaa Glu	ggc Gly	tgg Trp	aac Asn 235	ggc Gly	722
tcc Ser	ggc Gly	tct Ser	aga Arg 240	gac Asp	aaa Lys	act Thr	cac His	aca Thr 245	tgc Cys	cca Pro	ccg Pro	tgc Cys	cca Pro 250	gca Ala	cct Pro	770
gaa Glu	ctc Leu	ctg Leu 255	ggg Gly	gga Gly	ccg Pro	tca Ser	gtc Val 260	ttc Phe	ctc Leu	ttc Phe	ccc Pro	cca Pro 265	aaa Lys	ccc Pro	aag Lys	818
gac Asp	acc Thr 270	ctc Leu	atg Met	atc Ile	tcc Ser	cgg Arg 275	acc Thr	cct Pro	gag Glu	gtc Val	aca Thr 280	tgc Cys	gtg Val	gtg Val	gtg Val	866
gac Asp 285	gtg Val	agc Ser	cac His	gaa Glu	gac Asp 290	cct Pro	gag Glu	gtc Val	aag Lys	ttc Phe 295	aac Asn	tgg Trp	tac Tyr	gtg Val	gac Asp 300	914
ggc Gly	gtg Val	gag Glu	gtg Val	cat His	aat Asn	gcc Ala	aag Lys	aca Thr	aag Lys	ccg Pro	cgg Arg	gag Glu	gag Glu	cag Gln	tac Tyr	962

				305				310				315							
aac	agc	acg	tac	cgt	gtg	gtc	agc	gtc	ctc	acc	gtc	ctg	cac	cag	gac	1010			
Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp				
				320					325					330					
tgg	ctg	aat	ggc	aag	gag	tac	aag	tgc	aag	gtc	tcc	aac	aaa	gcc	ctc	1058			
Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu				
				335					340					345					
cca	gtc	ccc	atc	gag	aaa	acc	atc	tcc	aaa	gcc	aaa	ggg	cag	ccc	cga	1106			
Pro	Val	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg				
				350					355					360					
gaa	cca	cag	gtg	tac	acc	ctg	ccc	cca	tcc	cgg	gag	gag	atg	acc	aag	1154			
Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys				
				365					370					375					
aac	cag	gtc	agc	ctg	acc	tgc	ctg	gtc	aaa	ggc	ttc	tat	ccc	agc	gac	1202			
Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp				
				385					390					395					
atc	gcc	gtg	gag	tgg	gag	agc	aat	ggg	cag	ccg	gag	aac	aac	tac	aag	1250			
Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys				
				400					405					410					
acc	acg	cct	ccc	gtg	ctg	gac	tcc	gac	ggc	tcc	ttc	ttc	ctc	tat	agc	1298			
Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser				
				415					420					425					
aag	ctc	acc	gtg	gac	aag	agc	agg	tgg	cag	cag	ggg	aac	gtc	ttc	tca	1346			
Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser				
				430					435					440					
tgc	tcc	gtg	atg	cat	gag	gct	ctg	cac	aac	cac	tac	acg	cag	aag	agc	1394			
Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser				
				445					450					455					
ctc	tcc	ctg	tcc	ccg	ggc	aaa	tca	gga	atg	gca	tca	atg	aca	gga	ggc	1442			
Leu	Ser	Leu	Ser	Pro	Gly	Lys	Ser	Gly	Met	Ala	Ser	Met	Thr	Gly	Gly				
				465					470					475					
caa	caa	atg	ggc	tct	gga	tct	cat	cat	cat	cat	cat	cat	tct	gga	ggc	1490			
Gln	Gln	Met	Gly	Ser	Gly	Ser	His	His	His	His	His	His	Ser	Gly	Gly				
				480					485					490					
tga	gaattc															1499			
<210>	15																		
<211>	492																		
<212>	PRT																		
<213>	Homo sapiens																		
<400>	15																		
Met	Pro	Arg	Gly	Trp	Ala	Ala	Pro	Leu	Leu	Leu	Leu	Leu	Leu	Gln	Gly				
1				5				10						15					
Gly	Trp	Gly	Cys	Pro	Asp	Leu	Val	Cys	Tyr	Thr	Asp	Tyr	Leu	Gln	Thr				
				20					25					30					

Val Ile Cys₃₅ Ile Leu Glu Met Trp₄₀ Asn Leu His Pro Ser₄₅ Thr Leu Thr
 Leu Thr₅₀ Trp Gln Asp Gln Tyr₅₅ Glu Glu Leu Lys Asp₆₀ Glu Ala Thr Ser
 Cys₆₅ Ser Leu His Arg Ser₇₀ Ala His Asn Ala Thr₇₅ His Ala Thr Tyr Thr₈₀
 Cys His Met Asp Val₈₅ Phe His Phe Met Ala₉₀ Asp Asp Ile Phe Ser₉₅ Val
 Asn Ile Thr Asp₁₀₀ Gln Ser Gly Asn Tyr₁₀₅ Ser Gln Glu Cys Gly₁₁₀ Ser Phe
 Leu Leu Ala₁₁₅ Glu Ser Ile Lys Pro₁₂₀ Ala Pro Pro Phe Asn₁₂₅ Val Thr Val
 Thr Phe₁₃₀ Ser Gly Gln Tyr Asn₁₃₅ Ile Ser Trp Arg Ser₁₄₀ Asp Tyr Glu Asp
 Pro Ala Phe Tyr Met Leu₁₅₀ Lys Gly Lys Leu Gln₁₅₅ Tyr Glu Leu Gln Tyr₁₆₀
 Arg Asn Arg Gly Asp₁₆₅ Pro Trp Ala Val Ser₁₇₀ Pro Arg Arg Lys Leu₁₇₅ Ile
 Ser Val Asp Ser₁₈₀ Arg Ser Val Ser Leu₁₈₅ Leu Pro Leu Glu Phe₁₉₀ Arg Lys
 Asp Ser Ser₁₉₅ Tyr Glu Leu Gln Val₂₀₀ Arg Ala Gly Pro Met₂₀₅ Pro Gly Ser
 Ser Tyr₂₁₀ Gln Gly Thr Trp Ser₂₁₅ Glu Trp Ser Asp Pro₂₂₀ Val Ile Phe Gln
 Thr Gln Ser Glu Glu Leu₂₃₀ Lys Glu Gly Trp Asn₂₃₅ Gly Ser Gly Ser Arg₂₄₀
 Asp Lys Thr His Thr₂₄₅ Cys Pro Pro Cys Pro₂₅₀ Ala Pro Glu Leu Leu₂₅₅ Gly
 Gly Pro Ser Val₂₆₀ Phe Leu Phe Pro Pro₂₆₅ Lys Pro Lys Asp Thr₂₇₀ Leu Met
 Ile Ser Arg₂₇₅ Thr Pro Glu Val Thr₂₈₀ Cys Val Val Val Asp₂₈₅ Val Ser His

Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val
 290 295 300
 His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr
 305 310 315 320
 Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly
 325 330 335
 Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Val Pro Ile
 340 345 350
 Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val
 355 360 365
 Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser
 370 375 380
 Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu
 385 390 395 400
 Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro
 405 410 415
 Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val
 420 425 430
 Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met
 435 440 445
 His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser
 450 455 460
 Pro Gly Lys Ser Gly Met Ala Ser Met Thr Gly Gly Gln Gln Met Gly
 465 470 475 480
 Ser Gly Ser His His His His His Ser Gly Gly
 485 490

<210> 16
 <211> 1426
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (15)..(1415)

Ser	Gly	Ser	Arg 240	Asp	Lys	Thr	His	Thr 245	Cys	Pro	Pro	Cys	Pro 250	Ala	Pro		
gaa Glu	gcc Ala	ctg Leu 255	ggg Gly	gca Ala	ccg Pro	tca Ser	gtc Val 260	ttc Phe	ctc Leu	ttc Phe	ccc Pro	cca Pro 265	aaa Lys	ccc Pro	aag Lys	818	
gac Asp	acc Thr 270	ctc Leu	atg Met	atc Ile	tcc Ser	cgg Arg 275	acc Thr	cct Pro	gag Glu	gtc Val	aca Thr 280	tgc Cys	gtg Val	gtg Val	gtg Val	866	
gac Asp 285	gtg Val	agc Ser	cac His	gaa Glu	gac Asp 290	cct Pro	gag Glu	gtc Val	aag Lys	ttc Phe 295	aac Asn	tgg Trp	tac Tyr	gtg Val	gac Asp 300	914	
ggc Gly	gtg Val	gag Glu	gtg Val	cat His 305	aat Asn	gcc Ala	aag Lys	aca Thr	aag Lys 310	ccg Pro	cgg Arg	gag Glu	gag Glu	cag Gln 315	tac Tyr	962	
aac Asn	agc Ser	acg Thr	tac Tyr 320	cgt Arg	gtg Val	gtc Val	agc Ser	gtc Val 325	ctc Leu	acc Thr	gtc Val	ctg Leu	cac His 330	cag Gln	gac Asp	1010	
tgg Trp	ctg Leu	aat Asn 335	ggc Gly	aag Lys	gag Glu	tac Tyr	aag Lys 340	tgc Cys	aag Lys	gtc Val	tcc Ser	aac Asn 345	aaa Lys	gcc Ala	ctc Leu	1058	
cca Pro	gcc Ala 350	ccc Pro	atc Ile	gag Glu	aaa Lys	acc Thr 355	atc Ile	tcc Ser	aaa Lys	gcc Ala	aaa Lys 360	ggg Gly	cag Gln	ccc Pro	cga Arg	1106	
gaa Glu 365	cca Pro	cag Gln	gtg Val	tac Tyr	acc Thr 370	ctg Leu	ccc Pro	cca Pro	tcc Ser	cgg Arg 375	gag Glu	gag Glu	atg Met	acc Thr	aag Lys 380	1154	
aac Asn	cag Gln	gtc Val	agc Ser	ctg Leu 385	acc Thr	tgc Cys	ctg Leu	gtc Val	aaa Lys 390	ggc Gly	ttc Phe	tat Tyr	ccc Pro	agc Ser 395	gac Asp	1202	
atc Ile	gcc Ala	gtg Val	gag Glu 400	tgg Trp	gag Glu	agc Ser	aat Asn	ggg Gly 405	cag Gln	ccg Pro	gag Glu	aac Asn	aac Asn 410	tac Tyr	aag Lys	1250	
acc Thr	acg Thr	cct Pro 415	ccc Pro	gtg Val	ctg Leu	gac Asp	tcc Ser 420	gac Asp	ggc Gly	tcc Ser	ttc Phe	ttc Phe 425	ctc Leu	tat Tyr	agc Ser	1298	
aag Lys	ctc Leu 430	acc Thr	gtg Val	gac Asp	aag Lys	agc Ser 435	agg Arg	tgg Trp	cag Gln	cag Gln	ggg Gly 440	aac Asn	gtc Val	ttc Phe	tca Ser	1346	
tgc Cys 445	tcc Ser	gtg Val	atg Met	cat His	gag Glu 450	gct Ala	ctg Leu	cac His	aac Asn	cac His 455	tac Tyr	acg Thr	cag Gln	aag Lys	agc Ser 460	1394	
ctc Leu	tcc Ser	ctg Leu	tcc Ser	ccg Pro	ggc Gly 465	aaa Lys	tgagtgaatt c									1426	

<210> 17
 <211> 467
 <212> PRT

<213> Homo sapiens

<400> 17

Met Pro Arg Gly Trp Ala Ala Pro Leu Leu Leu Leu Leu Gln Gly
 1 5 10 15
 Gly Trp Gly Cys Pro Asp Leu Val Cys Tyr Thr Asp Tyr Leu Gln Thr
 20 25 30
 Val Ile Cys Ile Leu Glu Met Trp Asn Leu His Pro Ser Thr Leu Thr
 35 40 45
 Leu Thr Trp Gln Asp Gln Tyr Glu Glu Leu Lys Asp Glu Ala Thr Ser
 50 55 60
 Cys Ser Leu His Arg Ser Ala His Asn Ala Thr His Ala Thr Tyr Thr
 65 70 75 80
 Cys His Met Asp Val Phe His Phe Met Ala Asp Asp Ile Phe Ser Val
 85 90
 Asn Ile Thr Asp Gln Ser Gly Asn Tyr Ser Gln Glu Cys Gly Ser Phe
 100 105 110
 Leu Leu Ala Glu Ser Ile Lys Pro Ala Pro Pro Phe Asn Val Thr Val
 115 120 125
 Thr Phe Ser Gly Gln Tyr Asn Ile Ser Trp Arg Ser Asp Tyr Glu Asp
 130 135 140
 Pro Ala Phe Tyr Met Leu Lys Gly Lys Leu Gln Tyr Glu Leu Gln Tyr
 145 150 155 160
 Arg Asn Arg Gly Asp Pro Trp Ala Val Ser Pro Arg Arg Lys Leu Ile
 165 170 175
 Ser Val Asp Ser Arg Ser Val Ser Leu Leu Pro Leu Glu Phe Arg Lys
 180 185 190
 Asp Ser Ser Tyr Glu Leu Gln Val Arg Ala Gly Pro Met Pro Gly Ser
 195 200 205
 Ser Tyr Gln Gly Thr Trp Ser Glu Trp Ser Asp Pro Val Ile Phe Gln
 210 215 220
 Thr Gln Ser Glu Glu Leu Lys Glu Gly Trp Asn Gly Ser Gly Ser Arg
 225 230 235 240

$\langle 210 \rangle$	18
$\langle 211 \rangle$	741

<212> DNA
<213> Homo sapiens

<220>
<221> CDS
<222> (1)..(735)

<400> 18
atg ccg cgt ggc tgg gcc gcc ccc ttg ctc ctg ctg ctg ctc cag gga 48
Met Pro Arg Gly Trp Ala Ala Pro Leu Leu Leu Leu Leu Gln Gly
1 5 10 15

ggc tgg ggc tgc ccc gac ctc gtc tgc tac acc gat tac ctc cag acg 96
Gly Trp Gly Cys Pro Asp Leu Val Cys Tyr Thr Asp Tyr Leu Gln Thr
20 25 30

gtc atc tgc atc ctg gaa atg tgg aac ctc cac ccc agc acg ctc acc 144
Val Ile Cys Ile Leu Glu Met Trp Asn Leu His Pro Ser Thr Leu Thr
35 40 45

ctt acc tgg caa gac cag tat gaa gag ctg aag gac gag gcc acc tcc 192
Leu Thr Trp Gln Asp Gln Tyr Glu Glu Leu Lys Asp Glu Ala Thr Ser
50 55 60

tgc agc ctc cac agg tcg gcc cac aat gcc acg cat gcc acc tac acc 240
Cys Ser Leu His Arg Ser Ala His Asn Ala Thr His Ala Thr Tyr Thr
65 70 75 80

tgc cac atg gat gta ttc cac ttc atg gcc gac gac att ttc agt gtc 288
Cys His Met Asp Val Phe His Phe Met Ala Asp Asp Ile Phe Ser Val
85 90 95

aac atc aca gac cag tct ggc aac tac tcc cag gag tgt ggc agc ttt 336
Asn Ile Thr Asp Gln Ser Gly Asn Tyr Ser Ser Gln Glu Cys Gly Ser Phe
100 105 110

ctc ctg gct gag agc atc aag ccg gct ccc cct ttc aac gtg act gtg 384
Leu Leu Ala Glu Ser Ile Lys Pro Ala Pro Pro Phe Asn Val Thr Val
115 120 125

acc ttc tca gga cag tat aat atc tcc tgg cgc tca gat tac gaa gac 432
Thr Phe Ser Gly Gln Tyr Asn Ile Ser Trp Arg Ser Asp Tyr Glu Asp
130 135 140

cct gcc ttc tac atg ctg aag ggc aag ctt cag tat gag ctg cag tac 480
Pro Ala Phe Tyr Met Leu Lys Gly Lys Leu Gln Tyr Glu Leu Gln Tyr
145 150 155 160

agg aac cgg gga gac ccc tgg gct gtg agt ccg agg aga aag ctg atc 528
Arg Asn Arg Gly Asp Pro Trp Ala Val Ser Pro Arg Arg Lys Leu Ile
165 170 175

tca gtg gac tca aga agt gtc tcc ctc ctc ccc ctg gag ttc cgc aaa 576
Ser Val Asp Ser Arg Ser Val Ser Leu Leu Pro Leu Glu Phe Arg Lys
180 185 190

gac tcg agc tat gag ctg cag gtg cgg gca ggg ccc atg cct ggc tcc 624
Asp Ser Ser Tyr Glu Leu Gln Val Arg Ala Gly Pro Met Pro Gly Ser
195 200 205

tcc tac cag ggg acc tgg agt gaa tgg agt gac ccg gtc atc ttt cag 672
Ser Tyr Gln Gly Thr Trp Ser Glu Trp Ser Asp Pro Val Ile Phe Gln

210	215	220	
acc cag tca gag gag tta aag gaa ggc tgg aac aaa acc gaa acc tcc			720
Thr Gln Ser Glu Glu Leu Lys Glu Gly Trp Asn Lys Thr Glu Thr Ser			
225	230	235	240
cag gtt gct ccg gca taatga			741
Gln Val Ala Pro Ala			
	245		

<210> 19
 <211> 245
 <212> PRT
 <213> Homo sapiens

<400> 19

Met Pro Arg Gly Trp Ala Ala Pro Leu Leu Leu Leu Leu Gln Gly
1 5 10 15

Gly Trp Gly Cys Pro Asp Leu Val Cys Tyr Thr Asp Tyr Leu Gln Thr
20 25 30

Val Ile Cys Ile Leu Glu Met Trp Asn Leu His Pro Ser Thr Leu Thr
35 40 45

Leu Thr Trp Gln Asp Gln Tyr Glu Glu Leu Lys Asp Glu Ala Thr Ser
50 55 60

Cys Ser Leu His Arg Ser Ala His Asn Ala Thr His Ala Thr Tyr Thr
65 70 75 80

Cys His Met Asp Val Phe His Phe Met Ala Asp Asp Ile Phe Ser Val
85 90 95

Asn Ile Thr Asp Gln Ser Gly Asn Tyr Ser Gln Glu Cys Gly Ser Phe
100 105 110

Leu Leu Ala Glu Ser Ile Lys Pro Ala Pro Pro Phe Asn Val Thr Val
115 120 125

Thr Phe Ser Gly Gln Tyr Asn Ile Ser Trp Arg Ser Asp Tyr Glu Asp
130 135 140

Pro Ala Phe Tyr Met Leu Lys Gly Lys Leu Gln Tyr Glu Leu Gln Tyr
145 150 155 160

Arg Asn Arg Gly Asp Pro Trp Ala Val Ser Pro Arg Arg Lys Leu Ile
165 170 175

Ser Val Asp Ser Arg Ser Val Ser Leu Leu Pro Leu Glu Phe Arg Lys

180 185 190
 Asp Ser Ser Tyr Glu Leu Gln Val Arg Ala Gly Pro Met Pro Gly Ser
 195 200 205
 Ser Tyr Gln Gly Thr Trp Ser Glu Trp Ser Asp Pro Val Ile Phe Gln
 210 215 220
 Thr Gln Ser Glu Glu Leu Lys Glu Gly Trp Asn Lys Thr Glu Thr Ser
 225 230 235 240
 Gln Val Ala Pro Ala
 245

<210> 20
 <211> 1413
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)..(1410)

<400> 20
 atg ccg cgt ggc tgg gcc gcc ccc ttg ctc ctg ctg ctg ctc cag gga 48
 Met Pro Arg Gly Trp Ala Ala Pro Leu Leu Leu Leu Leu Gln Gly
 1 5 10 15
 ggc tgg ggc tgc ccc gac ctc gtc tgc tac acc gat tac ctc cag acg 96
 Gly Trp Gly Cys Pro Asp Leu Val Cys Tyr Thr Asp Tyr Leu Gln Thr
 20 25 30
 gtc atc tgc atc ctg gaa atg tgg aac ctc cac ccc agc acg ctc acc 144
 Val Ile Cys Ile Leu Glu Met Trp Asn Leu His Pro Ser Thr Leu Thr
 35 40 45
 ctt acc tgg caa gac cag tat gaa gag ctg aag gac gag gcc acc tcc 192
 Leu Thr Trp Gln Asp Gln Tyr Glu Glu Leu Lys Asp Glu Ala Thr Ser
 50 55 60
 tgc agc ctc cac agg tgc gcc cac aat gcc acg cat gcc acc tac acc 240
 Cys Ser Leu His Arg Ser Ala His Asn Ala Thr His Ala Thr Tyr Thr
 65 70 75 80
 tgc cac atg gat gta ttc cac ttc atg gcc gac gac att ttc agt gtc 288
 Cys His Met Asp Val Phe His Phe Met Ala Asp Asp Ile Phe Ser Val
 85 90 95
 aac atc aca gac cag tct ggc aac tac tcc cag gag tgt ggc agc ttt 336
 Asn Ile Thr Asp Gln Ser Gly Asn Tyr Ser Gln Glu Cys Gly Ser Phe
 100 105 110
 ctc ctg gct gag agc atc aag ccg gct ccc cct ttc aac gtg act gtg 384
 Leu Leu Ala Glu Ser Ile Lys Pro Ala Pro Pro Phe Asn Val Thr Val
 115 120 125
 acc ttc tca gga cag tat aat atc tcc tgg cgc tca gat tac gaa gac 432

Thr 130	Phe	Ser	Gly	Gln	Tyr	Asn 135	Ile	Ser	Trp	Arg	Ser 140	Asp	Tyr	Glu	Asp	
cct Pro 145	gcc Ala	ttc Phe	tac Tyr	atg Met	ctg Leu 150	aag Lys	ggc Gly	aag Lys	ctt Leu	cag Gln 155	tat Tyr	gag Glu	ctg Leu	cag Gln	tac Tyr 160	480
agg Arg	aac Asn	cgg Arg	gga Gly	gac Asp 165	ccc Pro	tgg Trp	gct Ala	gtg Val	agt Ser 170	ccg Pro	agg Arg	aga Arg	aag Lys	ctg Leu 175	atc Ile	528
tca Ser	gtg Val	gac Asp	tca Ser 180	aga Arg	agt Ser	gtc Val	tcc Ser	ctc Leu 185	ctc Leu	ccc Pro	ctg Leu	gag Glu	ttc Phe 190	cgc Arg	aaa Lys	576
gac Asp	tcg Ser	agc Ser 195	tat Tyr	gag Glu	ctg Leu	cag Gln	gtg Val 200	cgg Arg	gca Ala	ggg Gly	ccc Pro	atg Met 205	cct Pro	ggc Gly	tcc Ser	624
tcc Ser	tac Tyr 210	cag Gln	ggg Gly	acc Thr	tgg Trp	agt Ser 215	gaa Glu	tgg Trp	agt Ser	gac Asp	ccg Pro 220	gtc Val	atc Ile	ttt Phe	cag Gln	672
acc Thr 225	cag Gln	tca Ser	gag Glu	gag Glu	tta Leu 230	aag Lys	gaa Glu	ggc Gly	tgg Trp	aac Asn 235	gat Asp	gac Asp	gat Asp	gac Asp	aag Lys 240	720
ggc Gly	tcc Ser	ggc Gly	gac Asp	aaa Lys 245	act Thr	cac His	aca Thr	tgc Cys	cca Pro 250	ccg Pro	tgc Cys	cca Pro	gca Ala	cct Pro 255	gaa Glu	768
gcc Ala	ctg Leu	ggg Gly	gca Ala 260	ccg Pro	tca Ser	gtc Val	ttc Phe	ctc Leu 265	ttc Phe	ccc Pro	cca Pro	aaa Lys	ccc Pro 270	aag Lys	gac Asp	816
acc Thr	ctc Leu	atg Met 275	atc Ile	tcc Ser	cgg Arg	acc Thr	cct Pro 280	gag Glu	gtc Val	aca Thr	tgc Cys	gtg Val 285	gtg Val	gtg Val	gac Asp	864
gtg Val 290	agc Ser	cac His	gaa Glu	gac Asp	cct Pro	gag Glu 295	gtc Val	aag Lys	ttc Phe	aac Asn	tgg Trp 300	tac Tyr	gtg Val	gac Asp	ggc Gly	912
gtg Val 305	gag Glu	gtg Val	cat His	aat Asn	gcc Ala 310	aag Lys	aca Thr	aag Lys	ccg Pro	cgg Arg 315	gag Glu	gag Glu	cag Gln	tac Tyr	aac Asn 320	960
agc Ser	acg Thr	tac Tyr	cgt Arg	gtg Val 325	gtc Val	agc Ser	gtc Val	ctc Leu	acc Thr 330	gtc Val	ctg Leu	cac His	cag Gln	gac Asp 335	tgg Trp	1008
ctg Leu	aat Asn	ggc Gly	aag Lys 340	gag Glu	tac Tyr	aag Lys	tgc Cys	aag Lys 345	gtc Val	tcc Ser	aac Asn	aaa Lys	gcc Ala 350	ctc Leu	cca Pro	1056
gcc Ala	ccc Pro	atc Ile 355	gag Glu	aaa Lys	acc Thr	atc Ile	tcc Ser 360	aaa Lys	gcc Ala	aaa Lys	ggg Gly	cag Gln 365	ccc Pro	cga Arg	gaa Glu	1104
cca Pro 370	cag Gln	gtg Val	tac Tyr	acc Thr	ctg Leu	ccc Pro 375	cca Pro	tcc Ser	cgg Arg	gag Glu	gag Glu 380	atg Met	acc Thr	aag Lys	aac Asn	1152

cag Gln 385	gtc Val	agc Ser	ctg Leu	acc Thr	tgc Cys 390	ctg Leu	gtc Val	aaa Lys	ggc Gly 395	ttc Phe 395	tat Tyr	ccc Pro	agc Ser	gac Asp	atc Ile 400	1200
gcc Ala	gtg Val	gag Glu	tgg Trp	gag Glu 405	agc Ser	aat Asn	ggg Gly	cag Gln 410	ccg Pro	gag Glu	aac Asn	aac Asn	tac Tyr	aag Lys 415	acc Thr	1248
acg Thr	cct Pro	ccc Pro	gtg Val 420	ctg Leu	gac Asp	tcc Ser	gac Asp	ggc Gly 425	tcc Ser	ttc Phe	ttc Phe	ctc Leu	tat Tyr 430	agc Ser	aag Lys	1296
ctc Leu	acc Thr	gtg Val 435	gac Asp	aag Lys	agc Ser	agg Arg	tgg Trp 440	cag Gln	cag Gln	ggg Gly	aac Asn	gtc Val 445	ttc Phe	tca Ser	tgc Cys	1344
tcc Ser	gtg Val 450	atg Met	cat His	gag Glu	gct Ala	ctg Leu 455	cac His	aac Asn	cac His	tac Tyr	acg Thr 460	cag Gln	aag Lys	agc Ser	ctc Leu	1392
tcc Ser 465	ctg Leu	tcc Ser	ccg Pro	ggt Gly	aaa Lys 470	tga										1413

<210> 21
 <211> 470
 <212> PRT
 <213> Homo sapiens

<400> 21

Met 1	Pro	Arg	Gly	Trp 5	Ala	Ala	Pro	Leu	Leu 10	Leu	Leu	Leu	Leu	Gln 15	Gly
Gly	Trp	Gly	Cys 20	Pro	Asp	Leu	Val	Cys 25	Tyr	Thr	Asp	Tyr	Leu 30	Gln	Thr
Val	Ile	Cys 35	Ile	Leu	Glu	Met	Trp 40	Asn	Leu	His	Pro	Ser 45	Thr	Leu	Thr
Leu	Thr	Trp	Gln	Asp	Gln	Tyr 55	Glu	Glu	Leu	Lys	Asp 60	Glu	Ala	Thr	Ser
Cys 65	Ser	Leu	His	Arg	Ser 70	Ala	His	Asn	Ala	Thr 75	His	Ala	Thr	Tyr	Thr 80
Cys	His	Met	Asp	Val 85	Phe	His	Phe	Met	Ala 90	Asp	Asp	Ile	Phe	Ser	Val 95
Asn	Ile	Thr	Asp 100	Gln	Ser	Gly	Asn	Tyr 105	Ser	Gln	Glu	Cys	Gly 110	Ser	Phe
Leu	Leu	Ala 115	Glu	Ser	Ile	Lys	Pro 120	Ala	Pro	Pro	Phe	Asn 125	Val	Thr	Val

Thr Phe Ser Gly Gln Tyr Asn Ile Ser Trp Arg Ser Asp Tyr Glu Asp
 130 135 140
 Pro Ala Phe Tyr Met Leu Lys Gly Lys Leu Gln Tyr Glu Leu Gln Tyr
 145 150 155 160
 Arg Asn Arg Gly Asp Pro Trp Ala Val Ser Pro Arg Arg Lys Leu Ile
 165 170 175
 Ser Val Asp Ser Arg Ser Val Ser Leu Leu Pro Leu Glu Phe Arg Lys
 180 185 190
 Asp Ser Ser Tyr Glu Leu Gln Val Arg Ala Gly Pro Met Pro Gly Ser
 195 200 205
 Ser Tyr Gln Gly Thr Trp Ser Glu Trp Ser Asp Pro Val Ile Phe Gln
 210 215 220
 Thr Gln Ser Glu Glu Leu Lys Glu Gly Trp Asn Asp Asp Asp Asp Lys
 225 230 235 240
 Gly Ser Gly Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu
 245 250 255
 Ala Leu Gly Ala Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp
 260 265 270
 Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp
 275 280 285
 Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly
 290 295 300
 Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn
 305 310 315 320
 Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp
 325 330 335
 Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro
 340 345 350
 Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu
 355 360 365
 Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn
 370 375 380

Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile
385 390 395 400

Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr
405 410 415

Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys
420 425 430

Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys
435 440 445

Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu
450 455 460

Ser Leu Ser Pro Gly Lys
465 470

<210> 22
<211> 1754
<212> DNA
<213> Mus musculus

<220>
<221> CDS
<222> (1)..(720)

<400> 22	
atg ccc cgg ggc cca gtg gct gcc tta ctc ctg ctg att ctc cat gga	48
Met Pro Arg Gly Pro Val Ala Ala Leu Leu Leu Leu Ile Leu His Gly	
1 5 10 15	
gct tgg agc tgc ctg gac ctc act tgc tac act gac tac ctc tgg acc	96
Ala Trp Ser Cys Leu Asp Leu Thr Cys Tyr Thr Asp Tyr Leu Trp Thr	
20 25 30	
atc acc tgt gtc ctg gag aca cgg agc ccc aac ccc agc ata ctc agt	144
Ile Thr Cys Val Leu Glu Thr Arg Ser Pro Asn Pro Ser Ile Leu Ser	
35 40 45	
ctc acc tgg caa gat gaa tat gag gaa ctt cag gac caa gag acc ttc	192
Leu Thr Trp Gln Asp Glu Tyr Glu Glu Leu Gln Asp Gln Glu Thr Phe	
50 55 60	
tgc agc cta cac agg tct ggc cac aac acc aca cat ata tgg tac acg	240
Cys Ser Leu His Arg Ser Gly His Asn Thr Thr His Ile Trp Tyr Thr	
65 70 75 80	
tgc cat atg cgc ttg tct caa ttc ctg tcc gat gaa gtt ttc att gtc	288
Cys His Met Arg Leu Ser Gln Phe Leu Ser Asp Glu Val Phe Ile Val	
85 90 95	
aat gtg acg gac cag tct ggc aac aac tcc caa gag tgt ggc agc ttt	336
Asn Val Thr Asp Gln Ser Gly Asn Asn Ser Gln Glu Cys Gly Ser Phe	

100	105	110	
gtc ctg gct gag agc atc aaa cca gct ccc ccc ttg aac gtg act gtg Val Leu Ala Glu Ser Ile Lys Pro Ala Pro Pro Leu Asn Val Thr Val 115 120 125			384
gcc ttc tca gga cgc tat gat atc tcc tgg gac tca gct tat gac gaa Ala Phe Ser Gly Arg Tyr Asp Ile Ser Trp Asp Ser Ala Tyr Asp Glu 130 135 140			432
ccc tcc aac tac gtg ctg agg ggc aag cta caa tat gag ctg cag tat Pro Ser Asn Tyr Val Leu Arg Gly Lys Leu Gln Tyr Glu Leu Gln Tyr 145 150 155 160			480
cgg aac ctc aga gac ccc tat gct gtg agg ccg gtg acc aag ctg atc Arg Asn Leu Arg Asp Pro Tyr Ala Val Arg Pro Val Thr Lys Leu Ile 165 170 175			528
tca gtg gac tca aga aac gtc tct ctt ctc cct gaa gag ttc cac aaa Ser Val Asp Ser Arg Asn Val Ser Leu Leu Pro Glu Glu Phe His Lys 180 185 190			576
gat tct agc tac cag ctg cag gtg cgg gca gcg cct cag cca ggc act Asp Ser Ser Tyr Gln Leu Gln Val Arg Ala Ala Pro Gln Pro Gly Thr 195 200 205			624
tca ttc agg ggg acc tgg agt gag tgg agt gac ccc gtc atc ttt cag Ser Phe Arg Gly Thr Trp Ser Glu Trp Ser Asp Pro Val Ile Phe Gln 210 215 220			672
acc cag gct ggg gag ccc gag gca ggc tgg gac ggc tcc ggc tct aga Thr Gln Ala Gly Glu Pro Glu Ala Gly Trp Asp Gly Ser Gly Ser Arg 225 230 235 240			720
gagccccgcg gaccgacaat caagccctgt cctccatgca aatgcccagg taagtcacta			780
gaccagagct ccactcccgg gagaatggta agtgctataa acatccctgc actagaggat			840
aagccatgta cagatccatt tccatctctc ctcatcagca cctaacctcg aggggtggacc			900
atccgtcttc atcttccctc caaagatcaa ggatgtactc atgatctccc tgagcccat			960
agtcacatgt gtggtggtgg atgtgagcga ggatgacca gatgtccaga tcagctggtt			1020
tgtgaacaac gtggaagtac acacagctca gacacaaacc catagagagg attacaacag			1080
tactctccgg gtggtcagtg ccctcccat ccagcaccag gactggatga gtggcaaggc			1140
tttcgcatgc gccgtcaaca acaaagacct ccagcgcgc atcgagagaa ccatctcaaa			1200
acccaaaggt gagagctgca gcctgactgc atgggggctg ggatgggcat aaggataaag			1260
gtctgtgtgg acagccttct gcttcagcca tgacctttgt gtatgtttct accctcacag			1320
ggtcagtaag agctccacag gtatatgtct tgcctccacc agaagaagag atgactaaga			1380
aacaggtcac tctgacctgc atggtcacag acttcatgcc tgaagacatt tacgtggagt			1440
ggaccaâcaa cgggaaaaca gagctaaact acaagaacac tgaaccagtc ctggactctg			1500
atggttctta cttcatgtac agcaagctga gagtggaaaa gaagaactgg gtggaaagaa			1560
atagctactc ctgttcagtg gtccacgagg gtctgcacaa tcaccacacg actaagagct			1620

tctcccggac tccgggtaaa tgagctcagc acccacaataa ctctcagggtc caaagagaca 1680
 cccacactca tctccatgct tcccttgtat aaataaagca cccagcaatg cctggggacca 1740
 tgtaatagga attc 1754

<210> 23
 <211> 240
 <212> PRT
 <213> Mus musculus
 <400> 23

Met Pro Arg Gly Pro Val Ala Ala Leu Leu Leu Leu Ile Leu His Gly
 1 5 10 15
 Ala Trp Ser Cys Leu Asp Leu Thr Cys Tyr Thr Asp Tyr Leu Trp Thr
 20 25 30
 Ile Thr Cys Val Leu Glu Thr Arg Ser Pro Asn Pro Ser Ile Leu Ser
 35 40 45
 Leu Thr Trp Gln Asp Glu Tyr Glu Glu Leu Gln Asp Gln Glu Thr Phe
 50 55 60
 Cys Ser Leu His Arg Ser Gly His Asn Thr Thr His Ile Trp Tyr Thr
 65 70 75 80
 Cys His Met Arg Leu Ser Gln Phe Leu Ser Asp Glu Val Phe Ile Val
 85 90 95
 Asn Val Thr Asp Gln Ser Gly Asn Asn Ser Gln Glu Cys Gly Ser Phe
 100 105 110
 Val Leu Ala Glu Ser Ile Lys Pro Ala Pro Pro Leu Asn Val Thr Val
 115 120 125
 Ala Phe Ser Gly Arg Tyr Asp Ile Ser Trp Asp Ser Ala Tyr Asp Glu
 130 135 140
 Pro Ser Asn Tyr Val Leu Arg Gly Lys Leu Gln Tyr Glu Leu Gln Tyr
 145 150 155 160
 Arg Asn Leu Arg Asp Pro Tyr Ala Val Arg Pro Val Thr Lys Leu Ile
 165 170 175
 Ser Val Asp Ser Arg Asn Val Ser Leu Leu Pro Glu Glu Phe His Lys
 180 185 190

Asp Ser Ser Tyr Gln Leu Gln Val Arg Ala Ala Pro Gln Pro Gly Thr
 195 200 205

Ser Phe Arg Gly Thr Trp Ser Glu Trp Ser Asp Pro Val Ile Phe Gln
 210 215 220

Thr Gln Ala Gly Glu Pro Glu Ala Gly Trp Asp Gly Ser Gly Ser Arg
 225 230 235 240

<210> 24
 <211> 795
 <212> DNA
 <213> Mus musculus

<220>
 <221> CDS
 <222> (19)..(783)

<400> 24
 ctgcaggctcg acaccacc atg ccc cgg ggc cca gtg gct gcc tta ctc ctg 51
 Met Pro Arg Gly Pro Val Ala Ala Leu Leu 10
 1
 ctg att ctc cat gga gct tgg agc tgc ctg gac ctc act tgc tac act 99
 Leu Ile Leu His Gly Ala Trp Ser Cys Leu Asp Leu Thr Cys Tyr Thr 20 25
 15
 gac tac ctc tgg acc atc acc tgt gtc ctg gag aca cgg agc ccc aac 147
 Asp Tyr Leu Trp Thr Ile Thr Cys Val Leu Glu Thr Arg Ser Pro Asn 30 35 40
 ccc agc ata ctc agt ctc acc tgg caa gat gaa tat gag gaa ctt cag 195
 Pro Ser Ile Leu Ser Leu Thr Trp Gln Asp Glu Tyr Glu Glu Leu Gln 45 50 55
 gac caa gag acc ttc tgc agc cta cac agg tct ggc cac aac acc aca 243
 Asp Gln Glu Thr Phe Cys Ser Leu His Arg Ser Gly His Asn Thr Thr 60 65 70 75
 cat ata tgg tac acg tgc cat atg cgc ttg tct caa ttc ctg tcc gat 291
 His Ile Trp Tyr Thr Cys His Met Arg Leu Ser Gln Phe Leu Ser Asp 80 85 90
 gaa gtt ttc att gtc aat gtg acg gac cag tct ggc aac aac tcc caa 339
 Glu Val Phe Ile Val Asn Val Thr Asp Gln Ser Gly Asn Asn Ser Gln 95 100 105
 gag tgt ggc agc ttt gtc ctg gct gag agc atc aaa cca gct ccc ccc 387
 Glu Cys Gly Ser Phe Val Leu Ala Glu Ser Ile Lys Pro Ala Pro Pro 110 115 120
 ttg aac gtg act gtg gcc ttc tca gga cgc tat gat atc tcc tgg gac 435
 Leu Asn Val Thr Val Ala Phe Ser Gly Arg Tyr Asp Ile Ser Trp Asp 125 130 135
 tca gct tat gac gaa ccc tcc aac tac gtg ctg agg ggc aag cta caa 483
 Ser Ala Tyr Asp Glu Pro Ser Asn Tyr Val Leu Arg Gly Lys Leu Gln 140 145 150 155

tat gag ctg cag tat cgg aac ctc aga gac ccc tat gct gtg agg ccg 531
 Tyr Glu Leu Gln Tyr Arg Asn Leu Arg Asp Pro Tyr Ala Val Arg Pro
 160 165 170
 gtg acc aag ctg atc tca gtg gac tca aga aac gtc tct ctt ctc cct 579
 Val Thr Lys Leu Ile Ser Val Asp Ser Arg Asn Val Ser Leu Leu Pro
 175 180 185
 gaa gag ttc cac aaa gat tct agc tac cag ctg cag gtg cgg gca gcg 627
 Glu Glu Phe His Lys Asp Ser Ser Tyr Gln Leu Gln Val Arg Ala Ala
 190 195 200
 cct cag cca ggc act tca ttc agg ggg acc tgg agt gag tgg agt gac 675
 Pro Gln Pro Gly Thr Ser Phe Arg Gly Thr Trp Ser Glu Trp Ser Asp
 205 210 215
 ccc gtc atc ttt cag acc cag gct ggg gag ccc gag gca ggc tgg gac 723
 Pro Val Ile Phe Gln Thr Gln Ala Gly Glu Pro Glu Ala Gly Trp Asp
 220 225 230 235
 ggc agc gga cac cac cat cat cac cac ggt agc ggc gac tat aaa gac 771
 Gly Ser Gly His His His His His Gly Ser Gly Asp Tyr Lys Asp
 240 245 250
 gat gac gat aag tagtgagaat tc 795
 Asp Asp Asp Lys
 255

<210> 25
 <211> 255
 <212> PRT
 <213> Mus musculus

<400> 25

Met Pro Arg Gly Pro Val Ala Ala Leu Leu Leu Ile Leu His Gly
 1 5 10 15
 Ala Trp Ser Cys Leu Asp Leu Thr Cys Tyr Thr Asp Tyr Leu Trp Thr
 20 25 30
 Ile Thr Cys Val Leu Glu Thr Arg Ser Pro Asn Pro Ser Ile Leu Ser
 35 40 45
 Leu Thr Trp Gln Asp Glu Tyr Glu Glu Leu Gln Asp Gln Glu Thr Phe
 50 55 60
 Cys Ser Leu His Arg Ser Gly His Asn Thr Thr His Ile Trp Tyr Thr
 65 70 75 80
 Cys His Met Arg Leu Ser Gln Phe Leu Ser Asp Glu Val Phe Ile Val
 85 90 95
 Asn Val Thr Asp Gln Ser Gly Asn Asn Ser Gln Glu Cys Gly Ser Phe
 100 105 110

Val Leu Ala Glu Ser Ile Lys Pro Ala Pro Pro Leu Asn Val Thr Val
 115 120 125
 Ala Phe Ser Gly Arg Tyr Asp Ile Ser Trp Asp Ser Ala Tyr Asp Glu
 130 135 140
 Pro Ser Asn Tyr Val Leu Arg Gly Lys Leu Gln Tyr Glu Leu Gln Tyr
 145 150 155 160
 Arg Asn Leu Arg Asp Pro Tyr Ala Val Arg Pro Val Thr Lys Leu Ile
 165 170 175
 Ser Val Asp Ser Arg Asn Val Ser Leu Leu Pro Glu Glu Phe His Lys
 180 185 190
 Asp Ser Ser Tyr Gln Leu Gln Val Arg Ala Ala Pro Gln Pro Gly Thr
 195 200 205
 Ser Phe Arg Gly Thr Trp Ser Glu Trp Ser Asp Pro Val Ile Phe Gln
 210 215 220
 Thr Gln Ala Gly Glu Pro Glu Ala Gly Trp Asp Gly Ser Gly His His
 225 230 235 240
 His His His His Gly Ser Gly Asp Tyr Lys Asp Asp Asp Asp Lys
 245 250 255

<210> 26
 <211> 792
 <212> DNA
 <213> Mus musculus

<220>
 <221> CDS
 <222> (1)..(786)

<400> 26
 atg aaa ttc tta gtc aac gtt gcc ctt gtt ttt atg gtc gtg tac att 48
 Met Lys Phe Leu Val Asn Val Ala Leu Val Phe Met Val Val Tyr Ile
 1 5 10 15
 tct tac atc tat gcc ggc agc gga cac cac cat cat cac cac ggt agc 96
 Ser Tyr Ile Tyr Ala Gly Ser Gly His His His His His His Gly Ser
 20 25 30
 ggc gac tat aaa gac gat gac gat aag ggt tcc gga tgc ctg gac ctc 144
 Gly Asp Tyr Lys Asp Asp Asp Asp Lys Gly Ser Gly Cys Leu Asp Leu
 35 40 45
 act tgc tac act gac tac ctc tgg acc atc acc tgt gtc ctg gag aca 192
 Thr Cys Tyr Thr Asp Tyr Leu Trp Thr Ile Thr Cys Val Leu Glu Thr
 50 55 60

cgg agc ccc aac ccc agc ata ctc agt ctc acc tgg caa gat gaa tat	240
Arg Ser Pro Asn Pro Ser Ile Leu Ser Leu Thr Trp Gln Asp Glu Tyr	
65 70 75 80	
gag gaa ctt cag gac caa gag acc ttc tgc agc cta cac agg tct ggc	288
Glu Glu Leu Gln Asp Gln Glu Thr Phe Cys Ser Leu His Arg Ser Gly	
85 90 95	
cac aac acc aca cat ata tgg tac acg tgc cat atg cgc ttg tct caa	336
His Asn Thr Thr His Ile Trp Tyr Thr Cys His Met Arg Leu Ser Gln	
100 105 110	
ttc ctg tcc gat gaa gtt ttc att gtc aat gtg acg gac cag tct ggc	384
Phe Leu Ser Asp Glu Val Phe Ile Val Asn Val Thr Asp Gln Ser Gly	
115 120 125	
aac aac tcc caa gag tgt ggc agc ttt gtc ctg gct gag agc atc aaa	432
Asn Asn Ser Gln Glu Cys Gly Ser Phe Val Leu Ala Glu Ser Ile Lys	
130 135 140	
cca gct ccc ccc ttg aac gtg act gtg gcc ttc tca gga cgc tat gat	480
Pro Ala Pro Pro Leu Asn Val Thr Val Ala Phe Ser Gly Arg Tyr Asp	
145 150 155 160	
atc tcc tgg gac tca gct tat gac gaa ccc tcc aac tac gtg ctg agg	528
Ile Ser Trp Asp Ser Ala Tyr Asp Glu Pro Ser Asn Tyr Val Leu Arg	
165 170 175	
ggc aag cta caa tat gag ctg cag tat cgg aac ctc aga gac ccc tat	576
Gly Lys Leu Gln Tyr Glu Leu Gln Tyr Arg Asn Leu Arg Asp Pro Tyr	
180 185 190	
gct gtg agg ccg gtg acc aag ctg atc tca gtg gac tca aga aac gtc	624
Ala Val Arg Pro Val Thr Lys Leu Ile Ser Val Asp Ser Arg Asn Val	
195 200 205	
tct ctt ctc cct gaa gag ttc cac aaa gat tct agc tac cag ctg cag	672
Ser Leu Leu Pro Glu Glu Phe His Lys Asp Ser Ser Tyr Gln Leu Gln	
210 215 220	
gtg cgg gca gcg cct cag cca ggc act tca ttc agg ggg acc tgg agt	720
Val Arg Ala Ala Pro Gln Pro Gly Thr Ser Phe Arg Gly Thr Trp Ser	
225 230 235 240	
gag tgg agt gac ccc gtc atc ttt cag acc cag gct ggg gag ccc gag	768
Glu Trp Ser Asp Pro Val Ile Phe Gln Thr Gln Ala Gly Glu Pro Glu	
245 250 255	
gca ggc tgg gac tag tga gaattc	792
Ala Gly Trp Asp	
260	

<210> 27
 <211> 260
 <212> PRT
 <213> Mus musculus

<400> 27

Met Lys Phe Leu Val Asn Val Ala Leu Val Phe Met Val Val Tyr Ile
1 5 10 15

260

<210> 28
<211> 20
<212> DNA
<213> Artificial

<220>
<223> PCR Primer

<400> 28
gccagatcgc ctcttgatta 20

<210> 29
<211> 20
<212> DNA
<213> Artificial

<220>
<223> PCR Primer

<400> 29
catgctcaca gtgccccttt 20

<210> 30
<211> 20
<212> DNA
<213> Artificial

<220>
<223> PCR Primer

<400> 30
ctccccctt gaacgtgact 20

<210> 31
<211> 20
<212> DNA
<213> Artificial

<220>
<223> PCR Primer

<400> 31
ttgcccctca gcacgtagtt 20

<210> 32
<211> 28
<212> DNA
<213> Artificial

<220>
<223> PCR Primer

<400> 32
agagccagat tatctctttc tacctcag 28

<210> 33
<211> 20
<212> DNA
<213> Artificial

<220>
<223> PCR Primer

<400> 33
cctttttcgc cttgctgttg

20

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2006/013829

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07K14/715 A61P17/02 A61K38/17
ADD. C07K16/28 C07K16/24 A61K39/395

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, EMBASE, Sequence Search, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 03/028630 A (WYETH; CARTER, LAURA; WHITTERS, MATTHEW, J; COLLINS, MARY; YOUNG, DEBO) 10 April 2003 (2003-04-10) pages 2-14 pages 52,53; sequences 6,10, 22-39	1-28
X	DISTLER JORG H W ET AL: "Expression of interleukin-21 receptor in epidermis from patients with systemic sclerosis" ARTHRITIS & RHEUMATISM, vol. 52, no. 3, March 2005 (2005-03), pages 856-864, XP009070828 ISSN: 0004-3591 the whole document ----- --/--	23-28

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

11 August 2006

Date of mailing of the international search report

21/08/2006

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Bernhardt, W

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2006/013829

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	US 2006/039902 A1 (YOUNG DEBORAH A ET AL) 23 February 2006 (2006-02-23) paragraphs [0007] - [0031], [0210] - [0215]; sequences 2,10,23-40 -----	1-28
A	MEHTA DEVANGI S ET AL: "Biology of IL-21 and the IL-21 receptor" IMMUNOLOGICAL REVIEWS, vol. 202, no. December, December 2004 (2004-12), pages 84-95, XP009070837 ISSN: 0105-2896 the whole document -----	1-28
T	PESCE JOHN ET AL: "The IL-21 receptor augments Th2 effector function and alternative macrophage activation." THE JOURNAL OF CLINICAL INVESTIGATION. JUL 2006, vol. 116, no. 7, July 2006 (2006-07), pages 2044-2055, XP009070829 ISSN: 0021-9738 the whole document -----	1-22
A	WO 03/087320 A (BETH ISRAEL DEACONESS MEDICAL CENTER, INC; MOLL, THOMAS; STROM, TERRY,) 23 October 2003 (2003-10-23) pages 2-5 pages 28-30 -----	1-28

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2006/013829

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box II.1

Although claims 1-22 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition. In addition, although claims 25-28 are directed to a diagnostic method practised on the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Continuation of Box II.1

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

Rule 39.1(iv) PCT - Diagnostic method practised on the human or animal body

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2006/013829

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 03028630	A	10-04-2003	CA 2460916 A1	10-04-2003
			EP 1432431 A2	30-06-2004
			JP 2005508915 T	07-04-2005
US 2006039902	A1	23-02-2006	NONE	
WO 03087320	A	23-10-2003	AU 2003230834 A1	27-10-2003